





# 3 SUPER 3-WAY WEEDILIZER

a fertilizer with weed & insect control  
for established dichondra lawns only

DIPHENAMID (IN N-DIMETHYL 2,2-DIPHENYLACETAMIDE)	2.30%
MONURON 3-(P-CHLOROPHENYL)-1,1-DIMETHYLUREA	0.25%
<b>INERT INGREDIENTS</b>	
FERTILIZER 21-6-6	96.30%
<b>GUARANTEED FERTILIZER ANALYSIS</b>	
ORGANIC NITROGEN (DERIVED FROM UREA AND UREA-FORMALDEHYDE)	18.60%
SOLUBLE NITROGEN	2.40%
AMMONIC NITROGEN (DERIVED FROM AMMONIUM PHOSPHATE)	2.40%
TOTAL NITROGEN	21.00%
PHOSPHORIC ACID (AVAILABLE)	6.00%
(DERIVED FROM AMMONIUM PHOSPHATE)	
POTASH (WATER-SOLUBLE) (DERIVED FROM MURIATE OF POTASH)	6.00%
CALCIUM (DERIVED FROM GYPSUM)	1.00%
SULFUR (COMBINED) (DERIVED FROM GYPSUM)	1.50%
IRON (EXPRESSED AS ELEMENTAL)	1.00%
MANGANESE (EXPRESSED AS ELEMENTAL)	0.10%
ZINC (EXPRESSED AS ELEMENTAL)	0.10%
*DERIVED FROM SPECIALLY PROCESSED NATURAL MINERALS CONTAINING A MIXTURE WHICH GRADUALLY BECOMES SOLUBLE UPON EXPOSURE TO AIR AND WATER	

**CAUTION: KEEP OUT OF REACH OF CHILDREN AND PETS**

HARMFUL IF INHALED OR SWALLOWED  
 AVOID SKIN CONTACT  
 WASH HANDS THOROUGHLY BEFORE EATING  
 IF SWALLOWED INDUCE VOMITING AND CALL PHYSICIAN IMMEDIATELY  
 CARBARYL IS A MODERATE REVERSIBLE CHOLINESTERASE INHIBITOR. ATROPINE IS ANTIDOTAL  
 DO NOT CONTAMINATE FEED AND FOODSTUFFS. LAWN CLIPPINGS AFTER APPLICATION SHOULD NOT BE FED TO LIVESTOCK OR POULTRY  
 DO NOT USE ON BLADE GRASS LAWNS OR ON EDIBLE CROPS  
 CONTAINER DISPOSAL: DO NOT REUSE EMPTY CONTAINER. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

THIS PRODUCT IS IN A CLEAN, EASY TO HANDLE PELLET FORM AND FURNISHES A BALANCED FERTILIZER ESPECIALLY DEVELOPED FOR OUR ACID-NEEDING SOILS OF THE WEST.

- PREVENTS CRABGRASS, SPOTTED SPURGE AND OTHER WEEDS
  - CONTROLS OXALIS AND ANNUAL BLUEGRASS
  - RETARDS BERMUDA (DEVIL) GRASS
  - STOPS DAMAGING LAWN INSECTS
  - FEEDS DICONDRA LAWNS WITH COMPLETE FERTILIZER
  - CONDITIONS SOIL AND AIDS IN REPAIRING ALCALINITY
- THIS PRODUCT CONTROLS THE GERMINATING SEED AND SEEDLING STAGES OF THE FOLLOWING WEEDS:
- |                      |           |                  |              |
|----------------------|-----------|------------------|--------------|
| SPOTTED SPURGE       | CRABGRASS | BERMUDA GRASS    | LAMBSQUARTER |
| OXALIS               | CRABGRASS | SHEPHERD'S PURSE | ARRPETWEED   |
| CRABGRASS            | CRABGRASS | EVENING PRIMROSE | DIRTY NOSEL  |
| THICKWEED (COMMON)   | CRABGRASS | STINK GRASS      | CALLIPANCIUM |
| CRABGRASS (FRONT)    | CRABGRASS | GERMAN MOSS      | RED SOPRANO  |
| CRABWEED (MOUSE EAR) | CRABGRASS | WITCH GRASS      | WATER WALKER |
| ANNUAL BLUEGRASS     | CRABGRASS | FLORIDA FLEECE   | RED TAIL     |
| SPINY AMARANTH       | CRABGRASS | ANNUAL SEED      | RED TAIL     |
- THE INSECTICIDE IN THIS PRODUCT IS DESIGNED TO PREVENT THE HEAVY INFESTATION AND AID IN THE CONTROL OF:
- |                  |           |      |       |
|------------------|-----------|------|-------|
| LAWN MOTH LARVAE | CUT WORMS | FLY  | GRASS |
| ARMY WORMS       |           | ANTS |       |

**INSTRUCTIONS FOR USE: (FOR ESTABLISHED DICONDRA LAWNS ONLY)**

READ AND FOLLOW ALL DIRECTIONS CAREFULLY

NOTICE: DO NOT APPLY ON OR NEAR YOUNG TREES OR YOUNG SHRUBS. NOT RECOMMENDED FOR DENSELY SHADED AREAS. AS DICONDRA MAY THIN OUT, SHOULD NOT BE USED ON NEW DICONDRA LAWNS UNTIL 60 DAYS AFTER SEEDING.

- FOLIAGE SHOULD BE DRY BUT SOIL MODERATELY MOIST.
- SPREAD THE CONTENTS OF THIS PACKAGE EVENLY OVER 250 SQ. FT. USE A FERTILIZER SPREADER, BEING SURE NOT TO OVERLAP.
- WASH MATERIAL OFF FOLIAGE AND INTO THE SOIL IMMEDIATELY AFTER APPLICATION. PREFERABLY WITH MEDIUM SPRAY FROM HAND HELD NOZZLE. THEN WATER THOROUGHLY.

OCT NOV DEC APPLICATION: AIDS IN PREVENTION OF POA ANNUAL WINTER BLUEGRASS. DELAYS GROWTH OF SPRING BERMUDA. PREVENTS EARLY GERMINATION OF OXALIS AND SPOTTED SPURGE.

JAN FEB MAR APPLICATION: THIS APPLICATION IS A NECESSARY FOLLOW UP FOR THE PREVENTION OF SPOTTED SPURGE. CONTINUED RETARDING OF BERMUDA GRASS AND PREVENTION OF MANY OF THE SPRING WEEDS AS LISTED.

APR MAY JUN APPLICATION: THIS APPLICATION CONTROLS MOST GRASSY WEEDS. FURTHER PREVENTS SPOTTED SPURGE AND OXALIS.

JUL AUG SEP APPLICATION: THIS APPLICATION IS FOR FURTHER AID IN ELIMINATING OXALIS GRASSY WEEDS AND CONTROL OF INSECTS.

NOTE: AT LEAST 90 DAYS SHOULD ELAPSE BETWEEN ANY APPLICATIONS AND NO MORE THAN 4 APPLICATIONS SHOULD BE MADE IN ANY 12 MONTH PERIOD. DICONDRA LAWNS CANNOT BE RESEED FOR 60 DAYS AFTER APPLICATION AS THE MATERIALS THAT PROVIDE THE RESISTIVE WEED



**3**

# SUPER 3-WAY WEEDILIZER<sub>21-0-0</sub>

a fertilizer with weed & insect control  
for established dichondra lawns only

GRANTED

INSTRUCTIONS FOR USE

INSTRUCTIONS FOR USE (FOR ESTABLISHED DICONDRA LAWNS ONLY)