

PEDERAL INSECTICIDE PUNGICIDE AND RODENTICIDE ACT

VEIGHT 24 LBS.



# Bandini **SUPER 2-WAY** PLUSH a fertilizer & insect control

# FOR DICHONDRA & GRASS LAWNS

# **GUARANTEED FERTILIZER ANALYSIS**

ORGANIC NITROGEN (DERIVED FROM UREA AND UREA FORMALDEHYDE) 18.60% (SOLUBLE NITROGEN 12 60% INSOLUBLE NITROGEN 6 00%) AMMONIC NITROGEN (DERIVED FROM AMMONIUM PHOSPHATE) 2.40% TOTAL NITROGEN 21.00% PHOSPHORIC ACID (AVAILABLE) (DERIVED FROM AMMONIUM PHOSPHATE) 6.00% POTASH (WATER SOLUBLE) (DERIVED FROM MURIATE OF POTASH) 6.00% CALCIUM (DERIVED FROM GYPSUM) 1.00% "SULFUR (COMBINED) (DERIVED FROM GYPSUM) 1.50% "IRON (EXPRESSED AS ELEMENTAL) 1.00% 'MANGANESE (EXPRESSED AS ELEMENTAL) 0.10% 'ZINC (EXPRESSED AS ELEMENTAL) 0.10%

\*DERIVED FROM SPECIALLY PROCESSED NATURAL MINERALS CONTAINING A MIXTURE WHICH GRADUALLY BECOMES SOLUBLE UPON EXPOSURE TO AIR AND WATER.

# **GUARANTEED INSECTICIDE ANALYSIS**

**ACTIVE INGREDIENT:** 

CARBARYL (1-NAPHTHYL N-METHYLCARBAMATE)

2.30%

INERT INGREDIENT:

FERTILIZER 21-8-6

97.70%

# CAUTION: KEEP OUT OF REACH OF CHILDREN AND PETS

AVOID BILIN CONTACT.

WASH HANDS THOROUGHLY BEFORE EATING.

IF SWALLOWED HIDUCE VOMITTING AND CALL PHYSICIAN IMMEDIATELY.

CARBARYL IS A MODERATE, REVERSIBLE, CHOLINESTERASE BUHISITOR ATROPINE

DO NOT CONTAINMATE FEED AND POODSTUPPS. LAWN CLIPPINGS AFTER APPLICATION, SHOULD NOT SE PED TO LIVESTOCK OR POULTRY.

DO NOT USE ON EDIBLE CROPS. AVOID EMEATHING OF DUST.

CONTAINER DISPOSAL: DO NOT RELIES EMPTY CONTAINER. Destroy R by burying with waste or huming. Stay every from arrate or furnies.

COMPLETE. BALANCED FERTILIZER ESPECIALLY DEVELOPED FOR ALL WESTERN LAWNS. IT SUPPLIES IMPORTANT MINOR ELEMENTS SUCH AS IRON, SULFUR, AND OTHER MINERALS TO THE SOIL.

THE INSECTICIDE IN THIS PRODUCT IS DESIGNED TO CONTROL OR PREVENT LAWN MOTH LARVAE FLEAS

ARMY WORMS

# TREES-FLOWERS-BEDDING PLANTS-SHRUBS-GROUND COVER

AS A GENERAL PURPOSE FERTILIZER USE 1 LB PER 100 SQ FT OF SOIL AREA (A 1 LG. COFFEE CAN HOLDS ABOUT 1 LB. OF THIS MATERIAL) WATER THOROUGHLY IMMEDI-ATELY AFTER APPLICATION. APPLICATIONS SHOULD BE REPEATED EVERY 6 TO 8 WEEKS THROUGHOUT THE GROWING SEASON

# INSTRUCTIONS FOR USE: (DICHONDRA OR GRASS LAWNS)

NEAD AND FOLLOW ALL DIRECTIONS CAREFULLY

- 1. FOLIAGE SHOULD BE DRY, BUT SOIL MODERATELY MOIST.
- 2 SEI SET START THE MILL OF LIGHT STERMS SEASON ASSOLUTE THE ASSOCIATION





April 19 April 19 Company

'SULFUR (COMBINED) (DERIVED FROM GYPSUM) 1.50% "IRON (EXPRESSED AS ELEMENTAL) 'MANGANESE (EXPRESSED AS ELEMENTAL) 'ZINC (EXPRESSED AS ELEMENTAL) 0.10%

\*DERIVED FROM SPECIALLY PROCESSED NATURAL MINERALS CONTAINING A MIXTURE WHICH GRADUALLY BECOMES SOLUBLE UPON EXPOSURE TO AIR AND WATER.

# **GUARANTEED INSECTICIDE ANALYSIS**

**ACTIVE INGREDIENT** 

CARBARYL (1-NAPHTHYL N-METHYLCARBAMATE)

2.30%

INERT INGREDIENT:

FERTILIZER 21-6-6

97.70%

PPLICATION, SHOULD NOT BE PED TO LIVESTOCK OR POULTRY.

DO NOT USE ON SOULE CROPS.

ER CHAPCHAL: DO NOT RELIEE PROFTY CONTAINER. Dealery it by Incyling with wante or burning. Stay away from amohe or famos.

THIS PRODUCT IS IN A CLEAN, EASY TO HANDLE PELLET FORM, AND FURNISHES A COMPLETE, BALANCED FERTILIZER ESPECIALLY DEVELOPED FOR ALL WESTERN LAWNS. IT SUPPLIES IMPORTANT MINOR ELEMENTS SUCH AS IRON, SULFUR, AND OTHER MINERALS TO THE SOIL.

THE INSECTICIDE IN THIS PRODUCT IS DESIGNED TO CONTROL OR PREVENT:

LAWN MOTH LARVAE ARMY WORMS

ANTS

# TREES—FLOWERS—BEDDING PLANTS—SHRUBS—GROUND COVER

AS A GENERAL PURPOSE FERTILIZER USE 1 LB. PER 100 SQ. FT. OF SOIL AREA (A 1 LB. COFFEE CAN HOLDS ABOUT 1 LB. OF THIS MATERIAL) WATER THOROUGHLY IMMEDI-ATELY AFTER APPLICATION. APPLICATIONS SHOULD BE REPEATED EVERY 6 TO 8 WEEKS THROUGHOUT THE GROWING SEASON.

# INSTRUCTIONS FOR USE: (DICHONBRA OR GRASS LAWNS)

- 1. FOLIAGE SHOULD BE DRY, BUT SOIL MODERATELY MOIST.
- 2. SELECT EITHER THE FULL OR LIGHT FEEDING PROGRAM. SPREAD THE CONTENTS OF THIS PACKAGE EVENLY OVER 2500 SQ. FT. OR 5000 SQ. FT.
- 3. APPLY WITH SPREADER FOR EVEN DISTRIPTION, BEING CAREFUL NOT TO
- 4. WASH MATERIAL OFF THE FOLIAGE AND INTO THE SOIL IMMEDIATELY AFTER APPLICATION, PREFERABLY WITH MEDIUM SPRAY FROM HAND-HELD NOZZLE, THEN WATER THOROUGHLY.

# APPROXIMATE SPREADER SETTINGS

IF YOU WALK SLOWER THAN NORMAL USE A HALF NUMBER SMALLER, IF YOU WALK FASTER THAN NORMAL USE A HALF NUMBER LARGER.

FULL FEEDING  2,900 SQ. FT. (USE 4 TIMES PER YEAR)  FOR FAST DEEP GREENING		LIGHT FEEDING  5,000 SQ. FT. (USE 0 TIMES PER YEAR)  WHEN FREQUENT FEEDING IS PREFERRED					
				FOR UPGRADING ALL LAWNS		FOR WELL-MAINTAINED LAWNS	
				FOR BERMUDA AND GRASS LAWNS		FOR FIRST FEEDING OF ALL NEW LAWNS	
FOR MISECT CONTROL		FOR MISECT PREVENTION					
BANDINI	5%	BANDINI	41/2				
CYCLONE MODEL B	54	CYCLONE MODEL B.	4				
SCOTT MODEL 35	64	SCOTT MODEL 35	54				

PLEASE CHECK WITH YOUR DEALER FOR THE PROPER SETTING OF UTHER SPREADERS.

CAUTION: ALWAYS SHUT SPREADER OFF WHEN FILLING, MAKING SHARP TURNS. OR STOPPING AVOID SPILLING.

PRODUCT OF BANDINI FERTILIZER COMPANY LOS ANGELES, CALIFORNIA 90023. U S A



SUPER 2-WAY

a fertilizer & insect control

PLUSH

# .

CAUTION: KEEP OUT OF REACH OF CHILDREN AND PETS

HARMFUL IF INHALED OR SWALLOWED

AVOID SKIN CONTACT

WASH HANDS THOROUGHLY BEFORE EAT NO.

IF SWALLOWED INDUCE YOU THING AND CALL PHYS AND MAST ATE I.

CAUTION: A MODIFIED CONTINUE AND MAST ATE I.

CAUTION STATEMENTS OF SWALLOWED

AVOID SKIN CONTACT

WASH HANDS THOROUGHLY BEFORE EAT NO.

IF SWALLOWED INDUCE YOU TIME AND CALL PHYS AND MAST ATE I.

WASH MANDS THOROUGHLY BEFORE EAT NO.

IF SWALLOWED INDUCE VOMITING AND CALL PHYS AN IMMETIATE OF

CARBARYL IS A MODERATE REVERSIBLE HICK RESTERASE NH B TO BOTH PING

IS ANTIDOTAL

DO NOT CONTAMINATE FEED AND FOODSTUFFS LAWM LOPIN IS ASTER

APPLICATION SHOULD NOT SEFED TO LIVESTY ROOF POTORY

DO NOT USE ON EDIBLE CROPS

AVOID BREATHING OF DUST

CONTAINER DISPOSAL DO NOT REUSE EMPTY LIMITANER HIMMETING OF TAKENY

with winds or burning. Stay away from arreits in furnes.

THE FUR COMBINED OFFICE OF WELL HAS WE

 $(\mathbf{x}_{1},\ldots,\mathbf{x}_{n}) \in (\mathbf{x}_{1},\ldots,\mathbf{x}_{n}) \times (\mathbf{x}_{1},\ldots,\mathbf{x}_{n}) \times (\mathbf{x}_{1},\ldots,\mathbf{x}_{n}) \times (\mathbf{x}_{1},\ldots,\mathbf{x}_{n})$ 

THIS PRODUCT SIN A CLEAN FACE COLONIC COLONIC

THE NSK TORE NOW THE SECOND STORES OF SECOND SECOND

# TREES-FLOWERS-BEDDING PLANTS - SHRUBS-GROUND CO 'ER

AN A SEMENAL OF HE SEMENT OF SEMENT

# INSTRUCTIONS FOR USE (DICHONDRA OR GRASS LAWNS) READ AND FOLLOW ALL DIRECTIONS CAREFULL!

- 1 FOLIAGE SHOULD BE DRY BUT SOIL MODERATELY MOIST
- SELECT EITHER THE FULL OR LIGHT FEEDING PROGRAM SPHEAD TE THE PATENTS.

  OF THIS PACKAGE EVENUT OVER 2500 SQ FT OR 5000 SQ FT.
- 3 APPLY WITH SPREADER FOR EVEN DISTRIC ON BEING CAREFUL NOT TO OVERLAP
- WASH MATERIAL OFF THE FOL AGE AND INTO THE SOIL IMMETIATELY ACTED APPLICATION PHEFERABLY WITH MET IN SPRAY FROM HAND HELD NOT IN THEN WATER THOROUGHLY

# APPROXIMATE SPREADER SETTINGS

FY C WALK SLOWER THAN NORMAL SEA FA FR. WHER MA AF FASTER THAN NORMAL SSEA HALCH MREF. AS

FULL FEEDING

SME SOURCE

THE ME SEEDING

THE ME SEEDING

THE MESTER OF THE SEEDING

THE SEEDING SEEDING

THE SEEDING SEEDING

THE SEEDING SEE