

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

MAR 3 0 2007

Mr. Masuo Robinson Agent Britz Fertilizers, Inc. c/o Robinson Associates 583 Canyon Road Redwood City, CA 94062

RE: Correct the First Aid Statement for if Swallowed as well as Misspelled Words

EPA Registration Number: 10951-19 Date of Submission: January 23, 2007

Dear Mr. Robinson:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated January 23, 2007, for the product, Britz Fertilizers, Inc. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the actions requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Joyce Edwards of my staff at 703-308-5479.

Sincerely,

Linda Arrington

Notifications & Minor Formulations Team Leader Registration Division (7505P)

Office of Pesticide Programs

Please read instructions on reverse before c	ompleting form. Form	Approved, OMB N	lo. 2070-0060, Approva	l expires 05-31-98	
TOTAL A	United States		Registra	ition	OPP Identifier Number
EPA Environ	nmental Protection	Agency	Amenda		RA2002
	Washington, DC 20460		=	Notification	KA2002
	Appl	ication for Pe	sticide - Section		····
1. Company/Product Number					3. Proposed Classification
10951-	19	Z. EPA PI	oduct Manager		3. Proposed Classification
10001	• 7	John H	łebert		
4. Company/Product (Name)		PM#			None Restricted
Britz F	ertilizers, Inc.	7			
5. Name and Address of Applicant (Include	'e ZIP Code)	6. Expedi	ted Review. In accorda	nce with FIFRA Secti	on 3(c)(3)(b)(l), my product is similar
,	,	or identical	in composition and lab	eling to:	•
Britz Fertilizers, Inc.		FPA R	eg. No		
c/o Robinson Associates			·g. 110.		
583 Canyon Rd. Redwood City, CA 94062		Product	Name		
Check if this is a new ad	lduare				
Check if this is a new ad	aress		· · · · · · · · · · · · · · · · · · ·		
		Section	n - 11		NOTIFICATION
Amendment - Explain below.			Final printed lab	els in response to Age	ency letter dated
Resubmission in response to Agend	y letter dated	_	"Me Too" Appli	cation	MAR 3 0 2007
Notification - Explain below.		<u>.</u>	Other - Explain	below	
Explanation: Use additional pa	.ge(s) if necessary. (F	For Section I an	d Section II.)		
This notification is consistent with the the labeling or the confidential statement to EPA. I further understand be in violation of FIFRA and I may be	ent of formula of this prother that if this notification	oduct. I understa is not consistent	nd that it is a violatio with the terms of PR	n of 18 U.S.C. Sec. Notice 98-10 and 4	1001 to willfully make any false 0 CFR 152.46, this product may
	·····	Section	n - III		,
1. Material This Product Will Be Packa					
Child-Resistant Packaging	Unit Packaging		Water Soluble Pack	aging	2. Type of Container
Yes*	Yes		Yes		Metal
No *Certification must be	No If "Yes"	No. per	If "Yes"	No. per	Plastic
submitted	Unit Packaging wgt.	container	Package wgt.	container	Glass
Submittee					Other (Specify)
Location of Net Contents Information	4. Size(:) Retail Container	-	5. Location of 1	Label Directions
Label Contain	1	,,		On Label	
				On labeling	g accompanying product
6. Manner in Which Label is Affixed to	Product Life	thograph	Other_		
•		per glued			
	Ste	enciled			
		Section	ı - IV		
1. Contact Point (Complete items direct)	y below for identification c	of individual to be c	contacted, if necessary,	o process this applica	ntion)
Name		Title			Telephone No. ('nclude Area Code)
Masuo Robinson		Agent			650.361.1101
	Certific	ation		11	6. Date Application Received
I certify that the statements I have made that any knowingly false or misleading s	on this form and all attach	ments thereto are to			(Stamped)
2. Signature		3. Title		•	
Man 2	5	Agent		<u>,</u>	
4. Typed Name Masuo Robinso		5. Date Janua	ry 23, 2007		
Masuu Kudiiisu	j	vanua	m y 200, 200,		



BFR 440 SUPREME SPRAY OIL

ACTIVE INGREDIENT:	BY WT.
Petroleum Oil	99.00%
INERT INGREDIENTS	<u>1.00%</u>
TOTAL	100.00%
Unsulfonated Residue	95.00%
Aromatic Composition ASTM D2140	0.00%

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

ic. u	0.11
If swallowed:	•Call a poison control center or doctor
ļ	immediately for treatment advice.
	•Do not give ANY liquid to the person.
	•Do not induce vomiting unless told to do so
	by a poison control center or doctor.
	•Do not give anything by mouth to an
	unconscious person.
If in eyes:	 Hold eyes open and rinse slowly and gently
	with water for 15-20 minutes. Remove contact
	lenses, if present, after the first 5 minutes,
	continue rinsing.
	•Call a poison control center or doctor for
	treatment advice.
If on skin or	•Take off contaminated clothing.
clothing:	Rinse skin immediately with plenty of water
_	for 15-20 minutes.
	•Call a poison control center or doctor for
	treatment advice
In inhaled:	•Move person to fresh air.
	If person is not breathing, call 911 or an
	ambulance, and then give artificial respira-
	tion, preferably mouth to mouth if possible.
	•Call a poison control center or doctor for
	further treatment advice.

NOTE .TO PHYSICIAN: Contains petroleum distillates. Vomiting may pose aspiration pneumonia.

FOR MEDICAL EMERGENCY INVOLVING THIS PRODUCT, CALL: 1-800-228-5635, EXT. 136 OR CALL COLLECT, 612-851-8180, EXT. 136.

EPA EST. NO. 10951-CA-1		
NET CONTENTS	GAL. () L

EPA REG. NO. 10951-19



NOTIFICATION

MAR 3 0 2007

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin or inhaled. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. This product is a potential skin sensitizer. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

Personal Protective Equipment (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on the EPA chemical resistance category selection chart

Applicators and other handlers must wear: long-sleeved shirt and long pants, chemical-resistant gloves, such as barrier laminate, nitrile rubber, butyl rubber, neoprene or viton, and shoes plus socks.

Follow manufacturer's instruction for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering controls statements:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets with requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water. Apply this product only as specified on the label.

PHYSICAL OR CHEMICAL HAZARDS

Do not weld, heat, or drill this container. Residue may ignite with explosive violence if heated sufficiently.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or thorough drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

4/8

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR, Part 170. This Standard contain requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval of 4 hours.

PPE require for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is: coveralls, waterproof gloves, shoes plus socks.

STORAGE AND DISPOSAL

Do not contaminate water, food and feed and water by storage or disposal.

PESTICIDE STORAGE: Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, replace lids and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call collect: CHEMTREC (800) 424-9300.

To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or an approved waste disposal facility.

CONTAINER DISPOSAL: Metal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Plastic: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay of smoke.

APPLICATION

For applications, the spray tank must be free of any sulfur residue before using BFR 440 Supreme Spray Oil. Do not combine this product with any type of sulfur spray. Do not apply BFR 440 Supreme Spray Oil after a sulfur application until sufficient time has passed for safety (at least 30 days). Do not apply sulfur after an application of BFR 440 Supreme Spray Oil until sufficient time has passed for safety (at least 30 days). If doubts exist, spray BFR 440 Supreme Spray Oil mixture on a portion of one tree that was treated with sulfur. If no burn is noted in several days, enough time has passed. Do not spray when temperatures are above 95°F or when temperatures drop below 35°F. Do not use on foliage closer than 60 days after or 90 days before using Captan or Folpet products.

Application to trees weakened by disease, drought, drying winds, or high nitrogen application may result injury to the trees.

Do not apply this product through any type of irrigation system.

Do not apply if this product will drift to food forage or other crops that might be damaged.

BFR 440 Supreme Spray Oil must be diluted before an application is made.

MIXING DIRECTIONS

Check to make sure this product will emulsify. If the mixture is not milky after agitation, do not use this product. Do not use this material if it does not emulsify.

With the agitator running, start filling the spray tank with water. Add wettable powders or other products when the tank is one third full and always before adding BFR 440 Supreme Spray Oil. When the tank is one half full and with good agitation, add the proper amount of BFR 440 Supreme Spray Oil, then fill to the desired volume. Constant agitation must be maintained during mixing application.

DO NOT MIX with insecticides or fungicides incompatible with oil. Consult your State Agricultural Extension Service or Fieldman for specific compatibilities.

This product may be used as a carrier for appropriate insecticides and fungicides which are registered for use on deciduous fruits and nuts and which can be applied to give a uniform spray. For recommendations on timing and combinations with BFR 440 Supreme Spray Oil, Consult your local Agricultural Extension Service or Fieldman.

Use only in spray machines with powerful positive agitation. Keep agitator running at all times during spray operations.

This product is formulated as a "fast break" spray oil and will not stay in solution unless agitated constantly.

ORCHARD CONCENTRATE APPLICATION: Apply according to rates given in the table for each crop. All quantities of spray solution are for mature trees and vines with normal foliage cover. Quantities should be reduced proportionally for smaller immature trees and vines. Use only on crops listed on this label. Special concentrate spray equipment is necessary for this application.

NOTE: Use efficient equipment of the proper type. Avoid spraying during or immediately prior to hot or freezing weather (over 95°F or under 35°F), hot dry wind, rain or other unsuitable conditions; avoid over spraying or double spraying. Plants should be sprayed only when in vigorous condition and when their moisture condition is suitable.



ALMONDS, APRICOTS, CHERRIES, NECTARINES,

PISTACHIOS, PEACHES, PLUMS AND PRUNES: (Directions for use on plums and prunes in California's Sacramento Valley can be found in the table following this one):

PEST	APPLICATION RATE & TIMING	COMMENTS
Black Scale, Brown	DORMANT AND	Preferred timing of
Soft Scale,	DELAYED	application for
European Fruit	DORMANT: Use	European Fruit
Lecanium Scale,	1 to 2 gallons per	Lecanium Scale
Frosted Scale,	100 gals of water at	and European Red
Olive Scale,	300 to 500 gals per	Mite eggs is green
San Jose Scale,	acre. (Concentrate	up to Delayed
European Red Mite	Spray: Use 5 to 7	Dormant and for
Eggs,	gals per acre in a	San Jose Scale is
Aphid Eggs,	minimum of 20 gals	Dormant, Green tip
Brown Mite Eggs,	by air or 100 gals	to delayed dormant.
Fruitree Leafroller	by ground of water	
Eggs	рег асте).	
Terrapin Scale	DELAYED	
	DORMANT: Use	
	2 to 3 gals per 100	
	gals of water at 300	
	to 600 gals per acre.	
	(Concentrate Spray:	
	Use 4 to 6 gals per	
	acre in a minimum	İ
	of 20 gals by air or	
	100 gals by ground	
Cottony peach	of water per acre). DORMANT: Use	
Scale	2 to 3 gals per 100	
Scarc	gals of water at 300	
	to 600 gals per acre.	
	(Concentrate Spray:	
	Use 4 to 7 gals per	
	acre in a minimum	•
	of 20 gals by air or	
	100 gals by ground	
	Of water per acre).	
Two Spotted Spider	FOLIAR SPRAY:	Certain varieties of
Mites,	Use ½ to 1½ gals	plums and prunes
Pacific Mite,	per acre.	may be injured by
European Red Mite,	(Concentrate Spray:	all sprays, check for
Brown Mite	Do not use more	tolerance prior to
	than 4 gals per acre	treatment.
	in a minimum of 20	
	gals by air or 100	
	gals by ground of	
	water per acre).	
White Peach Scale	DORMANT: Use	Apply before buds
on Peaches: Only	1½ to 2 gais per	begin to swell in
for use in Georgia,	100 gais of water at	the spring. Make 2
Florida, South Carolina, North	300 to 600 gals per	applications 7 to 14
Carolina, North	(Concentrate Spray)	days apart,
Carolina and Alabama.	(Concentrate Spray:	according to local recommendations.
Alabama.	Use 5 to 7 gals per acre in a minimum	during the dormant
	of 20 gals by air or	period.
	100 gals by ground	portou.
	of water per acre).	
	or water per acre).	

PLUMS AND PRUNES: (Sacramento Valley of California Only)

PEST	APPLICATION RATE & TIMING	COMMENTS
Black Scale, European Fruit Lecanium Scale, San Jose Scale, Olive Scale, Frosted Scale, Mealy Plum Aphid Eggs, Leaf Curl Plum Aphid Eggs, Brown Mite Eggs, European Red Mite Eggs	PRE-BLOOM WHEN FRUIT BUDS SHOW GREEN TIPS: Use 1½ to 2 gal per 100 gals of water at 300 to 600 gals per acre. (Concentrate Spray: Use 4 to 6 gals per acre in a minimum of 20 gals by air or 100 gals by ground of water per acre.	Application at this time is preferred.
Two spotted Spider, Pacific Spider, European Red Mite, Brown Mite	FOLIAR SPRAY: Use ½ to 1 gal per 100 gals water at 300 to 600 gals per acre. (Concentrate Spray: Use 4 gals maximum per acre in a minimum of 20 gals by air or 100 gals by ground of water per acre).	Oil sprays may injure certain varieties of plums and prunes, check for tolerance prior to treatment.

For Aerial Application: Apply by air only when soil conditions do not permit regular ground application.

3

CITRUS: (California Only) – Lemons, Grapefruit, and Oranges (Navel and Valencia), Limes, Tangerines, Tangelos, and other citrus varieties.

PEST	APPLICATION RATE & TIMING	COMMENTS
Mites	Central California:	Do not apply to
-Citrus Bud,	Apply 0.5 to 1.4 gals per	lemons within 60
Citrus Flat.	100 gals of water as	days of harvest if
,	1 -	1 •
Citrus Red,	thorough coverage spray	lemons are to be
Citrus Rust	(TC).	sweated.
	Alternately, use a low	For lemons and
	volume spray (LV) to	limes, make
	apply 2.5 to 15 gals per	applications in
Scales	acre.	June through
-Black,		September. For
Brown, Soft,		grapefruits make
California		applications in
Red, Citricola		August through
Purple,	ĺ	September. For
Yellow, Red		
renow, kea		oranges, make
		applications on
i	1	navels and
	1	valencias - July
•		through August
1		and on tangelos,
		tangerines, and
1		other citrus
		varieties - July
		through August.
	•	unough tregues
		For navels and
	1	valencias.
		treatment can be
		made from Feb 15
		to 50% bloom, to
		1
		avoid injury, use
		low concentration
		(1.0%).
		On navel oranges,
		do Not apply
		during or after
		color break.
		COIOI DICAN.
Mites	Oxnard Plains Area:	For lemon,
-Citrus Bud,	Apply 0.5 to 1.4 gals per	grapefruits,
Citrus Flat.	100 gals of water as a	oranges (navels
Citrus Red.	thorough coverage spray	and valencias).
Citrus Rust	(TC). Alternately, use a	limes, tangelos,
Cidus Nust	low volume spray (LV) to	tangerines, and
	apply 1 to 10 gals per	other citrus
	acre.	varieties, make
i		applications as
		needed throughout
Scales		the year.
Black, Brown		On navels, do Not
Soft,		apply during or
California		after color break.
Red, Citriola,		Do not apply to
Purple,		lemons within 60
Yellow, Red		days of harvest.

Thorough coverage spray (TC): Applications achieving a uniform film wetting of all portions of the tree at 250 to 4500 gallons of spray solution per acre, the total volume is dependent of the size of the trees.

Low volume spray (LV): Applications with droplet depositions achieved on all interior and exterior parts of the tree using 20 to 100 gallons of spray solution per acre.

Do not apply prior to or during a period of excessively high temperatures above 95°F. (85° TO 95°F on the coast)

Do not apply oil sprays when trees show stress.

Consult with you local University of California Cooperative Extension Specialist or Fieldman for proper timing and spray volumes.

APPLES & PEARS

PEST	APPLICATION RATE & TIMING	COMMENTS
San Jose Scale, Italian Pear Scale	Dormant & Delayed Dormant: Apply 1½ to 2 gallons per 100 gallons of water in a minimum of 250 gallons per acre. (Concentrate sprays: do not use more than 6 gals per acre in a minimum of 20 gals by air or 100 gals by ground of water per acre).	An application of an insecticide labeled for use on apples and pears will improve scale control.
Leaf Roller Eggs	Delayed Dormant: Apply 2 gallons per 100 gallons of water at 300 to 500 gals per acre. (Concentrate Spray: Apply 5 to 6 gals per acre in a minimum of 20 gals by air or 100 gals. by ground of water per acre.)	
Pear Psylla	Apply 1 gallon per 100 gallons of water at 300 to 600 gals per acre. (Concentrate Spray: Apply 3 to 6 gals per acre in a minimum of 20 gals by air or 100 gals by ground of water per acre.)	:
Red Mites, San Jose Scale, Italian Pear Scale (immature summer stage of scales)	Summer Application (post harvest) Apply 1 gallon per 100 gallons of water at 300 to 600 gals per acre.	An application of an insecticide labeled for use on apples and pears will improve summer scale control.

AVOCADOS

PEST	APPLICATION RATE & TIMING	COMMENTS
Avocado Brown Mite, Thrips	Use 1 to 4 gals per 100 gals of water for thorough coverage in the 400 to 3000 gals per acre. (Concentrate Spray: Use 4 to 15 gals per acre in 50 to 100 gals of water per acre.) Do not exceed 30 gallons of oil per acre per application.	Apply through late fall. Consult with your local University of California Cooperative Extension Service or Fieldman for proper timing and application parameters. When an application of oil is more than 2% v/v, it is recommended that a small test area be treated and observed.

GRAPES (Dormant)

PEST	APPLICATION RATE & TIMING	COMMENTS
Mealy Bugs	Dormant: Use 1 to 2 gallons per 100 gals of water per acre with thorough coverage. Do not exceed 300 gals per acre per application	For improved control tank mix with a registered insecticide.

KIWI

PEST	APPLICATION RATE & TIMING	COMMENTS
Latonia Scale, Oleander Scale, San Jose Scale	Dormant: Apply 4 to 6 gallons per treated acre. Apply in a minimum 100 gallons of water per acre.	Do Not apply after bud break. Use the higher rate when scale populations are high.

OLIVES

PEST	APPLICATION RATE & TIMING	COMMENTS
Parlatoria Scale, Oleander scale, Black Scale	Apply 1½ gallons per 100 gallons of water as a dilute spray in 400 to 800 gallons of water per acre.	Apply prebloom or post bloom up to buckshot. Do not apply any oil between Aug 20 and harvest to Olives used for Spanish or green-ripe processing be- cause of the risk of fruit spotting.

PECANS

PEST	APPLICATION RATE & TIMING	COMMENTS
Scales, Aphid Eggs, Mite Eggs	Apply 2 to 3 gallons in 100 gallons of water using minimum of 200 gallons water per acre. (Concentrate Spray: Apply 6 to 8 gallons per acre in a minimum of 20 gals by air or 100 gals by ground of water per acre).	Apply from late February up to bud break.

WALNUTS

PEST	APPLICATION RATE & TIMING	COMMENTS
Brown Apricot Scale, San Jose Scale, Olive Scale, Oleander Scale, Frosted Scale, Italian Pear Scale, European Fruit Lecanium, Putnam Scale, Brown Soft Scale, Red Spider Mites, Aphid Eggs	Delayed Dormant (as buds begin to swell): Apply 1 gallon per 100 gallons of water. Apply as a dilute spray in 400 to 3,000 gallons per acre. As a concentrate spray: Use 4 to 6 gallons in a minimum of 100 gals of water per acre.	Oils should not be used on walnuts during the dormant season, between bud break and shoot elongation, or on drought stressed trees.
Red Spider Mites	Summer Use: Apply 1 ½ to 2 gallons per 100 gallons of water. Apply as a dilute spray in 400 to 3,000 gallons per acre. As a concentrate spray: Use 4 to 6 gallons in a minimum of 100 gals of water per acre.	Use only on trees that have not suffered from lack of moisture during the growing season. Do Not apply if the temperature will exceed 90°F at or following application
Brown Apricot Scale, San Jose Scale (summer stage)	Summer Use: Apply 1 gallon per 100 gallons of water. Apply as a dilute spray in 400 to 3,000 gallons per acre.	Do not apply after husk split. Do Not apply if the temperature will exceed 90°F at or following application.

Manufactured By BRITZ FERTILIZERS, INC. P.O. Box 366 Five Points, CA 93624

CONDITIONS OF SALE AND WARRANTY

1.Britz Fertilizers, Inc. warrants that this material conforms to the chemical description on the label and is reasonably fit for use as a directed hereon. To the extent consistent with applicable law, Britz Fertilizers, Inc. neither makes, nor authorizes any agent or representative to make any other warranty of fitness or of merchantability, guarantee or representation, expressed or implied, concerning this material.

2. Critical and unforeseeable factors beyond the control of Britz Fertilizers, Inc prevent it from elimination all risk in connection with the use of chemicals. Such risks include, but are not limited to damage, to plants and crops to which the material is applied, lack of complete control and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Britz Fertilizers, Inc. under 1 above) resulting from handling, storage, and use of this material.

"Britz for Results"

January 23, 2007

Document Processing Desk [NOTIF]
Office of Pesticide Programs (7504C)
U.S. Environmental Protection Agency
One Potomac yard
277 S. Crystal Drive
Arlington, VA 222021

Attn:

John Hebert

Product Manager, Team 7

Subject:

BFR 440 Supreme Spray Oil,

EPA Reg. No. 10951-19

Notification of Minor Changes to the Label.

Dear Mr. Hebert:

On behalf of Britz Fertilizers, Inc., we are submitting a notification to inform you of minor changes to the label. We have corrected the required language in the first aid statement for "If swallowed" and corrected several misspelled words.

In support of this notification, please find the following:

- EPA Form 8570-1; Application for Pesticide Registration dated January 23, 2007.
- Two copies of the revised label (one clean and one marked label).

Should you have any questions regarding this submission or require additional information, please call me. Please send all correspondence to Robinson Associates.

Sincerely,

Masuo Robinson Regulatory Specialist

attachments

cc: Tim Brown, Britz Fertilizers, Inc.

ra2002

Regulatory Affairs Pesticide Registration 583 Canyon Road

Redwood City, CA 94062

Telephone 650.361.1101

Facsimile 650.361.0111

Email

masuo@ix.netcom.com