

# PROKIL Cryolite 50 Dus

Active Ingredients	by Wt
Sodium Fluoroaluminate*	50
Inert Ingredients	50
	100
* Fluorine expressed as elemental	27.1

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
Avoid Inhalation of Dust

THIS PRODUCT IS HIGHLY TOXIC TO BEES exposed to treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

**NET WEIGHT 50 LBS.**

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. USES, TIMING AND DOSAGE MAY VARY AS A CONSEQUENCE OF LOCAL WEATHER OR CONDITIONS. WE RECOMMEND REFERENCE TO STATE AGRICULTURAL EXTENSION SERVICE CONCERNING SPECIFIC USES.

KEEP PESTICIDE IN ORIGINAL CONTAINER!

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.


This product is formulated for application by ground or air equipment

Use at rate of 25 to 45 pounds per acre depending on plant size and insect population. Repeat application as necessary. Remove residues remaining at harvest by washing, brushing or other effective means. Do not apply under conditions involving possible drift to food, forage or other planting that might be damaged or the crops thereof rendered unfit for sale, use or consumption

3-71-113 Proof Set

U.S.D.A. Reg. No. 10163 22

CALIFORNIA (Plant)  
P. O. Box 406  
340 East Main Street  
Calipatria, Calif. 92232  
Telephone: 348-5150



**The Dune Company**  
AGRICULTURAL CHEMICALS

Route 1,  
Somerton, Arizona  
Telephone: 62

BEANS: Beetles, Bean Leafroller — Repeat at two week intervals..

CABBAGE, CAULIFLOWER, BROCCOLI: Flea Beetles, Cabbage Caterpillar — NOT APPLY to Broccoli after the edible parts start to form. DO NOT APPLY to Cabbage after the heads start to form. DO NOT APPLY to Cauliflower the seedling stage.

CARROTS: Vegetable Weevil.

CUCUMBERS: Striped Cucumber Beetle, Twelve-Spotted Beetle, Flea Beetle — Dust both sides of foliage at first.

EGGPLANT: Flea Beetle — Dust thoroughly.

LETTUCE: Climbing Cutworms, Alfalfa Looper, Corn Earworm — Apply when worms first appear. DO NOT APPLY after the seedling stage.

MELONS: Cucumber Beetle — Dust both sides of foliage. Repeat if necessary.

PEPPERS: Pepper Weevil — Dust thoroughly at first set of fruit. Repeat 7 to 10 days until about 6 applications have been made.

STRAWBERRIES: Leaf Beetle — Dust both sides of leaves.

TOMATOES: Flea Beetle — Apply when plants are first set out and necessary. If applied after fruit starts to set, remove residue at harvest by washing, brushing or other effective means.

## NOTICE ON CONDITIONS OF SALE

The representations for use of this product are based upon tests conducted under controlled conditions. The use of this product being beyond the control of the manufacturer, no warranty is made as to the effects of such or the results of its use. It is sold as a general purpose product and is not intended for use in conjunction with any other material.