

CAUTION: Keep Out Of Reach Of Children. Use this product only in accordance with label directions. Keep children and pets away from treated areas until completely dry. Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing spray mist. Avoid contact with skin, eyes and clothing. In case of contact, wash skin with soap and water promptly. Flush eyes repeatedly with running water and get medical attention. Wash contaminated clothing with soap and hot water before re-use. Do not contaminate ornamental fish ponds or other bodies of water by direct application, cleaning of equipment or disposal of wastes and containers. Do not use on edible plants. Do not contaminate or store near feed or foodstuffs. Wash after handling and before eating or smoking. When container is empty, wash thoroughly and destroy. Do not transfer to unlabeled container.

This Product Is Toxic To Fish, Birds And Other Wildlife.

NOTE TO PHYSICIAN: Atropine sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine. **WARNING-FLAMMABLE! KEEP AWAY FROM HEAT OR OPEN FLAME. NOTICE TO BUYER:** Seller makes no warranty expressed or implied concerning the use of this product. Buyer assumes all risk of using or handling.

HACIENDA ENTERPRISES
565 Charles St
San Jose, Calif. 95112

Hacienda

INSECT-MILDEW SPRAY

ACTIVE INGREDIENTS:

Dimethyl (2,2,2-trichloro-1-hydroxyethyl) phosphonate	7.00%
S-(2-Ethylsulfinyl)ethyl 0,0-dimethyl phosphorothioate	5.00%
2,4-Dinitro-6-octyl* phenyl crotonate	} 1.25%
2,6-Dinitro-4-octyl* phenyl crotonate	
Nitrooctyl* phenols (principally dinitro)	0.08%
INERT INGREDIENTS	86.67%

*A mixture of 1-methylheptyl, 1-ethylhexyl and 1-propylentyl isomers.

NET CONTENTS: 1 PINT

APPLICATION:

Apply as full coverage and fine mist spray. Partly fill sprayer with water, add insecticide under agitation, add balance of water. Cover all plant surfaces, top leaf, under leaf and stems. Avoid spraying at mid-day when temperature is above 85°F. Water plants one day previous to spraying. Always prepare fresh spray solution at time of use.

DIRECTIONS FOR USE

DILUTION: Mix 1 oz. in one gallon water.
ORNAMENTAL TREES, ROSES, FLOWERS AND SHRUBS: Apply every 2 or 3 weeks or as required to control Aphis; Leafhoppers; Leafminers-Birch, Dipterous and ~~leaf~~; Lygus Bugs; Mites; Thrips; Whiteflies. For control of Armyworms; Bagworms; Cutworms-Varigated and Climbing; Stinkbug; Tarnished Plant Bug; Webworms; apply twice at 1 week intervals for heavy infestations, otherwise every 2 or 3 weeks or when required.

POWDERY MILDEW: To clean up heavy infestations, apply several times at 4 to 7 day intervals. For maintenance under mild conditions, apply every 10 days or as necessary.

DO NOT APPLY THIS MATERIAL TO BEARING TREES OR VEGETABLES.

CAUTION: Keep Out Of Reach Of Children. Use this product only in accordance with label directions. Keep children and pets away from treated areas until completely dry. Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing spray mist. Avoid contact with skin, eyes and clothing. In case of contact, wash skin with soap and water promptly. Flush eyes repeatedly with running water and get medical attention. Wash contaminated clothing with soap and hot water before re-use. Do not contaminate ornamental fish ponds or other bodies of water by direct application, cleaning of equipment or disposal of wastes and containers. Do not use on edible plants. Do not contaminate or store near feed or foodstuffs. Wash after handling and before eating or smoking. When container is empty, wash thoroughly and destroy. Do not transfer to unlabeled container.

This Product Is Toxic To Fish, Birds And Other Wildlife.

NOTE TO PHYSICIAN: Atropine sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine. **WARNING-FLAMMABLE! KEEP AWAY FROM HEAT OR OPEN FLAME. NOTICE TO BUYER:** Seller makes no warranty expressed or implied concerning the use of this product. Buyer assumes all risk of using or handling.

HACIENDA ENTERPRISES

505 S. ...
... ..

EPA Reg. No. 11037-3

Hacienda

INSECT-MILDEW SPRAY

**SYSTEMIC - RESIDUAL - CONTACT
FUNGICIDAL ACTION**

ACTIVE INGREDIENTS:

Dimethyl (2,2,2-trichloro-1-hydroxyethyl) phosphonate 5.00%
S-(2-Ethylsulfanyl)ethyl) 0,0-dimethyl phosphorothioate 5.00%
2,4-Dinitro-6-octyl-phenyl crotonate 1.25%
2,6-Dinitro-4-octyl-phenyl crotonate 1.25%
Nitro-octyl-phenols (principally dinitro) 0.08%
INERT INGREDIENTS 86.67%

A mixture of 1-methylheptyl, 1-ethylhexyl and 1-propylpentyl isomers

NET CONTENTS: 1 PINT

**CAUTION: KEEP OUT OF REACH OF CHILDREN.
READ OTHER CAUTIONS ON BACK PANEL.**

APPLICATION:

Apply as full coverage and fine mist spray. Partially fill sprayer with water, add insecticide under agitation, add balance of water. Cover all plant surfaces, top leaf, under leaf and stems. Avoid spraying at mid-day when temperature is above 85° F. Water plants one day previous to spraying. Always prepare fresh spray solution at time of use.

DIRECTIONS FOR USE

DILUTION: Mix 1 oz. in one gallon water.
ORNAMENTAL TREES, ROSES, FLOWERS AND SHRUBS: Apply every 2 or 3 weeks or as required to control Aphis; Leafhoppers; Leafminers-Birch, Dipterous and Holly; Lygus Bugs; Mites; Thrips; Whiteflies. For control of Armyworms; Bagworms; Cutworms-Varigated and Climbing; Stinkbug; Tarnished Plant Bug; Webworms; apply twice at 1 week intervals for heavy infestations, otherwise every 2 or 3 weeks or when required.

POWDERY MILDEW: To clean up heavy infestations, apply several times at 4 to 7 day intervals. For maintenance under mild conditions, apply every 10 days or as necessary.

DO NOT APPLY THIS MATERIAL TO BEARING TREES OR VEGETABLES.