

Hacienda

POST-EMERGENT CRABGRASS CONTROL

CAUTION: Keep Out Of Reach Of Children. Use this product only in accordance with label directions. Harmful if swallowed. Avoid contact with skin. Avoid breathing spray mist. Wash hands after using. Avoid storage near feed or food products. Do not contaminate waters used for domestic consumption, or by animals, wildlife and aquatic life, or for irrigation purposes. Do not feed treated foliage to livestock or graze treated areas. Keep children and domestic animals off treated areas until this material has been washed into the soil. Do not apply directly, or allow to come in contact with desirable trees or ornamentals.

Do not re-use empty container. Wash container and equipment thoroughly with water and detergent. Dispose of wastes and containers by burying in non-crop lands away from water supplies. Do not store spray solution in tank for a prolonged period.

ANTIDOTE: If taken internally, induce vomiting and call physician at once.

NOTICE TO BUYER: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions GIVEN HEREWITH.

HACIENDA ENTERPRISES
565 Charles Street
San Jose, Calif. 95112

EPA Reg. No. 11037-5

ACTIVE INGREDIENTS:		Total Arsenic (as elemental)	
Monosodium Acid		all in water soluble form	8.8%
Methanearsonate*	19.0%	*Product contains 1.8	
INERT INGREDIENTS	81.0%	MSMA per gallon.	

NET CONTENTS: 1 PINT

**CAUTION: KEEP OUT OF REACH OF CHILDREN.
READ OTHER CAUTIONS ON BACK PANEL.**

HACIENDA Post-Emergent Crabgrass Control is a combination of herbicide and surfactant, IT IS UNNECESSARY TO ADD ADDITIONAL SURFACTANT. Its phytotoxic properties are quickly inactivated on contact with the soil. Best results are obtained on young actively growing weeds at air temperatures above 70°F.

APPLICATION—Easily applied as a spray with tank sprayer, trombone type, or hose end insecticide sprayer. Partly fill sprayer with water, add proper amount of material, add balance of water. Apply as a spray, thoroughly wetting weeds to point just short of run-off.

DIRECTIONS—LAWN and ORNAMENTAL TURF: To Control CRABGRASS, CHICKWEED, NUTSEGE, DALLISGRASS, SANDBUR, BAHAGRASS and WOOD SOBREL. On established turf little or no damage will occur other than a temporary discoloration. Do not treat new lawns until after three mowings. DO NOT USE on St. Augustine or Centipede grasses.

Irrigate and mow lawns to height of 1-1½ inches before treatment. Mix 4.0 fluid ounces in 2½ gallons of water (¾ table-spoons in 1 gallon) and apply this amount to 1000 sq. ft. Spray thoroughly to wet all undesirable plants then withhold irrigation of treated areas for two days following application. Repeat applications, 10 to 14 days apart may be needed for good control.

DRIVE CROP: To control JOHNSONGRASS, NUTSEGE, DALLISGRASS, COCKLEBUR, RAGWEED, SANDBUR PUNCTURE VINE on Drainage Ditchbanks, Rights-Of-Way, Fence Rows and Storage Yards. Mix 4 to 8 ounces in 2½ gallons of water (¾ - 6½ tablespoons in 1 gallon). Spray unwanted vegetation with low volume coverage, thoroughly wetting weeds to a point just short of run-off. If regrowth occurs, repeat as necessary.

COVERAGE ONE PINT — 4,000 sq. ft.

