


Hacienda



# MITE-MILDEW CONTROL

**CAUTION:** Keep Out Of Reach Of Children. Use this product only in accordance with label directions. Keep children and pets away from treated areas until completely dry. Avoid contact with skin or eyes. May cause skin irritation or sensitization. Wash thoroughly after use. Avoid inhalation of spray mist. Harmful if swallowed. If ingested, get medical attention. Do not store near food or foodstuffs. Never re-use containers.

"This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label."

Do not apply when temperature will exceed 90 F during the day. **WARNING-FLAMMABLE! KEEP AWAY FROM HEAT OR OPEN FLAME.** NOTICE: Buyer assumes all risk of use, storage or handling of this material not in strict accordance with label directions.

## HACIENDA ENTERPRISES

565 Charles Street  
San Jose, Ca. 95112

ACTIVE INGREDIENTS  
2,4-Dinitro-6-octyl\*phenyl crotonate 4.4%  
2,6-Dinitro-4-octyl\*phenyl crotonate 0.3%  
Nitrooctyl\*phenols (principally dinitro) 95.3%  
INERT INGREDIENTS

\*A mixture of 1-methylheptyl  
1-ethylhexyl and 1-propyl entyl isomers.

NET CONTENTS: 1 PINT

**ROSES, BEGONIA, BELLADONNA, DAHLIA, CHRYSANTHEMUM, CALENDULA, DELPHINIUM, SNAPDRAGON, ORNAMENTAL TREES and SHRUBS, LILAC, HYDRANGEA, EUONYMUS** — To Control: Powdery Mildew, Red Spider, Clover Mites. Mix 2 teaspoons in 1 gallon of water. Apply as a full coverage spray when mildew or mites first appear and repeat at 7-10 day intervals.

**APPLES, PEARS** — To Control: Powdery Mildew, European Red, Clover and Two Spotted Mites. Mix 2 teaspoons in 1 gallon of water. Apply as a full coverage spray. Begin spraying at delayed dormant to pink stage. Repeat at 7-14 day intervals until mildew activity is completed. Apply when mites first appear and repeat at 7-10 day intervals. Do not graze livestock in treated orchards or feed cover crops from treated orchards to livestock. Do not use with oil based pesticides or apply to foliage within 30 days of an oil spray. Do not apply within 21 days of harvest.


**PEACHES, APRICOTS** — To Control: Powdery Mildew, European Red, Clover, Plum Rust and Peach Silver Mites. Mix 2 teaspoons in 1 gallon water. Apply as full coverage spray at pink bud, petal fall, shuck fall and early covers. Apply when mites first appear and repeat at 7-10 day intervals. Do not graze livestock in treated orchards or feed cover crops from treated orchards to livestock. Do not apply within 45 days of harvest.

**CANTALOUPE, CUCUMBERS, MELONS, PUMPKINS, SQUASH** — To Control: Powdery Mildew, European Red, Two Spotted Mites. Mix 2 teaspoons in 1 gallon of water. Apply as full coverage spray when mildew or mites first appear and repeat at 7-10 day intervals. Do not apply within 7 days of harvest. 3-10 applications may be required during the season. Cucumber, Summer Squash — Do not apply more than 4 times to any one set of fruit.

11031-1



Hacienda




# MITE-MILDEW CONTROL

leach Of Children. Use this prod-  
h label directions. Keep children  
ted areas until completely dry.  
r eyes. May cause skin irritation  
roughly after use. Avoid inhala-  
ul if swallowed. If ingested, get  
store near food or foodstuffs.

fish. Keep out of lakes, streams,  
here runoff is likely to occur. Do  
conditions favor drift from areas  
ate water by cleaning of equip-  
tes. Apply this product only as

erature will exceed 90°F during  
**FLAMMABLE! KEEP AWAY FROM HEAT**  
Buyer assumes all risk of use,  
material not in strict accordance

**ENTERPRISES**  
Charles Street  
se, Ca. 95112



**ACTIVE INGREDIENTS**  
2,4-Dinitro-6-octyl\*phenyl crotonate ..... 4.4%  
2,6-Dinitro-4-octyl\*phenyl crotonate  
Nitrooctyl\*phenols (principally dinitro) ..... 0.3%  
**INERT INGREDIENTS** ..... 95.3%

\*A mixture of 1-methylheptyl  
1-ethylhexyl and 1-propylpentyl isomers.

**ROSES, BEGONIA, BELLADONNA, DAH  
CALENDULA, DELPHINIUM, SNAPDR  
TREES and SHRUBS, LILAC, HYDRAN**

Control: Powdery Mildew, Red Spider,  
spoons in 1 gallon of water. Apply  
when mildew or mites first appear  
intervals.

**APPLES, PEARS**—To Control: Powdery  
Clover and Two Spotted Mites. Mix 2  
water. Apply as a full coverage spray  
laid dormant to pink stage. Repeat  
mildew activity is completed. Apply  
and repeat at 7-10 day intervals. Do  
treated orchards or feed cover crops  
livestock. Do not use with oil based  
foliage within 30 days of an oil spray  
days of harvest.

**PEACHES, APRICOTS**—To Control: Po  
Red, Clover, Plum Rust and Peach  
spoons in 1 gallon water. Apply as fu  
bud, petal fall, shuck fall and early  
first appear and repeat at 7-10 day  
livestock in treated orchards or feed  
orchards to livestock. Do not apply w  
**CANTALOUPE, CUCUMBERS, MELONS,**

To Control: Powdery Mildew, European  
Mix 2 teaspoons in 1 gallon of water  
spray when mildew or mites first ap  
day intervals. Do not apply within 7  
plications may be required during the  
mer Squash — Do not apply more  
set of fruit.

**NET CONTENTS: 1 PINT**