

PM 13 11037-13

10214

448234

SEP 17 1993

Mr. Don Watts  
Hacienda Enterprises  
565 Charles Street  
San Jose, CA 95112

Dear Mr. Watts:

Subject: Amendment - Change to "Caution" Formula  
Hacienda Diazinon 25%  
EPA Reg. No. 11037-13  
Your submission dated August 19, 1993

The amendment referred to above, submitted in connection with a registration under the Federal Insecticide, Fungicide, and Rodenticide Act is acceptable provided the following revisions are made prior to release for shipment under the amended labeling:

**CONTAINER LABEL:**

- 1) The front panel referral statement directs the user to the "booklet under cap". Is that really where the booklet will be? If not, revise the referral statement accordingly.
- 2) Under the heading DIRECTIONS FOR USE, capitalize the "F" of "Federal".
- 3) The referral statement under the heading "General Information" directs the user to the "booklet under cap". Is that really where the booklet will be? If not, revise the referral statement accordingly.

**BOOKLET LABEL:**

- 1) Throughout the CONDITIONS OF SALE AND WARRANTY, replace the name "CIBA-GEIGY" with the name of your company.
- 2) Under the heading DIRECTIONS FOR USE, capitalize the "F" of "Federal".
- 3) In item "5" of the "Notes" under the heading "GENERAL INFORMATION", revise "green houses" to "greenhouses".

- 4) Replace the column heading "D.z.n" with the name of your product under "Rate" in the Lawn Pest Control table.
- 5) Under the heading "Vegetables", replace "D.z.n Lawn and Garden Insect Control" with the name of your product.
- 6) For the crop "Cabbage" add the resistance limitation as follows to the "Timing/Special Directions" for use against the pests group including "Aphids" to "Dipterous Leafminers":  
  
NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides."
- 7) Delete "D.z.n trademark of CIBA-GEIGY" from beneath the Precautionary Statements block.

A stamped copy of the label is enclosed for your records. The replacement Confidential Statement of Formula (CSF) dated August 19, 1993 is acceptable and has been added to this product's registration file.

**NOTE - The label and associated Confidential Statement of Formula accepted at this time supersede all previous labels and formula statements on file for this product.**

Labeling released for shipment after the deadline imposed by the Diazinon Registration Standard, as amended, must be the revised amended label. As stipulated in the letter dated August 2, 1993, from the Agency's Special Review and Reregistration Division, products containing 25% EC formulations of diazinon may not be released after August 31, 1993 and no pesticide product containing diazinon 25% EC formulation may be distributed or sold after August 31, 1994, unless the product is the new CAUTION formula with the appropriate approved Toxicity Category III labeling.

Sincerely,

George T. LaRocca  
Product Manager (15)  
Insecticide Rodenticide Branch  
Registration Division (H7505C)

30414

ACCEPTED  
with COMMENTS  
in EPA Letter Dated

Container

SEP 17 1993

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.  
11037-13

Hacienda

# Diazinon 25%

## Lawn & Garden Insect Control

This product must not be used on golf courses and sod farms.

For outdoor use on the home lawn and the home garden only.

Spray concentrate. Contains diazinon insecticide.

**One Pint**  
Net Contents

Active Ingredient:  
Diazinon: O, O-Diethyl  
O-(2-isopropyl-6-  
methyl-4-pyrimidinyl)  
phosphoro-  
thioate . . . . . 25.0%

---

Inert  
Ingredients\* : 75.0%

---

Total: 100.0%

\* Contains aromatic petroleum distillates.

KEEP OUT OF REACH OF CHILDREN.

**CAUTION**  
See side panel for additional precautionary statements.

EPA Reg. No. 11037-13  
EPA Est. No. 11037-CA-1

Manufactured by  
HACIENDA ENTERPRISES  
565 Charles Street • San Jose, CA 95112

**Protects**  
Lawns  
Vegetables  
Fruits and Nuts  
Roses, Flowers  
Trees, Shrubs

**Controls**  
Ants, Aphids  
Chinch Bugs  
Cutworms  
Fire Ants<sup>1</sup>  
Flea Beetles  
Grubs, Mites  
Sod Webworms  
...and many others  
<sup>1</sup>Aids in control

See full label in booklet under cap for complete directions for mixing and applying this product and a complete list of pests controlled.

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.**

**General Information**

Hacienda Diazinon is an emulsifiable concentrate product which must be diluted with water before application. This product kills many outdoor pests which damage lawns, roses, flowers, trees, shrubs, vegetables and fruits and insects which annoy pets and people.

SEE THE BOOKLET UNDER CAP FOR COMPLETE DIRECTIONS FOR USE AND PESTS CONTROLLED.

Notes: (1) USE CONTENTS WITHIN 6 MONTHS AFTER OPENING. SEE STORAGE AND DISPOSAL DIRECTIONS. (2) Do not use in the home. (3) Do not use on ferns, poinsettia, hibiscus, papaya, pilea, gardenia or orchids since plant injury may occur. (4) Use this product on succulent peas and beans only. Do not use on dried pea or bean varieties. (5) Do not apply to food crops grown in greenhouses.

After spraying, wait the following number of days before harvesting fruits, nuts, or vegetables:

1 day	Tomatoes
3 days	Melons, Winter Squash
5 days	Peppers, Strawberries
7 days	Beans, Broccoli, Blackberries, Boysenberries, Cauliflower, Cucumbers, Dewberries, Loganberries, Peas, Raspberries, Summer Squash, Sweet Corn
10 days	Chinese Broccoli, Chinese Cabbage, Collards, Kale
14 days	Carrots, Endive (Escarole), Lettuce, Onions (Bulb and Green), Parsnips, Radishes, Spinach, Turnips
21 days	Apricots, Cabbage, Cherries, Nectarines, Peaches, Plums, Prunes
28 days	Grapes
35 days	Potatoes (Irish)
45 days	Walnuts
*	Almonds (one dormant spray only)

Precautions: (1) Wear a long-sleeved shirt and long-legged pants when spraying. (2) Spray with the wind to your back. (3) Do not spray on windy days. (4) If clothing becomes wet from spray during use, remove clothing after spraying, wash affected body areas thoroughly with soap and water and launder clothing before wearing again.

\* This table is only a partial listing of the pests controlled by this product. See booklet under cap for complete directions for use and pests controlled.

Problem Area	Pests*
Nuisance Pests in Outside Areas	Ants, Brown Dog Ticks, Chiggers, Cockroaches, Crickets, Fleas, Fire Ants, Spiders and others*
Lawns	Chinch Bugs, Ants, Sod Webworms, Brown Dog Ticks, Fleas, White Grubs, Lawn Billbugs and others*
<b>Vegetables: Foliar Application</b>	
Beans, Peas (except dried varieties)	Aphids, Cucumber Beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Spider Mites
Broccoli, Chinese Broccoli, Cabbage, Chinese Cabbage, Cauliflower, Collards, Kale	Aphids, Flea Beetles, Dipterous Leafminers, Diamondback Moths, Imported Cabbageworms
Endive (Escarole), Lettuce, Spinach	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites

Peppers. Tomatoes	Aonids. Colorado Potato Beetles. Fall Armyworms (Tomatoes). Flea Beetles. Leafhoppers. Southern Armyworms (Tomatoes). Vinegar Flies - <i>Drosophila</i> spp. (Tomatoes)
Onions (Sub and Green)	Onion Thrips
Radishes. Turnips. Carrots. Parsnips. Potatoes	Aonids. Flea Beetles. Leafhoppers. Dipterous Leafminers. Colorado Potato Beetles (Potatoes). Southern Armyworms (Potatoes)
Sweet Corn	Aonids. Flea Beetles. Grasshoppers. Spider Mites
Melons. Squash (Summer, Winter)	Aonids. Thrips. Spider Mites
<b>Vegetables: Preplant Soil Application</b>	
Beans. Peas (except dried varieties)	Cutworms. Mole Crickets. Wireworms (Snap Beans)
Beets. Red (Table)	Cutworms. Mole Crickets
Cabbage	Cutworms. Root Maggots. Wireworms
Carrots. Sweet Corn	Cutworms. Mole Crickets. Wireworms
Celery. Tomatoes*	Cutworms. Wireworms
Radishes. Turnips	Cutworms
<b>Fruits &amp; Nuts</b>	
Almonds (California only)	Scale Crawlers
Walnuts (California only)	Aonids. Scale Crawlers. Spider Mites and others*
Apricots. Peaches. Nectarines	Aonids. Apricot Mealybugs (Apricots). Spider Mites. Scale Crawlers
Cherries	Black Cherry Aonids. Cherry Fruitflies. Cherry Rust Mites Scale Crawlers and others*
Plums. Prunes	Aonids. Spider Mites
Caneberries (CA, OR, WA only) Blackberries. Boysenberries. Dewberries. Loganberries. Raspberries	Aonids. Leafhoppers. Two-Spotted Spider Mites. Thrips. Raspberry Fruitworms
Strawberries	Aonids. Two-Spotted Spider Mites. Strawberry Leafrollers
Grapes	Aonids. Grape Berry Moths. Grape Leafrollers. Leafhoppers
<b>Roses &amp; Flowers</b>	
Azaleas. Camellias. Chrysanthemums. Carnations. Gladiol. Lilac. Rhododendron. Roses	Aonids. Flea Beetles. Dipterous Leafminers. Leafhoppers. Mealybugs. Two-spotted Spider Mites. Thrips. Whiteflies and others*

**Ornamental Trees & Shrubs**

Arborvitae. Aohios. Mites. Scale Crawlers. Whiteflies. Tent Caterpillars and others\*  
 Azaleas. Birch.  
 Boxwood. Camemias.  
 Douglas Fir. Elm.  
 Hawthorn Holly.  
 Juniper. Lilac.  
 Locust. Maple.  
 Mimosa. Oak. Pine.  
 Ornamental Plum.  
 Poplar.  
 Rhododendron.  
 Soruce. Willow

**Storage and Disposal****Storage**

Do not store near heat or open flame. Store in original container in an area inaccessible to children or pets. **USE CONTENTS WITHIN 6 MONTHS AFTER OPENING.**

**Disposal**

Do not reuse empty container. Thoroughly rinse container, securely wrap in several layers of newspaper, and discard in trash. If disposal of a partially full container is necessary, make sure cap is tightly closed, securely wrap the container in several layers of newspaper and discard in trash.

**Precautionary Statements****Hazards to Humans and Domestic Animals****CAUTION**

Harmful if swallowed, inhaled, or absorbed through skin. Causes moderate eye injury. Avoid breathing spray mist. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not use inside the home. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not allow children or pets to go onto a treated area until the treated area, such as grass or soil, has dried. Do not contaminate ornamental fish ponds. Food utensils such as spoons and measuring cups must not be used for food purposes after use in measuring pesticides.

**Statement of Practical Treatment**

**If swallowed:** Call a physician immediately. **DO NOT INDUCE VOMITING UNLESS NO OTHER TREATMENT IS AVAILABLE.** Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

**If inhaled:** Remove victim to fresh air and apply artificial respiration if indicated. Get medical attention.

**If on skin:** Wash promptly with soap and water. Rinse thoroughly. Get medical attention.

**If in eyes:** Rinse eyes with plenty of water and call a physician immediately.

**Note to Physician:** This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. This product contains aromatic petroleum

distillates which may present an aspiration hazard. Gastric lavage may be indicated if product was taken internally.

**Environmental Hazards**

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays for lawns, where irrigation (watering) is recommended in the directions, or if waterfowl, i.e., ducks or geese, can be expected in the treated area after treatment (except spot treatment), apply at least 1/4 inch of water immediately after spraying this product, however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

**Physical or Chemical Hazards**

Do not use, pour, spill, or store near heat or open flame.

**BEST AVAILABLE COPY**

7044

ACCEPTED  
with COMMENTS  
in EPA Letter Dated

Booklet

SEP 17 1993

~~Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.  
11037-13~~

*Hacienda*

# Diazinon 25%

## Lawn & Garden Insect Control

This product must not be used on golf courses and sod farms.

For outdoor use on the home lawn and the home garden only.

Spray concentrate. Contains diazinon insecticide.

**One Pint  
Net Contents**

Active Ingredient:  
Diazinon: O, O-Diethyl  
O-(2-isopropyl-6-  
methyl-4-pyrimidinyl)  
phosphoro-  
thioate . . . . . 25.0%

Inert  
Ingredients \* : 75.0%

Total: 100.0%

\* Contains aromatic petroleum distillates.

KEEP OUT OF REACH OF CHILDREN.

### CAUTION

See end of booklet for additional precautionary statements

EPA Reg. No. 11037-13  
EPA Est. No. 11037-CA-1

Manufactured by  
HACIENDA ENTERPRISES  
565 Charles Street • San Jose, CA 95112

**Protects**  
Lawns  
Vegetables  
Fruits and Nuts  
Roses, Flowers  
Trees, Shrubs

**Controls**  
Ants, Aphids  
Chinch Bugs  
Cutworms  
Fire Ants<sup>1</sup>  
Flea Beetles  
Grubs, Mites  
Sod Webworms  
...and many others

<sup>1</sup>Aids in control

**DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY**

**IMPORTANT:** Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product. If terms are not acceptable, return the unopened product container at once.

**Conditions of Sale and Warranty**

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of CIBA-GEIGY or the Seller. All such risks shall be assumed by the Buyer.

CIBA-GEIGY warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. CIBA-GEIGY makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall CIBA-GEIGY or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product.

CIBA-GEIGY and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of CIBA-GEIGY.

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.**

**General Information**

Hacienda Diazinon is an emulsifiable concentrate product which must be diluted with water before application. Refer to the section How To Mix and Apply. This product kills many outdoor insects which damage lawns, roses, flowers, trees, shrubs, vegetables fruits and nuts, and insects which annoy pets and people. Contains diazinon insecticide which provides positive control of many "hard-to-kill" sucking and chewing insects. Diazinon works fast and applications should be repeated only as directed to maintain control.

**Notes:** (1) USE CONTENTS WITHIN 6 MONTHS AFTER OPENING. SEE STORAGE AND DISPOSAL DIRECTIONS. (2) Do not use in the home. (3) Do not use on ferns, poinsettia, hibiscus, papaya, pine, geranium or orchids since injury may occur. (4) Use this product on succulent peas and beans only. Do not use on dried pea or bean varieties. (5) Do not apply to food crops grown in green houses.

**Precautions:** (1) Wear a long-sleeved shirt and long-legged pants when spraying. (2) Spray with the wind to your back. (3) Do not spray on windy days. (4) If clothing becomes wet in spray during use, remove clothing after spraying, wash affected body areas in soap and water, and launder clothing before wearing again.

**How to Mix and Apply**

**Step #1:** Determine size of spraying job. For lawns and vegetables, measure the area to be sprayed. For all other applications, apply as a thorough cover spray. Make applications when insects first appear. Applications should be repeated only as directed to maintain control.

**Step #2:** Use clean sprayer. Carefully measure and mix the amount of Diazinon and water as suggested in the rate chart.

**Step #3:** Spray as directed in rate chart. Thorough coverage is important. Flush sprayer with clean water after each use.

See chart for complete list of use sites, crops, and insects controlled. Use the rate given in the chart for the type of insect you want to control on the site or crop you are spraying.

**Measurement Conversions**

- 1 tablespoon = 3 teaspoons
- 1 fl. oz. = 6 teaspoons or 2 tablespoons
- 8 fl. oz. = 1 cup

- 1 1/2 tablespoons in 3 gals. of water = 1/2 teaspoons in 1 gal. of water
- 2 tablespoons in 3 gals. of water = 2 teaspoons in 1 gal. of water

**Lawn Pest Control**

**Precautions:** Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Apply this product only as specified on the label. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid over-

lapping of sprays. Where irrigation (watering) is recommended in the directions, or if waterfowl, i.e., ducks or geese, can be expected in the treated area after treatment (except spot treatment), water lawn with at least 1/4 inch of water immediately after spraying this product; however, stop watering before puddling occurs.

Use on Bent, Bermuda, Bluegrass, Dichondra, Fescue, Mennon and St. Augustinegrass.

To control pests listed below, apply recommended rates.

Pests	Rate		Remarks
	D+zin	Water	
Lawn Chinch Bugs	1 1/2 tablespoons   per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals.  24 gals.	Thoroughly wet down grass a few hours before applying.  In the North, if Chinch Bugs are the only problem, a lower rate of application can be used: 3 teaspoons in 3 gals. of water covers 125 sq. ft. or 4 fl. oz. in 24 gals. of water per 1,000 sq. ft.
Ants, Armyworm, Bermudagrass Mites, Clover	1 1/2 tablespoons   per 125 sq. ft. or	3 gals.	Thoroughly wet down grass a few hours before applying.
Mites, Springtails (Collembola), Crickets, Cutworms, Digger Wasps, Earwigs, Frit Flies, Lawn Billbugs, Leathoppers, Sod Webworms (Lawn Moths), Sowbugs (Pillbugs)	6 fl. oz. per 1,000 sq. ft.	24 gals.	For ants and digger wasps, spot spray ant mites and wasp ground nest openings. For digger wasps, apply at dusk, when wasps are not active.  Billbug control: Apply when activity is first observed or when chewed or brown grass is present indicating damage from the insect.  For frit flies, mow grass and water thoroughly before treatment, but delay application until grass is dry.
Brown Dog Ticks, Chiggers, Fleas	1 1/2 tablespoons   per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals.  24 gals.	NOTE: Do not apply the product to animals.  For brown dog ticks, spray grass and under shrubber; preferably near house.
Hyperoos Weevils	1 1/2 tablespoons   per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals.  24 gals.	Treat problem areas in mid-April and again in mid-May. Water grass thoroughly after application.
Milkweed	1 1/2 tablespoons   per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals.  24 gals.	Thoroughly wet grass a few hours before applying.
Rhodesgrass Scapes	1 1/2 tablespoons   per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals.  24 gals.	Apply when crawlers first emerge.
White Grubs of Japanese Beetle, European Chaffer, Southern Chaler	1 1/2 tablespoons   per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals.  24 gals.	Apply early in the season, late July and early October, after grass has begun to grow.

**BEST AVAILABLE COPY**

Nuisance Pests Outside the Home

Ants, Boxelder Bugs, Brown Dog Ticks, Chiggers, Clover Mites, Cockroaches, Crickets, Earwigs, Fleas, Flies, Millipedes, Sowbugs (Pillbugs), Spiders, Springtails (Collembola)	Mix 6 fl. oz. in 3 gals. of water and thoroughly spray window and door screens, sills, foundations, porch and patio walls, entranceways, walls, garbage cans, tree trunks, and into cracks and other places where insects can hide.  To prevent entry into houses, spray a five-foot band of soil around the house next to the foundation and spray the foundation of the house to a height of 2-3 feet. Repeat application as necessary.  Do not treat animals with this product. Do not treat plants (other than tree trunks) at this nuisance pest rate.
---	---

Fire Ants	To aid in the control of fire ants in home lawns and other outdoor home recreation areas, apply 2 tablespoons in a minimum of 1 gal. of water as a drench. Apply over and 6 inches around each mound. Apply gently to avoid disturbing ants. Reapply as necessary. Larger quantities of solution can be prepared for multiple mound treatments. Use equipment capable of delivering the diluted product as a gentle rain at the rate of 1 gal. per mound. High pressure sprays may disturb the ants and cause migration, reducing product effectiveness.  NOTE: For best results, apply in cool weather 65-80°F or in early morning or late evening hours. Treat new mounds as they appear.
-----------	---

Insect Control on Ornamental Trees, Shrubs and Flowers

Tree and Shrubs	Insects	Rate per 3 Gal. of Water	Timing/Special Directions
Arborvitae, Azaleas, Birch, Boxwood, Camellias, Douglas Fir, Elm, Hawthorn, Holly, Juniper, Lilac, Locust, Maple, Mimosa, Oak, Pine, Ornamental Plum, Potted Rhododendron, Rose, Willow	Aphids, Bagworms, European Pine Shoot Moths, Holly Bud Moths, Dipterous Leafminers (Arborvitae, Azalea, Birch, Boxwood, Holly, Oak), Leafhoppers, Mealybugs, Mites (Clover, Cyclamen, European Red, Privet, Two-spotted Spider), Obscure Root Weevils, Omnivorous Leaf-tiers, Scale Crawlers (Cottoncushion), Euonymus, Lecanium, Pine Needle, Pt. San Jose, Soft	2 table- spoons	Apply as a thorough foliar spray. Wet leaves and branches to the drooping point. Try to penetrate dense foliage with spray.
	Webworms (Juniper, Mimosa), Whiteflies	2 table- spoons	Apply as a thorough foliar spray. Mimosa Webworms: apply as a thorough foliar spray when adults first appear. Make two additional applications at 4-5 week intervals.
	Apple-and-Thorn Skeletonizers, Hemlock Chermes, Oak Looper, Obliquebanded Leafhopper, Pear Slugs, Tent Caterpillars, Webworms (Cottoncushion, Fall)	2 table- spoons	Apply as a thorough foliar spray.

Roses and Flowers	Insects	Rate per 3 Gal. of Water	Timing/Special Directions
Azaleas, Camellias, Chrysanthemums, Carnations, Gladiol, Lilac, Rhododendron, Roses	Aphids, Euonymus Scale Crawlers, Flea Beetles, Dipterous Leafminers, Leafhoppers, Mealybugs, Mites (Carnation Bud, Carnation Shoot, Cyclamen, European Red, Two-spotted Spider), Thrips, Whiteflies	2 table- spoons	Apply as a thorough foliar spray. Try to spray underside of leaves and penetrate dense foliage.

Fruits and Nuts

Start spraying after blooming. Applications should be made according to infestation or when aphids, mites, and crawler stages of scale insects are first noted.

Observe stated time intervals between last application and harvest of the crop, as well as intervals between applications and total number of applications that can be made per season.

Crop	PHI*
Almonds	**
Apples	21
Blackberries	7
Boysenberries	7
Cherries	21
Dewberries	7
Grapes	28
Loganberries	7
Nectarines	21
Peaches	21
Plums	21
Prunes	21
Raspberries	7
Strawberries	7
Walnuts	15

\* PHI = Pre-Harvest Interval, or the number of days between the last application and harvest.

\*\* Dormant spray only

Note: To protect bees, do not apply to fruit trees when trees or substantial numbers of weeds around the trees are in bloom

Crop	Pest	Rate	Timing/Special Directions
Almonds (CA only)	San Jose scale, Pearstone scale	2 tablespoons in 3 gal. of water	Apply only as a dormant spray. Do not apply more than 6 gallons per 500 sq. ft. of orchard.  This product may be used on almonds only in CA. Do not use this product on almonds grown in other states.

<p>Aproncos</p>	<p>Aphids. Apricot Mealybugs. Mites (Brown. Two-spotted Spider). Scale Crawlers (Olive, San Jose)</p>	<p>4 teaspoons in 2 gals. of water</p>	<p>Apply as a full cover spray. Do not apply more than 4 gallons per 500 sq. ft. of orchard. Do not make more than 3 applica- tions per season a minimum of 14 days apart. Do not apply within 21 days of harvest.</p>
<p>Cane- bermes (CA, OR, WA only); Black- bermes. Boysen- bermes. Caw- bermes. Logan- bermes. Raso- bermes</p>	<p>Aphids. Leafhoppers. Raspberry Fruitworms. Two-spotted Spider Mites. Thrips</p>	<p>2 tablespoons in 3 gals. of water</p>	<p>Apply as a full cover spray. Do not apply more than 5 gallons per 500 sq. ft. of field. Do not make more than 5 applications per season a minimum of 14 days apart. Do not apply within 7 days of harvest.  This product may be used on caneberries only in CA, OR and WA. Do not use this product on caneber- ries grown in other states.</p>
<p>Cherries</p>	<p>Black Cherry Aphids Cherry Fruitflies Cherry Fruitworms (Maggots). Fruitree Leafrollers Mites (Cherry Rust). Scale Crawlers (San Jose)</p>	<p>4 teaspoons in 2 gals. of water</p>	<p>Apply as a full cover spray. Do not apply more than 4 gallons per 500 sq. ft. of orchard. Do not make more than 3 applica- tions per season a minimum of 7 days apart. Do not apply within 21 days of harvest.</p>
<p>Grapes</p>	<p>Aphids. Leafhoppers. Grape Berry Moths Grape Leafrollers</p>	<p>4 1/2 teaspoons in 3 gals. of water</p>	<p>Apply as a full cover spray. Do not apply more than 3 gallons per 500 sq. ft. of vineyard. Do not make more than 5 applica- tions per season a minimum of 7 days apart. Do not apply within 28 days of harvest.</p>
<p>Strawberries</p>	<p>Aphids. Mites (Brown Two-spotted Spider). Scale Crawlers (Olive, San Jose)</p>	<p>4 teaspoons in 2 gals. of water</p>	<p>Apply as a full cover spray. Do not apply more than 4 gallons per 500 sq. ft. of orchard. Do not make more than 3 applica- tions per season a minimum of 14 days apart. Do not apply within 21 days of harvest.</p>
<p>Peaches</p>	<p>Aphids. Mites (Brown. Two-spotted Spider). Scale Crawlers (Olive, San Jose)</p>	<p>4 teaspoons in 2 gals. of water</p>	<p>Apply as a full cover spray. Do not apply more than 4 gallons per 500 sq. ft. of orchard. Do not make more than 3 applica- tions per season a minimum of 14 days apart. Do not apply within 21 days of harvest.</p>

<p>Plums and Prunes</p>	<p>Aphids (Leatcut Plum. Mealy Plum, Thistle). Mites (Brown. European Red)</p>	<p>4 teaspoons in 2 gals. of water</p>	<p>Apply as a full cover spray. Do not apply more than 4 gallons per 500 sq. ft. of orchard. Do not make more than 3 applica- tions per season a minimum of 7 days apart. Do not apply within 21 days of harvest.</p>
<p>Straw- berries</p>	<p>Aphids. Two-spotted Spider Mites. Strawberry Leafrollers</p>	<p>2 tablespoons in 3 gals. of water</p>	<p>Apply as a full cover spray. Do not apply more than 3 gallons per 500 sq. ft. of field. Do not make more than 3 applications per season a minimum of 7 days apart. Do not apply within 5 days of harvest.</p>
<p>Walnuts (CA only)</p>	<p>Aphids. Scale Crawlers (Walnut, Italian Pear, Frested). Spider Mites. Twig Borers</p>	<p>2 tablespoons in 3 gals. of water</p>	<p>Apply as a full cover spray. Do not apply more than 6 gallons per 500 sq. ft. of orchard. Do not make more than 3 applica- tions per season 14 days apart. Do not apply after husks open or split. Do not apply within 45 days of harvest.  This product may be used on walnuts only in CA. Do not use this product on walnuts grown in other states.</p>

BEST AVAILABLE COPY

**Vegetables**

For foliar or soil applications, Diazinon Lawn and Garden Insect Control should be applied in the volume of water recommended in the table below to assure thorough coverage of the foliage or soil.

Observe stated time intervals between last application and harvest as well as intervals between applications and maximum numbers of applications per season. Do not apply to food crops grown in greenhouses.

**Pre-Harvest Interval (PHI\*) for Foliar Applications**

Crop	PHI* (Days)
Beans <sup>1</sup> (succulent varieties only: Lima, Pole and Snap)	7
Broccoli	7
Cabbage	21
Cantaloupes	3
Carrots	14
Cassabas	3
Cauliflower	7
Chinese Broccoli, Chinese Cabbage, Collards	10
Crenshaw	3
Cucumbers	7
Escarole	14
Honeydew Melons	3
Kale	10
Lettuce	14
Muskmelons	3
Onions (Bulb and Green), Parsnips	14
Peas <sup>1</sup>	7
Peppers	5
Potatoes (Irish)	35
Radishes, Spinach	14
Squash, Summer	7
Squash, Winter	3
Sweet Corn	7
Tomatoes	1
Turnips (Roots and Tops)	14
Watermelons	3

\* PHI = Pre-Harvest Interval or the number of days between the last application and harvest.

<sup>1</sup> Do not use this product on dried pea or bean varieties such as pinto beans, dried limas, split peas or dried black-eyed peas.

**Vegetables**

For application to the foliage, thoroughly spray all plant parts. Application to the soil should be mixed into the soil immediately after application.

Crop	Pest	Rate	Timing/Special Directions
Beans, Lima, Pole and Snap (succulent varieties only)	Aphids, Cucumber beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 5 days up to 3 applications per season. Do not apply within 7 days of harvest.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil 1-2 days before planting. Immediately work into soil 1-2 inches.
	Wireworms (snap beans only)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 4-6 inches.

NOTES: (1) Do not apply this product to dried bean varieties such as pinto beans or dried limas. (2) To protect bees, do not apply bean foliage treatments if the beans or weeds in the treatment area are in bloom.

Beets, Red (Table)	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil 1-2 days before planting. Immediately work into soil 1-2 inches.
Broccoli	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest.
			NOTE: Repeat applications will not be controlled by diazinon or similar type insecticides.

Cabbage	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 21 days of harvest.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Root Maggots	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 4-6 inches.

**BEST AVAILABLE COPY**

Carrots	Aphids. Flea Beetles. Leafhoppers. Dipterous Leafminers. Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 14 days of harvest.  NOTE: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecti- cides.
Cutworms (surface, subterranean)		6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for sur- face cutworms. 3-6 inches for subterra- nean cutworms.
Mole Crickets		2 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil 1-2 days before plant- ing. Immediately work into soil 1-2 inches.
Wireworms		6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 4-8 inches
Cauliflower	Aphids Diamondback Moths Flea Beetles. Imported Cabbageworms Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 7 days of harvest.  NOTE: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecti- cides.
Chinese Broccoli	Aphids Diamondback Moths Flea Beetles Imported Cabbageworms Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 10 days of harvest.  NOTE: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecti- cides.
Chinese Cabbage (Box Choy and Napa)	Aphids. Diamondback Moths Flea Beetles Imported Cabbageworms Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 10 days of harvest.  NOTE: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecti- cides.
Collards	Aphids. Diamondback Moths. Flea Beetles Imported Cabbageworms Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden.	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 10 days of harvest.  NOTE: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecti- cides.

Corn (see sweet corn)			
Cucumbers	Aphids. Thrips. Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 7 days of harvest.
Endive (Escarole)	Aphids. Flea Beetles. Leafhoppers. Dipterous Leafminers. Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 14 days of harvest.  NOTE: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecti- cides.
Kale	Aphids. Diamondback Moths Flea Beetles Imported Cabbageworms Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 10 days of harvest.  NOTE: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecti- cides.
Lettuce	Aphids. Flea Beetles Leafhoppers. Dipterous Leafminers. Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 14 days of harvest.  NOTE: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecti- cides.
Cutworms (surface, subterranean)		6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for sur- face cutworms. 3-6 inches for subterra- nean cutworms.
Wireworms		6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 4-8 inches.
Melons (Cantaloupe, Casabas, Cren- snaws Honey- dews, Must- melons, Persians and Hybrids of these) and Water- melons	Aphids. Thrips. Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 3 days of harvest.

**BEST AVAILABLE COPY**

Onions (Bulb and Green)	Onion Thrips	2 tablespoons in 3 gals. of water to cover 1 000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest.
Peas (succulent only)	Aphids. Flea Beetles. Leafhoppers. Dipterous Leafminers. Spider Mites	2 tablespoons in 3 gals. of water to cover 1 000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat applications every 7 days up to 5 applications per season. Do not apply within 14 days of harvest.  NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms. 3-6 inches for subterranean cutworms.
Peas	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil 1-2 days before planting. Immediately work into soil 1-2 inches.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 4-8 inches.
NOTES. (1) Do not apply this product to dried pea varieties such as split peas or dried black-eyed peas. (2) To protect bees, do not apply foliage treatments if bees or weeds in the treatment area are in bloom.			
Peas	Aphids. Colorado Potato Beetles. Flea Beetles. Leafhoppers	2 tablespoons in 3 gals. of water to cover 1 000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 5 days of harvest.
Potatoes	Aphids. Flea Beetles. Leafhoppers. Colorado Potato Beetles. Southern Armyworms. Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1 000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 35 days of harvest.  NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.

Radishes	Aphids. Flea Beetles. Leafhoppers. Imponed Cabbageworms. Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1 000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 3 applications per season. Do not apply within 14 days of harvest.  NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Squash Summer	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms. 3-6 inches for subterranean cutworms.
	Aphids. Thrips. Spider Mites	2 tablespoons in 3 gals. of water to cover 1 000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest.
Squash Winter	Aphids. Thrips Spider Mites	2 tablespoons in 3 gals. of water to cover 1 000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 3 days of harvest.
Sweet Corn	Aphids, Thrips Flea Beetles Grasshoppers Spider Mites	2 tablespoons in 3 gals. of water to cover 1 000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Corn may be treated 7 days following last application.
Sweet Corn	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms. 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil 1-2 days before planting. Immediately work into soil 1-2 inches.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 4-8 inches.
NOTE: To protect bees, do not apply to corn foliage during the pollen shed period.			

Tomatoes	Aphids. Colorado Potato Beetles. Fall Armyworms. Flea Beetles. Leafhoppers. Southern Armyworms. Vinegar Flies (Drosophila spp.)	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 1 day of harvest.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms. 3-6 inches for subterranean cutworms.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 4-8 inches.
Turnips and Turnip Tops	Aphids. Flea Beetles. Leafhoppers. Imperial Cabbageworms Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest.  NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms. 3-6 inches for subterranean cutworms

## Storage and Disposal

### Storage

Do not store near heat or open flame. Store in original container in an area inaccessible to children and pets. **USE CONTENTS WITHIN 6 MONTHS AFTER OPENING.**

### Disposal

Do not reuse empty container. Thoroughly rinse container, securely wrap it in several layers of newspaper and discard in trash. If disposal of a partially full container is necessary, make sure cap is tightly closed, securely wrap the container in several layers of newspaper and discard in trash.

## Precautionary Statements

### Hazards to Humans and Domestic Animals

#### CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Causes moderate eye injury. Avoid breathing spray mist. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not use inside the home. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not allow children or pets to go onto a treated area until the treated area, such as grass or soil, has dried. Do not contaminate ornamental fish ponds. Food utensils such as spoons and measuring cups must not be used for food purposes after use in measuring pesticides.

### Statement of Practical Treatment

If swallowed: Call a physician immediately. **DO NOT INDUCE VOMITING UNLESS NO OTHER TREATMENT IS AVAILABLE.** Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

If inhaled: Remove victim to fresh air and apply artificial respiration if indicated. Get medical attention.

If on skin: Wash promptly with soap and water. Rinse thoroughly. Get medical attention.

If in eyes: Rinse eyes with plenty of water and call a physician immediately.

Note to Physician: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. This product contains aromatic petroleum distillates which may present an aspiration hazard. Gastric lavage may be indicated if product was taken internally.

### Environmental Hazards

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. A C.C. overlapping of sprays. For lawns, where irrigation (watering) is recommended in the direction of waterfowl, i.e., duck or geese, can be expected in the treated area after treatment (except spot treatment), water lawn with at least 1/2 inch of water immediately after spraying this product, however, stop watering before budding occurs. Keep off lakes, streams, ponds, bogal marshes and estuaries. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on the label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

### Physical or Chemical Hazards

Do not use, pour, spill, or store near heat or open flame

© 1984, Bayer Crop Science, a Division of Bayer Corporation

Hacienda Enterprises  
565 Charles Street  
San Jose, CA 95112

**BEST AVAILABLE COPY**