CENTER PANEL U LIQUID COPPER FUNGICIDE SPRAY

> H A C I E N D A LIQUID COPPER FUNGICIDE SPRAY

A FIXED COPPER SPRAY

ACTIVE INGREDIENTS Copper Oleate*	15.00%
INERT INGREDIENTS	85.00%
*Copper expressed as elemental Contains petroleum distillates	1.34%

KEEP OUT OF REACH OF CHILDREN CAUTION SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS NET CONTENTS

ACCEPTED with COMMENTS in EPA Letter Dated

APR 3 0 1984

Under the Federal Insocticide, Fungicide, and Rodenticide Act as amended, for the postregistered under EVA I. //037-/5

EPA Reg. No. 11037-15 EPA Est. 10 7-CA-1

143

a an a ar ar aga a

•

:

ŧ

-- -

LIQUID COPPER FUNGICIDE SPRAY RIGHT PANEL

DIRECTIONS FOR USE

SHADE TREES, FLOWERS, SHRUBS; (Azaleas, Begonias, Camellias, Chrysanthemums, Crepe Myrtle, Dahlias, Delphinium, Fuchsias, Roses, Snapdragons, Stocks, Sweet Peas) - Black Spot of Rose, Leaf Spots, Powdery Mildew, Rust: Mix 1 Tbsp. per gallon water. Spray when foliage or disease appear. Repeat at 7-10 day intervals or as necessary.

APPLES, PEARS - Powdery Mildew: Mix 1 Tbsp. per gallon water. Spray when buds swell. Repeat when fruit buds show color and again at petal fall.

APRICOTS - Shothole: Mix 1 Tsp. per gallon or water. Apply in delayed dormant period (during January in West Coast areas). Repeat at 7-10 day intervals as necessary.

NECTARINES, PEACHES, PLUMS - Leaf Blight, Leaf Curl, Powdery Mildew, Shothole: Mix 1 Tbsp. per gallon water. Apply in late fall (Nov.15-Dec.15 in West Coast areas). Repeat when buds begin to swell in the spring. Repeat at 7-10 day intervals as necessary.

WALNUTS - Leaf Spots: Mix 1 Tbsp. per gallon water. Spray after blooms drop and nut is well set. Repeat at 2 week intervals as necessary.

CITRUS ~ Blue Mold, Brown Rot: Mix 3 Tbsp. per gallon water. Spray lower part of tree during winter months. Sooty Mold: Mix 1 Tbsp. per gallon water. Spray when disease appears and repeat as necessary.

BLACKBERRIES, RASPBERRIES, STRAWBERRIES - Leaf Spots, Powdery Mildew: Mix $1-1\frac{1}{2}$ Tbsp. per gallon water. Spray when disease appears and repeat as necessary.

TOMATOES - Powdery Mildew: Mix 1 Tbsp. per gallon water. Spray before disease appears. Repeat at 7-10 day intervals as necessary.

BEANS - Powdery Mildew, Rust: Mix 12 Tbsp. per gallon water. Spray before disease appears. Repeat at 7-10 day intervals as necessary.

CUCUMBERS - Powdery Mildew: Mix $1\frac{1}{2}$ Tbsp. per gallon water. Begin when foliage appears. Repeat at 7-10 day intervals as necessary.

MELONS; Cantaloupe, Cassaba, Honey Ball, Honeydew, Muskmelon, Persian Melon, Watermelon - Powdery Mildew: Mix 1¹/₂ Tbsp. per gallon water. Spray before disease appears. Repeat at 7-10 intervals as necessary.

HACIENDA ENTERPRISES 565 Charles Street, San Jose, CA 95112

EPA Reg. No. 11037-15

11037-15

•

ŧ

ł

LIQUID COPPER FUNGICIDE SPRAY LEFT PANEL

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION: Harmful if swallowed. Do not breathe spray mist. Wash thoroughly after using.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label.

(3)

PHYSICAL OR CHEMICAL HAZARDS Do not use, pour, spill or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistant with its labeling.

Do not reuse container. Wrap partially filled container in several layers of newspaper, or rinse empty container before discarding in trash.

्_र)

EPA Reg: No. 11037-15) EPA Est. 11037-CA-1

- - - - ----

301 - 119612. 17 1 APR 3 0 1984 Hacienda Enterprises 565 Charles Street San Jose, CA 95112 Attention: D. L. Watts Gentlemen: Subject: Hacienda Liquid Copper Fungicide Spray EPA Registration No. 11037-15 Your Letter dated Harch 2, 1984 The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, is acceptable subject to the comments listed below. Five copies of the finished labeling must be submitted. 1. Insert the heading "Storage and Disposal" above pertinent information utilizing "Storage:" and "Disposal:" subheadings. Include information pertaining to product storage. 2. Change "Active Ingredients" to the singular "Active Ingredient." A stamped copy of the label is enclosed for your records. Sincerely yours, Henry M. Jacoby Product Hanager (21) Fungicide-Herbicide Branch Registration Division (TS~767) Enclosue PR Notice 83-3 RD-DIS:DCR-03271:Jacoby:GG:Raven: 57-2226:RD-48:4/19/84/Del.5/3/84

1

CONCURRENCES									
SYMBOL .									
AURHAME					***********				

