

Sanitizing of Food Processing Equipment and other Hard Non-Porous Surfaces in Food Contact Locations.

For sanitizing food processing equipment, dairy equipment, food utensils, dishes, silverware, glasses, sink tops, counter tops, refrigerated storage and display equipment and other hard non-porous surfaces.

Wash and rinse all articles thoroughly, then apply a solution of 1 oz. of cleaner to 4 gallons of water (200 ppm active). Surfaces should remain wet for at least one minute followed by adequate draining and air drying. Fresh solution should be prepared daily or when use solution becomes visibly dirty. For mechanical application, use solution may not be re-used for sanitizing applications.

Dishes, silverware, glasses, cooking utensils and other similar size food processing equipment can be sanitized by immersing in a 1 oz. to 4 gallon dilution of cleaner. No Potable water rinse is required.

At 1 oz. to 4 gallon dilution, this cleaner has a hard water tolerance of 750 ppm of hardness calculated as CaCO₃ when evaluated by the AOAC Germicidal and Detergent Sanitizer Method against Escherichia coli and Staphylococcus aureus.

The udders, flanks, and teats of dairy cows can be sanitized by washing with a solution of 1 oz. of cleaner in 4 gallons of warm water.

Use a fresh towel for each cow. Avoid contamination of sanitizing solution by dirt and soil. Do not dip used towel back into sanitizing solution. When solution becomes visibly dirty, discard and provide fresh solution.

**PRECAUTIONARY STATEMENTS • Hazards to Humans and Domestic Animals
DANGER**

Keep out of reach of children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL.
OPEN DUMPING IS PROHIBITED
DO NOT REUSE EMPTY CONTAINER

PESTICIDE DISPOSAL—Pesticide or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

CONTAINER DISPOSAL—Triple rinse (or equivalent) and dispose in an incinerator or landfill approved for pesticide containers, or bury in a safe place

GENERAL—Consult federal, State or Local disposal authorities for approved alternative procedures such as limited open burning.

ACCEPTED
DEC 3 0 1980
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under 11082-7 EPA Reg. No.

**LONG
CHEMICAL
INC.**

AG 99

DISINF
SANIT
FUNGI
DEOD

**DISINFECTANT • SANITIZER • FUNGICIDE • DEODORIZER
HOSPITAL, INSTITUTIONAL, INDUSTRIAL, SCHOOL,
AND OTHER FARM AND HOME USE**

ACTIVE INGREDIENTS:

- Axyl (C14, 60%; C16, 30%; C12, 5%; C18, 5%)
- Dimethyl Benzyl Ammonium Chlorides
- Axyl (C12, 68%; C14, 32%)
- Dimethyl Ethylbenzyl Ammonium Chlorides

INERT INGREDIENTS

**KEEP OUT OF REACH OF CHILDREN
DANGER
Statement of Practical Treatment**

In case of contact, immediately flush eyes or skin with plenty of water for 15 minutes. For eyes, call a physician, remove and wash contact lenses, and remove clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg white solution; or if these are not available, drink large quantities of water or alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

40 NET 1 GALLON

BEST DOCUMENT AVAILABLE

AA AG 99

DISINFECTANT
SANITIZER
FUNGICIDE
DEODORIZER

SANITIZER • FUNGICIDE • DEODORIZER FOR
INSTITUTIONAL, INDUSTRIAL, SCHOOL, DAIRY
OTHER FARM AND HOME USE

CONTENTS:

10% (12.5% ClO 5%)	5.0%
Sodium Chloride	
(12%)	
1 Ammonium Chloride	5.0%
	90.0%
	100.0%

KEEP OUT OF REACH OF CHILDREN

DANGER

Prevention of Practical Treatment

Immediately flush eyes or skin with plenty of water for at least 15 minutes. Call a physician. Remove and wash contaminated clothing.

Apply a large quantity of milk, egg whites, gelatin or other food if available. Drink large quantities of water. Avoid further exposure.

Obvious mucous membrane damage may contraindicate the use of this product against circulatory shock, respiratory depression, or other conditions. See first aid section.

SEE ADDITIONAL PRECAUTIONARY STATEMENTS

NET 1 GALLON

DIRECTIONS FOR USE • General Classification

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Apply cleaner with a cloth, mop or mechanical spray device. When applied with a mechanical spray device, surface must remain wet for 10 minutes. Fresh solution should be prepared daily or when the use solution becomes visibly dirty.

Disinfection in Hospitals, Nursing Homes and Other Health Care Institutions: For disinfecting floors, walls, countertops, bathing areas, lavatories, bedframes, tables, chairs, garbage pails and other hard non-porous surfaces add 4 1/2 oz. of cleaner to 5 gallons of water. Apply to previously cleaned hard non-porous surface with mop or cloth. At this level the solution is effective against *Pseudomonas aeruginosa*.

Disinfectant in Institutions, Industry, Schools and Homes: For disinfecting floors, walls, bedframes, countertops, tables, chairs, garbage pails, bathroom fixtures and other hard non-porous surfaces add 3 oz. of cleaner to 5 gallons of water. Apply to previously cleaned hard non-porous surface with mop or cloth. At 3 oz./5 gallon use-level, this solution is effective against *Staphylococcus aureus*, and *Salmonella choleraesuis*.

Fungicidal Activity: At 8 oz./5 gallons use-level, this cleaner is effective against *Trichophyton interdigitale* (athlete's foot fungus) on inanimate hard non-porous surfaces.

Disinfection of Pre-Cleaned Barber Tools: Barber tools (such as combs, brushes, razors, and scissors) can be disinfected by immersing in a .6 oz./gallon solution.

Disinfection of Poultry Equipment, Animal Quarters and Kennels: Poultry brooders, watering fountains, feeding equipment and other animal quarters (such as stalls and kennel areas) can be disinfected after thorough cleaning by applying a solution of 4 1/2 oz. cleaner to 5 gallons of water, with a mop, cloth or brush. Small utensils should be immersed in this solution.

Prior to disinfection, all poultry, other animals and their feeds must be removed from the premises. This includes emptying all troughs, racks and other feeding and watering appliances. Remove all litter and droppings from floor, walls and other surfaces occupied or traversed by poultry or other animals.

After disinfection, ventilate buildings, coops and other closed spaces. Do not house poultry, or other animals, or employ equipment until treatment has been absorbed, set or dried. All treated equipment that will contact feed or drinking water must be rinsed with potable water before reuse.

E.P.A. Reg. No. 11082-7
E.P.A. Est. 11082-CA-1

41

LONG CHEMICAL INC., 6231 S. MANHATTAN PL., LOS ANGELES, CA 90047

BEST DOCUMENT AVAILABLE