



VALCOR BRAND PRODUCTS

BOTRAN 6 SULPHUR 25 DUST

EPA EST. No. 11169-CA-01

EPA REG. No. 11169-2

ENVIRONMENTAL AND DISPOSAL CAUTIONS

Keep out of any body of water. Fish may be killed. Do not use on germinating seeds or annual seedlings except as suggested. To avoid drift to adjacent areas, do not apply dust when wind velocities are greater than 2-3 mph. Do not feed treated forage to livestock.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal of cleaning of equipment. Open dumping is prohibited.

Pesticide or rinsate that cannot be used or chemically reprocessed should be disposed of in a way approved for pesticides or treated in a water pipe away from water supplies.

Dispose of containers in an approved or approved alternate container disposal site.

Contact factory representative for information on approved alternate disposal methods for this open burning.

DIRECTIONS FOR USE

1. Spray after the following diseases have been identified and confirmed by a reliable laboratory.

For Powdery Mildew: Apply with a backpack sprayer or hand pump sprayer. For the control of powdery mildew on grapes, apply 25 lbs. per acre. For the control of powdery mildew on other plants, apply 10 lbs. per acre. Repeat applications at 7 to 14 day intervals.

BLACKBERRY (7), BOYSENBERRY (7), RASPBERRY (7), Pacific Northwest Only: For the control of powdery mildew, apply 25 lbs. per acre. Repeat applications at 7 to 14 day intervals.

FUNGICIDE

| ACTIVE INGREDIENTS: | BY WT |
|-------------------------------------|---------------|
| 2,4-Dichlorophenyl dimethyl sulfide | 25.00 |
| Other inert ingredients | 75.00 |
| Total | 100.00 |

Botran is a Registered Trademark of San Joaquin Sulphur Co., Lodi, California. For more information, contact San Joaquin Sulphur Co., Lodi, California, 95240.

CAUTION

Keep Out of Reach of Children
 Harmful if swallowed. Avoid inhalation of dust. Avoid Contact with skin and eyes. Flush with water and call a physician.

NON WARRANTY

San Joaquin Sulphur Co. warrants that the product is as labeled and that it is free from adulteration. It does not warrant that the product will be effective under all conditions. The user assumes all responsibility for the proper use of the product.

USAGE CAUTIONS

1. Do not use on plants listed in the directions for use.

2. Do not use on plants listed in the directions for use if the plant is stressed or if the weather is unfavorable.

3. Apply Botran 6 Sulphur 25 Dust to the plant at the rate specified in the directions for use.

4. Do not use Botran 6 Sulphur 25 Dust on plants listed in the directions for use if the plant is stressed or if the weather is unfavorable.

GRAPES: For the control of powdery mildew, apply 25 lbs. per acre. Repeat applications at 7 to 14 day intervals.

HEAD LETTUCE, Greenhouse (14): For the control of powdery mildew, apply 10 lbs. per acre. Repeat applications at 7 to 14 day intervals.

PEACHES (1), NECTARINES (1): Brown Rot, Stem Blight, Monilia Powdery Mildew. Use 25 lbs. per acre. Apply Botran 6 Sulphur 25 Dust to the plant at the rate specified in the directions for use.

SWEET CHERRIES: Blossom Blight. Use 50 pounds per acre. Apply Botran 6 Sulphur 25 Dust to the plant at the rate specified in the directions for use.

TOMATO: For the control of stem rot, Blight, Green Blight. Repeat applications 30 lbs. per acre. Dust stem of plant.

ONION, GARLIC: For the control of white rot, Sclerotium blight, apply 250 lbs. per acre. Do not use on plants listed in the directions for use.

ROSE, HYDRANGEA: For the control of plant Botrytis, apply 10 lbs. per acre. Repeat applications at 7 to 14 day intervals.

CHRYSANTHEMUM, GERANIUM, ROSE: For the control of plant Botrytis, apply 10 lbs. per acre. Repeat applications at 7 to 14 day intervals.

GLADIOLUS: For the control of plant Botrytis, apply 10 lbs. per acre. Repeat applications at 7 to 14 day intervals.

SAN JOAQUIN SULPHUR CO.

LODI, CALIFORNIA
 NET WEIGHT 50 LBS.

