



VALOR BRAND PRODUCTS

BT SULFUR 15-50 DUST

EPA REG. No. 11169-6

EPA EST. No. 11169-CA-01

ACTIVE INGREDIENTS:

Bacillus thuringiensis Berliner, 320 International Units of Potency per mg (at least 0.5 billion viable spores per gm)*..... 0.064%
Sulfur 50.000%
INERT INGREDIENTS: 49.936%
100.000%

*Equivalent to potency of 0.145 billion International Units of Potency per pound of this product.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Reasonable care should be exercised to avoid inhalation or contact with open wounds.

ENVIRONMENTAL AND DISPOSAL CAUTIONS

Keep out of lakes, ponds or streams.
Do not apply when weather conditions favor drift from the areas treated. Do not apply where runoff is likely to occur.
DISPOSE OF WASTES by burying in non-crop lands away from water supplies. Empty bags should be buried with wastes or burned. Keep out of smoke.

USAGE CAUTIONS

Do not use for dried fruit.
Application of materials containing sulfur should not be made just prior to or during periods of excessively high temperatures or within 10 days to 2 weeks of an oil spray.
Do not use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Alfalfa: Alfalfa Caterpillar, Powdery Mildew, Rust. Apply 10 to 20 lbs. per acre.
Broccoli, Cabbage, Cauliflower: Powdery Mildew, Cabbage Looper, Diamond-back Moth Larvae, Imported Cabbage Worm. Apply 30 to 50 lbs. per acre.
Celery, Collards, Cotton, Kale, Lettuce, Mustard Greens, Potatoes, Spinach, Turnip Greens: Cabbage Looper, Early and Late Blight, Powdery Mildew, Rust. Apply 25 to 50 lbs. per acre.
Grapes: Grape Leaf Folder, Omnivorous Leaf Roller, Powdery Mildew, Rust. Apply 20 to 50 lbs. per acre. Start treatment as soon as possible after hatching and before larvae are protected by leaf folds. Do not allow workers to enter treated areas to engage in any activity requiring substantial contact with treated foliage for 1 day following application.
Oranges: Fruit Tree Leafroller. Apply 25 to 50 lbs. per acre. Orange Dog. Apply 20 to 50 lbs. per acre.
Strawberries: Cabbage Looper, Rough Skin Cutworm, Powdery Mildew. Apply 25 to 50 lbs. per acre. Sulfur may injure strawberries under certain climatic conditions.
Tomatoes: Tomato Hornworm. Apply 20 to 50 lbs. per acre.

Do not apply to sulfur sensitive crops. Do not apply during high temperatures. Do not use within 2 weeks of an oil treatment, on citrus within 3 to 6 weeks. When grown for processing, consult the processor before applying sulfur.

NON WARRANTY


Seller makes no warranty, expressed or implied, concerning the use of this product other than as indicated on the label. Buyer assumes all risk of use and/or handling of this material, when such use and/or handling is contrary to label instructions.



SAN JOAQUIN SULPHUR CO.

LODI, CALIFORNIA
NET WEIGHT 50 LBS.

BEST DOCUMENT AVAILABLE



VALO BRAND PRODUCTS

BT SULFUR 15-50 DUST



EPA REG. No. 11169-6

EPA EST. No. 11169-CA-01

ACTIVE INGREDIENTS:

Bacillus thuringiensis Berliner, 320 International Units of Potency per mg (at least 0.5 billion viable spores per gm)*..... 0.064%
Sulfur 50.000%
INERT INGREDIENTS: 49.936%
100.000%

*Equivalent to potency of 0.145 billion International Units of Potency per pound of this product.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Reasonable care should be exercised to avoid inhalation or contact with open wounds.

ENVIRONMENTAL AND DISPOSAL CAUTIONS

Keep out of lakes, ponds or streams.

Do not apply when weather conditions favor drift from the areas treated. Do not apply where runoff is likely to occur.

DISPOSE OF WASTES by burying in non-crop lands away from water supplies. Empty bags should be buried with wastes or burned. Keep out of smoke.

USAGE CAUTIONS

Do not use for dried fruit.

Application of materials containing sulfur should not be made just prior to or during periods of excessively high temperatures or within 10 days to 2 weeks of an oil spray.

Do not use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Alfalfa: Alfalfa Caterpillar, Powdery Mildew, Rust. Apply 10 to 20 lbs. per acre.
Broccoli, Cabbage, Cauliflower: Powdery Mildew, Cabbage Looper, Diamond-back Moth Larvae, Imported Cabbage Worm. Apply 30 to 50 lbs. per acre.
Celery, Collards, Cotton, Kale, Lettuce, Mustard Greens, Potatoes, Spinach, Turnip Greens: Cabbage Looper, Early and Late Blight, Powdery Mildew, Rust. Apply 25 to 50 lbs. per acre.

Grapes: Grape Leaf Folder, Omnivorous Leaf Roller, Powdery Mildew, Rust. Apply 20 to 50 lbs. per acre. Start treatment as soon as possible after hatching and before larvae are protected by leaf folds. Do not allow workers to enter treated areas to engage in any activity requiring substantial contact with treated foliage for 1 day following application.

Oranges: Fruit Tree Leafroller. Apply 25 to 50 lbs. per acre. Orange Dog. Apply 20 to 50 lbs. per acre.

Strawberries: Cabbage Looper, Rough Skin Cutworm, Powdery Mildew. Apply 25 to 50 lbs. per acre. Sulfur may injure strawberries under certain climatic conditions.

Tomatoes: Tomato Hornworm. Apply 20 to 50 lbs. per acre.

Do not apply to sulfur sensitive crops. Do not apply during high temperatures. Do not use within 2 weeks of an oil treatment, on citrus within 3 to 6 weeks. When grown for processing, consult the processor before applying sulfur.

NON WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than as indicated on the label. Buyer assumes all risk of use and/or handling of this material, when such use and/or handling is contrary to label instructions.

SAN JOAQUIN SULPHUR CO.

LODI, CALIFORNIA

NET WEIGHT 50 LBS.

BEST DOCUMENT AVAILABLE

