

SBP-1382 0.10% Insecticide Spray

PLANTS

Active Ingredients

Inert Ingredients

CAUTION: KEEP OUT OF REACH OF CHILDREN

Additional of swappers and Europe transfer in the Application of feed and foods to the total provide or specific along treatment. As food processing surface stold be removed or specific during treatment. As food processing surface stold be removed or thoroughly released before using the hospitals patients should be removed from room prior to treatment. Room should be ventilated for 2 hours after spraying. Do not return patients to room units after ventilation. Remove pets birds and cover fish aquarium: before spraying

DIRECTIONS FOR USE

FOR FLIES, MOSQUITOES AND GNATS: Close doors and windows. Spray freely toward ceiling in all directions until room is filled with mist, then leave the room. Keep doors closed for at least 15 minutes. Ventilate the room when treatment is completed.

WASPS: Application should be made in the late, vening when insects are at rest. Spray liberally into hiding and breeding places, conticting as many insects as possible. Repeat as necessary.

CONTAINER DISPOSAL

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place away from water supplies.

Manufactured by

TARGET CHEMICAL CO.

17710 STUDEBAKER ROAD
CERRITOS, CALIFORNIA 90701

E.P.A. Reg. No. 11214-13-44

NET CONTENTS