DIRECTIONS FOR USE

App, wy A. Speritled

GREENHOUSE PESTS

Figure 1 stations of a hiteties on beganial fuchsia, salvial poinsettia, correspondent members, one of geraniums ageratum, petutia, aster, ralenda, a mai gotal roses, azalea and avy.

Contact Spray — For whiteflies apply SBP 1382 2 E.C. spray at 1 pint per 100 gations of water or 1 teaspoonful per gallon. Spray upper and lower surfaces of plant folioge to point of runoff. Spray in late afternoon and at night only, when the temperature ranges from 50°F to 72°F. Repeat application weekly for four weeks or longer for control of adults, pupae, and nymphs. Complete spray schedule on poinsettia before blooms form.

Where fewer applications are preferred, spray at the rate of 1 qt. per 100 gallons of water or 2 teaspoonfuls per gallon. Repeat as necessary. Apply sprays as mentioned above.

Non-Thermal Applications. Direct mist first toward upper corners of the greenhouse, then apply evenly throughout the area. Keep nozzle at least three feet from plant foliage. Apply SBP-1382 2 E.C. at the rate of 1½ fl. oz. thoroughly mixed in 3 fl. ozs. of methylene chloride per 10,000 cu. ft. This is equivalent of 1 gram of active ingredient per 1,000 cu. ft. Keep treated area closed off for at least two hours. Ventilate before re-entry. Apply at weekly intervals for at least four weeks. Applications should be made during the late afternoon or at night when the temperature range is 50°F to 72°F.

HOUSEHOLD PESTS:


For use in warehouses, food processing plants, business offices and industrial plants.

Contact Spray — Roaches, mix 1 1/3 fluid ounces of 24.3 % SBP-1382-2 E.C. in a gallon of water to make a 0.25% emulsion. Mix well before using. Spray thoroughly into hiding places such as cracks, crevices, and moist area around pipes, under refrigerators and along basebiased. Spray directly on insects when possible. Repeat as necessary.

Space Spray — For control of flies, mosquitoes and gnats. Remove or cover exposed food and food handling surfaces including utensils.

.

ACCEPTED



SBP-1382 24.3% E.C.

(CONTAINS 2 POUNDS SBP-1382 R PER GALLON)

ACTIVE INGREDIENTS:

*1 (5-Benzyl-3-furyl) methyl
2,2-dimethyl-3- (2-methylpropenyl)
cyclopropanecarboxylate
Related compounds
Aromatic petroleum hydrocarbons

INERT INGREDIENTS:

*Cis trans isomers ratio: max. 30 (+) cis
and min. 70 (+) trans
:U.S. Patent No. 3 465,007

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid contact with skin. Avoid breathing spray mist. DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME. Keep container closed and store at temperatures above 40°F.

LPA Reg No 11214-14-AA Net C

Net Contents

Gallons

DIRECTIONS FOR USE . . . (Continued)

Remove pets, birds, and cover fish aquariums before spraying. Shut off ventilating systems, air ducts and air conditioning equipment. Mix 1/2 fluid oz. of 24.3% SBP-1382 2 E.C. in a gallon of water to make 0.10% emulsion. Apply as a fine spray at the rate of 2 fluid ozs. per 1,000 cubic feet or 1 gallon per 64,000 cubic feet. Mix well before using. Direct nozzle to upper corner of the room and along ceiling. Avoid inhaling the spray mist. Close off room for at least one-half hour after spraying. Ventilate room before re-entry. Repeat as necessary. Do not remain in treated areas. Ventilate the area after treatment is completed.

CAUTION: KEEP OUT OF REACH OF CHILDREN

This product is toxic to fish and birds. Keep out of lakes, streams and ponds Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

FOOD AREAS — Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using this product in these areas, apply only when the facility is not in operation.

CONTAINER DISPOSAL -- Do not reuse container. C ush used container and bury in non-cropland soil away from wat-

NOTICE — Do not store below 40°F. If exposed to temperatures below 40°F warm to 80°F and mix well before using

Manufactured by

TARGET CHEMICAL CO.

17710 STUDEBAKER ROAD
CERRITOS, CALIF. 90701