

[11273-8]



50 pounds net

thuricide®-HP

144M* DUST

ACTIVE INGREDIENT:

Bacillus thuringiensis
Berliner, potency of 320
International Units (at least
600 thousand viable spores)
per milligram* 0.064%
INERT INGREDIENTS 99.936%
100.000%


*Equivalent to 144 Million International
Units per pound

**CAUTION: KEEP OUT
OF REACH OF CHILDREN.**

STORE IN COOL PLACE

For use only in south central
and western United States
(Texas, New Mexico, Arizona, Oregon,
Washington and California).

Reg. Trade Marks International Minerals & Chemical Corporation



ACCEPTED

UNDER THE FEDERAL INSECTICIDE
ACT, 1947, AS AMENDED
FOR ECONOMIC ANALYSIS AND
CONTROL

RECOMMENDATIONS

Thorough coverage essential when using THURICIDE®HP-144M Dust

Crop	Pest	Lbs. per Acre	Crop	Pest	Lbs. per Acre
BROCCOLI	Cabbage looper	20-40	KALE	Cabbage looper	20-40
	Imported cabbage-worm	20-35		Imported cabbage-worm	20-35
CABBAGE	Cabbage looper	20-40	LETTUCE	Cabbage looper	20-50
	Imported cabbage-worm	20-35			
CAULIFLOWER	Cabbage looper	20-50	MELONS	Cabbage looper	20-40
	Imported cabbage-worm	20-40	MUSTARD GREENS	Cabbage looper	20-40
CELERY	Cabbage looper	20-50		Imported cabbage-worm	20-35
COLLARDS	Cabbage looper	20-40	POTATOES	Cabbage looper	20-40
	Imported cabbage-worm	20-35			
GRAPES	Grape leaf folder	20-50	SPINACH	Cabbage looper	20-40
GRAPES (California)	Omnivorous leafroller	20-50	TOMATOES	Cabbage looper	20-40
				Tomato hornworm	20-35
			TURNIP GREENS	Cabbage looper	20-40
				Imported cabbage-worm	20-35

Start treating as soon as possible after hatching
and before larvae are protected by leaf folds
Apply by ground equipment using 20-50 pounds
per acre to obtain thorough coverage of leaf
surfaces

DIRECTIONS FOR USE

Apply recommended amounts when air is
calm, using ground equipment, aircraft or
hand duster. Apply at first sign of infesta-
tion and repeat at weekly intervals when
needed to maintain control. Good under-
leaf coverage is essential since the insect
must ingest THURICIDE®HP-144M Dust.
After eating THURICIDE®HP-144M Dust,
some insects stop feeding but may other-

wise appear to be unaffected for several days.

Use the higher recommendations against
very heavy infestations, or where it is diffi-
cult to obtain good coverage due to rank
foliage. THURICIDE®HP-144M Dust is
exempt from tolerance requirements, and
may be applied in recommended amounts
up to the day of harvest.

CONTAINER DISPOSAL:

**DO NOT REUSE EMPTY CONTAINER.
DESTROY OR DISCARD IN SAFE PLACE.**

EPA Reg. No. 4456-548

INTERNATIONAL MINERALS & CHEMICAL CORP
IMC PLAZA
Libertyville, Illinois

LOT