


NET CONTENTS 50 LBS.

LOT NO. _____

evital

A GRANULAR HERBICIDE FOR CRANBERRIES



For the control of certain annual and perennial weeds in cranberries grown in New Jersey, Massachusetts, Oregon, Washington, and Wisconsin.

ACTIVE INGREDIENTS: norflurazon (4-chloro-5-(methylamino)-2-(alpha, alpha, alpha-trifluoro-m-tolyl)-3 (2H)-pyridazinone) 5%

INERT INGREDIENTS 95%

TOTAL 100%

EPA Registration No. 11273-12

WARNING: Keep Out of Reach of Children

In case of skin contact, wash thoroughly with soap and water. Wear clean clothes and wash before reuse. Wash hands thoroughly with soap and water before eating, drinking, or smoking.

Dispose of empty bag by burning in a safe place. Do not breathe smoke from burning bag.

Do not contaminate streams, ponds, or watershed areas with unused materials, or with hopper washings when cleaning equipment.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. The manufacturer makes no other warranties of any kind, express or implied, including warranties of merchantability or fitness for a particular use.

GENERAL DIRECTIONS

APPLICATION DIRECTIONS

Apply EVITAL in the early spring as a single application after removal of winter flood and before weed growth resumes. Apply EVITAL as a dry granular treatment through any suitable ground equipment that will ensure uniform coverage. Flooding after application may reduce weed control. Cranberry vines may be injured when application patterns overlap, in treated low wet areas of the bog, or where water, standing over treated soils is splashed onto the vines.

WEEDS CONTROLLED

EVITAL controls the following weeds when used according to the directions:

AT 120 lbs/A:

Little bluestem (Poverty grass)
Nutsedge (*Cyperus dentatus*)
Smokegrass
Barnyardgrass
Needlegrass
Spikerush

AT 80 lbs/A:

Rice cutgrass (Sicklegrass)

NOTE: In a newly planted bog, the maximum amount applied in any one year should not exceed 80 lbs/A. In an established bog, the maximum amount applied in any one year should not exceed 120 lbs/A.



Sandoz, Inc.
Crop Protection
Homestead, Florida 33030