

11273-3

5-31-78

1 of 3

THURONIDE HP SPRAY CONCENTRATE (WETTABLE POWDER)

(EPA REG. NO. 11273-3)

LABEL ADDITION

CROP	PEST	LBS. PER ACRE
Cotton	European budworm cotton bollworm	4-1

For the suppression of light to moderate infestations apply a first crop of spraying or early instar larvae (1st and 2nd instar larval). Reapply as needed to suppress worm suppression of inadequate reduction of worm populations achieved. Use THURONIDE HP Wettable Powder under a pest management program that includes close monitoring of early instar populations. If they are observed to multiply or increase with application and control measures are not sufficient.

Apply THURONIDE HP Wettable Powder spray per acre

and ground spray per acre with 247

gallon spray per acre with 247

ground equipment

MAY 31 1978

11273-3

11273-3



thuricide®-HP

SPRAY CONCENTRATE (Wettable Powder)

ACTIVE INGREDIENT: *Bacillus thuringiensis* Berliner, potency of 16,000 International Units (at least 30 million viable spores) per milligram

INERT INGREDIENTS

3.2%

96.8%

100.0%

*Equivalent to 7.2 billion International Units per pound

STORE IN A COOL PLACE

KEEP OUT OF REACH OF CHILDREN

CAUTION

**AVOID INHALATION OR CONTACT WITH EYES OR OPEN WOUNDS
APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL**

DIRECTIONS FOR USE

GENERAL: Apply at first sign of infestation when worms are small and repeat at weekly intervals as needed to maintain control. Thuricide HPSC wettable powder attacks the larval gut and must be eaten by the insect to be effective. Thorough under leaf coverage is, therefore, essential for best results. After eating Thuricide HPSC wettable powder, worms stop feeding and die within 2-3 days. Since the active ingredient, *Bacillus thuringiensis*, is exempt from tolerance requirements, Thuricide HPSC wettable powder may be applied up to the day of harvest.

MIXING: Pour recommended amount onto surface of partially filled spray tank. Maintain agitation until tank is filled to desired level. Spray tanks equipped with by-pass agitation only should maintain pump circulation at full capacity until mixing is complete. Use the higher rates against very heavy worm infestations, or where it is difficult to get good coverage due to rank foliage. Do not allow dilute sprays to stand in the tank for more than 12 hours. Combinations with chemical insecticides, fungicides, spreader stickers, and stickers in the tank

are generally not deleterious to Thuricide HPSC wettable powder if the mix is used promptly.

GROUND APPLICATION: Use recommended amount of Thuricide HPSC wettable powder in a minimum of 5 to 50 gallons of water, depending on the type of crop and requirements of state regulations. Add a sticker, adhesive or deposit builder where heavy dew or rain is encountered.

GROUND APPLICATION, TREES: Use recommended amount of Thuricide HPSC wettable powder in 100 gallons of water in high pressure, high gallonage, hydraulic sprayers. Wet foliage thoroughly with spray but not to the point of excessive runoff. For mist blowers use recommended amount in 10 gallons of water. To prevent washoff by rain add a sticker to the finished spray.

AERIAL APPLICATION: Use recommended amount of Thuricide HPSC wettable powder in 5 to 20 gallons of water. A sticker should be added under conditions of heavy dew or rain. Apply early morning or evening when the air is calm.

CONTAINER DISPOSAL: Do not reuse empty container. Destroy or discard in safe place.

NOTICE: Sandoz, Inc. makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose concerning this material except those which are contained on Sandoz, Inc.'s label.

EPA

EPA Reg. No. 11272-1

Sandoz, Inc.
Crop Protection

Kenilworth, New Jersey 07033

NET

EPA Est. No. 36299 - CA - 1

3043

RECOMMENDATIONS: Thorough coverage is essential when using Thuricide-HPSC Wettable Powder

Crop	Pest	Lbs. per Acre
ALFALFA (Arizona, West Coast)	Alfalfa caterpillar	1/8 - 1/4
BROCCOLI, BRUSSELS SPROUTS, CABBAGE	Cabbage looper	1/2 - 1
CAULIFLOWER, COLLARDS, KALE, MUSTARD GREENS, TURNIP GREENS	Imported cabbage worm	1/4 - 3/4
BEANS, CELERY, LETTUCE, SPINACH	Diamondback moth	1/4 - 1/2
CUCUMBERS (Southeast only), MELONS	Cabbage looper	1/2 - 1
COTTON (Arizona, California), POTATOES	Cabbage looper	1/4 - 3/4
SOYBEANS	Cabbage looper	1/2 - 1
	Soybean looper	1/2 - 3/4
	Cabbage looper, Green cloverworm,	
	Velvetbean caterpillar	1/4 - 3/4
TOBACCO	Cabbage looper	1/2 - 3/4
	Tobacco budworm, Tobacco	
	hornworm	1/4 - 3/4
TOMATOES	Cabbage looper, Tomato	
	hornworm	1/2 - 3/4
WATERMELONS (Florida)	Rindworm complex	1/4 - 3/4
GRAPES	Grape leaf folder	1/2 - 1
Start treating as soon as possible after hatching and before larvae are protected by leaf folds. Apply by ground equipment in up to 200 gallons total spray per acre to obtain thorough coverage of leaf surfaces.		
GRAPES (California)	Omnivorous leafroller	1/2 - 1
The following must be applied by ground equipment only:		
ORANGES (Arizona, California)	Fruit tree leafroller	1/2 - 1
	Orange dog	1/4 - 1/2
WALNUTS (California)	Redhumped caterpillar	1/4 - 1
SHADE TREES, ORNAMENTALS, FOREST	Spring & Fall cankerworm, Fall	
	webworm, Great Basin Tent	
	caterpillar (West U.S.) Tent	
	caterpillar	1/4 - 1/2
	Redhumped caterpillar	
	(California)	1/4 - 1
	Oakmoth larvae (California),	
	Gypsy Moth (Northeast)*	1/2 - 1

*** GYPSY MOTH: Aerial Application**

Thuricide HPSC wettable powder may also be applied by air to prevent excessive defoliation by gypsy moth. For best results, spray when leaf expansion reaches 40% - 50%, and respray 14 days later. Use one pound of Thuricide HPSC wettable powder in 1-4 gallons finished spray per acre. Incorporate in the spray mix an evaporation retardant, such as molasses, Cargill's Insecticide Base, polyethylene glycol E-400, and a film-forming sticker. To obtain thorough coverage use spray equipment suitable to deliver droplets with an average diameter of less than 300 microns.

ACCEPTED
MAY 31 1978
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. **11273-3**

mc