thuricide® 24B

AQUEOUS CONCENTRATE FOR AERIAL APPLICATION

ACTIVE INGREDIENT: Bacillus thuringiensis Berliner 24.0%
Equivalent to 6,000 International Units per milligram
INERT INGREDIENTS 76.0%

Equivalent to 1.2 billion International Units per pint

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans: Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Applications must be made at the recommended rate of THURICIDE 24B in the recommended amount of water per acre. Use sufficient water to obtain adequate coverage of the foliage. Maintain agitation in the tank during mixing and application of THURICIDE 24B to provide uniform suspension of the product. Add recommended adjuvants specified on the label to the tank mix as necessary.

For control of forest insects, apply THURICIDE 24B by aerial application.

RECOMMENDATIONS FOR FOREST INSECT CONTROL

For control of forest insects with THURICIDE 24B use the following application rates and directions:

Pest	Rate per Acre	DIRECTIONS FOR USE
Spruce budworm	1 - 3	Apply when leaf expansion reaches 40% to 50% as infestation warrants. If eggs hatch over a long period of time or if reinfestation occurs, spray about 14 days after first application.
Jack pine budworm	1 - 3	Apply when leaf expansion reaches 40% to 50% as infestation warrants. If eggs hatch over a long period of time or if reinfestation occurs, spray about 14 days after first application.
Forest tent caterpillar	1 - 3	Apply when leaf expansion reaches 40% to 50% as infestation warrants. If eggs hatch over a long period of time or if reinfestation occurs, spray about 14 days after first application.
Gypsy moth	1 - 3	Apply when leaf expansion reaches 40% to 50% as infestation warrants. If eggs hatch over a long period of time or if reinfestation occurs, spray about 14 days after first application.
Eastern spruce budworm	1 - 3	Apply when leaf expansion reaches 40% to 50% as infestation warrants. If eggs hatch over a long period of time or if reinfestation occurs, spray about 14 days after first application.
Western spruce budworm	1 - 3	Apply when leaf expansion reaches 40% to 50% as infestation warrants. If eggs hatch over a long period of time or if reinfestation occurs, spray about 14 days after first application.
Western hemlock looper	1 - 3	Apply when leaf expansion reaches 40% to 50% as infestation warrants. If eggs hatch over a long period of time or if reinfestation occurs, spray about 14 days after first application.
Jack pine budworm	1 - 3	Apply when leaf expansion reaches 40% to 50% as infestation warrants. If eggs hatch over a long period of time or if reinfestation occurs, spray about 14 days after first application.

STORAGE AND DISPOSAL

Storage: Store in a cool, dry place.
Disposal: Do not reuse empty container. Triple rinse container. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill.

NOTICE: Sandoz Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use. To the extent consistent with applicable law, Sandoz Inc. makes no other express or implied warranty concerning this product and specifically disclaims any implied warranty of merchantability or fitness for a particular purpose. Sandoz Inc. shall not be liable for consequential damages resulting from the use or handling of this product. Use of this product constitutes acceptance of these conditions.

NET U.S. GALLONS

SANDOZ

Sandoz, Inc
Crop Protection
480 Camino Del Rio So
San Diego, California 92108

E.P.A. Reg. No. 11273-29

22

DIRECTIONS FOR USE

GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

"Remove patients from hospital rooms before spraying. Ventilate rooms for at least two hours before returning patients."

STORAGE AND DISPOSAL

Storage	Do not store near heat or open flame
Disposal	Do not use empty container Wrap container and put in trash collection

To kill flies mosquitoes and gnats Close room and discharge spray mist vigorously in all parts of rooms using about 2 or 3 ounces for each 1 000 cubic feet or room space Leave room closed for at least 15 minutes Ventilate room thoroughly before re entry Sweep up and discard fallen insects Repeat as needed

To kill cockroaches and waterbugs Apply in all cracks and crevices in woodwork underneath sinks behind pipes and in all places suspected of harboring these insects Contact as many insects as possible Repeat as needed

Insecticide #922

For Use Only in Household and Non-food Areas of Industrial Buildings: schools, warehouses, office buildings, theatres, garages, hotels, motels, hospitals, kennels.

ACTIVE INGREDIENTS:	
Tetramethrin [(1-Cyclohexene 1, 2-dicarboximido) methyl 2, 2-dimethyl-3-(2-methylpropenyl) cyclopropanecarboxylate]	0.075%
*3-Phenoxybenzyl d-cis and trans** 2, 2-dimethyl 2-(2-methylpropenyl) cyclopropanecarboxylate	0.024%
*Other isomers	0.001%
Petroleum distillate	99.776%
INERT INGREDIENTS:	0.124%
	100 000%

*d (cis, trans) phenothrin

*cis trans isomer ratio Max 25% (+ or -) cis
Min 75% (+ or -) trans

CAUTION

KEEP OUT OF REACH OF CHILDREN

See side panel for additional precautionnary statements

Manufactured By

EUREKA LABORATORIES
2033 W FULTON ST
CHICAGO, IL 60612
EPA Reg No 37346-6
EPA Est 37346-IL-1