



thuricide® 24B

AQUEOUS CONCENTRATE FOR AERIAL APPLICATION

ACTIVE INGREDIENT *Bacillus thuringiensis* Berliner, potency of 6,000 International Units
(at least 9 million viable spores) per milligram* 1.03%
INERT INGREDIENTS 98.97%
100.00%

*Equivalent to 6.0 billion International Units per quart

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS: Avoid inhalation or contact with eyes or open wounds

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

Suspension must be shaken or stirred before use. Pour the recommended amount of THURICIDE 24B into the required volume of water in the spray tank. Maintain agitation. Do not allow diluted sprays to remain in the tank for more than 24 hours. THURICIDE 24B is formulated to provide desirable coverage and stickability on leaf surfaces. Additional adjuvants, spreaders, or stickers may be added but are not essential.

For control of the forest insects listed opposite, apply THURICIDE 24B by air, diluting with water to achieve a spray volume of at least 1/2 gallon per acre. Best results are obtained when THURICIDE 24B is applied to dry foliage with a calibrated aircraft capable of obtaining droplet sizes below 300 microns and preferably in the range of 50-150 microns.

Thorough coverage is essential when using THURICIDE 24B. Use the higher rates against heavy worm infestations.

RECOMMENDATIONS FOR SHADE TREES, ORNAMENTALS & FOREST

Pest	Pints per Acre	Directions for Use
Spring and Fall cankerworm	1 1/2-3	Apply when leaf expansion reaches 40%-50% as infestation warrants. If eggs hatch over a long period of time or if reinfestation occurs, spray about 14 days after first application.
Elm spanworm	1 1/2-3	
Tent caterpillars	1 1/2-2 1/2	
Gypsy moth	2 1/2-4	
Spruce budworms	2-4	Apply when most larvae are 3rd-4th instar. Also consider the opening of the bud cap to ensure foliage exposure.
Douglas fir tussock moth	1 1/2-3	Apply after eggs have hatched and early instar larvae are feeding on exposed foliage.
Jack pine budworm	1 1/2-3	
Bagworm	1 1/2-3	
California oak moth	1 1/2-3	
Western tussock moth	1 1/2-3	
Fruit tree leafroller	1 1/2-3	
Mimosa webworm	1 1/2-3	
Redhumped caterpillar	1/2	1-1 1/2
Fall webworm	1-1 1/2	

STORAGE AND DISPOSAL

STORAGE Store in a cool place.

DISPOSAL Do not reuse empty container. Destroy or discard in a safe place.

NOTICE Sandoz Inc. makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material except those which are contained on this label.

LOT
NET

U.S. GALLONS



Sandoz, Inc.
Crop Protection
480 Camino del Rio So.
San Diego, California 92108

EPA Reg. No. 11273-29
EPA Est. No. 36299-CA-1

APR 20 1982

11273-29