

teknar™

AQUEOUS CONCENTRATE
A LARVICIDE FOR MOSQUITO CONTROL

ACTIVE INGREDIENTS:
Bacillus thuringiensis Berliner var. israelensis, potency of 1500 AA Units (Aedes aegypti units) per milligram* 0.8%
INERT INGREDIENTS 99.2%
100.0%

*Equivalent to 1.4 billion AA Units per quart.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

Hazardous to Humans: Avoid inhalation or contact with eyes or open wounds.

NET _____ U.S. GALLONS
LOT _____



Sandoz, Inc.
Crop Protection
488 Camino Del Rio So.
San Diego, California 92108

EPA Est. No. 26299-CA-1
EPA Reg. No. 11273-20

39

TEKNAR™

LIQUID CONCENTRATE

KILDER FOR MOSQUITO CONTROL

ANALYSIS:


As Sandoz var. israelensis, potency of 1500 AA Units (a) per milligram	0.8%
ITS	99.2%
	100.0%

1 billion AA Units per quart.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS
 Avoid inhalation or contact with eyes or open wounds.

 Sandoz, Inc.
 Crop Protection
 480 Camino Del Rio So.
 San Diego, California 92108

EPA EN No. 30290-CA-1
 EPA Reg. No. 11273-30

39

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Storage: Store in a cool place. Activity may be impaired by prolonged storage at temperatures above 90°F. Protect from freezing.

Disposal: Do not reuse empty container. Destroy or discard in a safe place.

GENERAL: Apply TEKNAR Aqueous Concentrate when larvae are in 1st to early 4th instar. Larvicidal action is expected within 24 hours. Reapply as needed.

MIXING: Always shake or stir thoroughly before use. Pour recommended amount of TEKNAR onto surface of water in a nearly filled spray tank. Maintain agitation. Do not allow dilute sprays to stand in the tank for more than 12 hours.

APPLICATION: Apply with standard spray equipment using recommended amount of TEKNAR. Use sufficient water to assure complete coverage at recommended rate. For aerial application, a volume of 3 to 5 gallons/surface acre is recommended.

RECOMMENDATIONS: Use lowest rate when 1st to 3rd instar larvae are predominant, and highest rate when late 3rd to early 4th instar larvae are predominant in the mosquito population. Rate ranges, according to habitat are:

Habitat	Rate Required For Control
floodwater, standing ponds, and ditches.	0.5-1.0 pts./A
tidal water and salt marshes, catch basins, and storm water retention areas.	1.0-2.0 pts./A
water polluted with sewage (moderate organic content).	2.0 pts./A

NOTICE: Sandoz, Inc. makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material except those which are contained on this label.

ACCEPTED

APR 2 1981

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 11273-30

11273-30

40