

sollicam[®]

FLOWABLE HERBICIDE

11273 32

FOR WEED CONTROL IN ESTABLISHED TREE FRUIT AND NUT CROPS
(Apricots, Cherries, Filberts, Nectarines, Peaches, Plums, Prunes, Walnuts)

ACTIVE INGREDIENT:
norflurazon [4-chloro-5-(methylamino)-2-(alpha,
alpha, alpha-trifluoro-m-tolyl)-3(2H)-pyridazinone] . . . 50.0%
INERT INGREDIENTS 50.0%
100.0%

(1 gallon contains 5 lbs. of norflurazon)

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
AND DOMESTIC ANIMALS

WARNING: Causes eye irritation. Do not get in eyes, on skin, or on clothing. In case of contact with eyes, immediately flush eyes with water for 15 minutes. Get medical attention.

ENVIRONMENTAL HAZARDS

Keep out of lakes, streams and ponds. Do not apply when weather conditions favor runoff or drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. The manufacturer makes no other warranties of any kind, express or implied, including warranties of merchantability or fitness for a particular use.

LOCATION OF USE

For use in California and West of the Cascade Mountains in Oregon and Washington

WEEDS CONTROLLED

Grass Weeds
Annual bluegrass, Barnyardgrass, Crabgrass, Italian ryegrass, Rippgut brome, Wild barley, Witchgrass.
Broadleaf Weeds
Chickweed, Common purslane, False dandelion, Fiddleneck, Filaree (red stem and white stem), Horseweed (maretail), Little mallow*** (cheeseweed), London rocket, Pineapple weed, Puncturevine, Redmaids, Russian thistle, Shepherdspurse.

WEEDS SUPPRESSED

Bermudagrass, Common lambsquarters, Groundsel, Pigweeds (redroot, tumble, and green amaranth), Purple nutsedge, Quackgrass, Sowthistle.

* For control only on medium to fine soils
** For control only on coarse soils
*** For control only on coarse and medium soils

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

GENERAL INFORMATION

Crops

Use SOLICAM Flowable Herbicide on the crops below that have been established at least one year in the field. Do not use on nursery stock.

Apricots	Peaches
Cherries	Plums
Filberts	Prunes
Nectarines	Walnuts

Directions

Apply SOLICAM Flowable to the soil surface before the weeds emerge. The orchard floor must be free of weeds and must be relatively free of plant residues.

Apply as a directed spray. Avoid contact with fruit and foliage. Make only one application per year. Do not apply when nuts or fruit are on the ground at harvest.

Apply SOLICAM Flowable anytime from fall to early spring in California. In other states, apply before weed growth resumes in the spring.

Sprinkler irrigation is recommended if no rainfall occurs within two weeks after application.

Do not graze livestock on treated areas. Do not feed or graze cover crops grown in treated orchards.

STORAGE AND DISPOSAL

Prohibition: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

Pesticide Disposal: Pesticide, spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

Container Disposal: Triple rinse (or equivalent) and dispose in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

General: Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burning.

APPLICATION DIRECTIONS

Broadcast Rates per Acre

Use 1 1/2 - 3 quarts of SOLICAM Flowable with 20-100 gallons of water per acre.

Use the lower rates on coarse soils.

Use the higher rates on medium to fine soils or soils having organic matter content of greater than 3%.

Strip or Band Treatment

When applying a strip treatment of SOLICAM Flowable, the following formula may be used to calculate the amount per acre:

$$\begin{array}{rcccl} \text{Width of strip} & & \text{Amount per} & & \text{Amount per} \\ \text{in feet} & & \text{acre for} & = & \text{acre for} \\ \text{Distance between} & \times & \text{broadcast} & & \text{strip} \\ \text{rows in feet} & & \text{treatment} & & \text{treatment} \end{array}$$

MIXING AND EQUIPMENT INSTRUCTIONS

Apply accurately with a fixed boom sprayer. Equip sprayer with 8003 to 8006 Tee-jet type nozzle tips or equivalent for application. Use screen size 50 mesh or larger. If a by-pass line is used, discharge at the bottom of the tank to minimize foaming.

Clean and calibrate the sprayer. Fill the spray tank with one-half the required amount of clean water and add the recommended amount of SOLICAM Flowable while maintaining agitation. Triple rinse empty container and add all rinsate to spray tank. Maintain agitation while filling and spraying.



Sandoz, Inc. Crop Protection
480 Camino del Rio South, San Diego, California 92108

Net Wt. 1 gal.

EPA Reg. No. 11273-32

36

BEST DOCUMENT AVAILABLE