


Solicom 5F

FLOWABLE HERBICIDE

FOR WEED CONTROL IN ESTABLISHED TREE FRUIT AND NUT CROPS
(Apricots, Cherries, Filberts, Nectarines, Peaches, Plums, Prunes, Walnuts)



ACTIVE INGREDIENT:

norflurazon [4-chloro-5-(methylamino)-2-(alpha,

alpha, alpha-trifluoro-m-tolyl)-3(2H)-pyridazinone] .. 50.0%

INERT INGREDIENTS 50.0%

100.0%

(1 gallon contains 5 lbs. of norflurazon)

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS
AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation.
Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

in eyes: Flush with plenty of water.
Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

Do not apply directly to any body of water.
Do not apply when weather conditions favor runoff or drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

NOTICE: Sandoz, Inc. makes no warranty, expressed or implied, including the warranties of merchantability and/or fitness for any particular purpose concerning this material, except those which are contained on this label.

NET WT.

LOCATION OF USE

For use in California and West of the Cascade Mountains in Oregon and Washington.

WEEDS CONTROLLED

Grass Weeds

Annual bluegrass, Barnyardgrass, Crabgrass, Italian ryegrass; Ripgut brome, Wild barley, Witchgrass.

Broadleaf Weeds

Chickweed, Common purslane, False dandelion; Fiddleneck, Filaree (red stem and white stem), Horseweed (maretail), Little mallow*** (cheeseweed), London rocket, Pineapple weed, Puncturevine, Redmaids, Russian thistle, Shepherdspurse.

WEEDS SUPPRESSED

Bermudagrass, Common lambsquarters, Groundsel, Pig-weeds (redroot, tumble, and green amaranth), Purple nightshade, Quackgrass, Sowthistle.

* For control only on medium to fine soils.

** For control only on coarse soils.

*** For control only on coarse and medium soils.

EPA Reg. No. 11273-32
EPA Est. No.

Sandoz Inc., Crop Protection, 480 Del Rio So., San Diego, CA 92108

LE

DIRECTIONS FOR USE

✓ It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL INFORMATION

Crops

Use SOLICAM® Flowable Herbicide on the crops below that have been established at least one year in the field. Do not use on nursery stock.

Apricots	Peaches
Cherries	Plums
Filberts	Prunes
Nectarines	Walnuts

Directions

Apply SOLICAM Flowable to the soil surface before the weeds emerge. The orchard floor must be free of weeds and must be relatively free of plant residues.

Apply as a directed spray. Avoid contact with fruit and foliage. Make only one application per year. Do not apply when nuts or fruit are on the ground at harvest.

Apply SOLICAM Flowable anytime from fall to early spring in California. In other states, apply before weed growth resumes in the spring.

Sprinkler irrigation is recommended if no rainfall occurs within two weeks after application.

Do not graze livestock on treated areas. Do not feed or graze cover crops grown in treated orchards.

STORAGE AND DISPOSAL

Storage: Do not store near or contaminate feed or foodstuffs.

Pesticide Disposal: Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State or local procedures.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by incineration if allowed by state and local authorities.

APPLICATION DIRECTIONS

Broadcast Rates per Acre

Use 1½-3 quarts of SOLICAM Flowable with 20-100 gallons of water per acre.

Use the lower rates on coarse soils.

Use the higher rates on medium to fine soils or soils having organic matter content of greater than 3%.

Strip or Band Treatment

When applying a strip treatment of SOLICAM Flowable, the following formula may be used to calculate the amount per acre:

Width of strip in feet	Amount per acre for broadcast treatment	Amount per acre for strip treatment
Distance between rows in feet		

MIXING AND EQUIPMENT INSTRUCTIONS

Apply accurately with a fixed boom sprayer. Equip sprayer with 8003 to 8006 Tee-jet type nozzle tips or equivalent for application. Use screen size 50 mesh or larger. If a by-pass line is used, discharge at the bottom of the tank to minimize foaming.

Clear and calibrate the sprayer. Fill the spray tank with one-half the required amount of clean water and add the recommended amount of SOLICAM Flowable while maintaining agitation. Triple rinse empty container and add all rinsate to spray tank. Maintain agitation while filling and spraying.