

BEST DOCUMENT AVAILABLE

Since 1855
GROWELL

TOMATO AND VEGETABLE SPRAY

Kills Tomato Hornworm,
Whitefly, Beetles (Flea,
Cucumber & Colorado
Potato) And Other
Insects on Tomatoes
& Vegetables *

May Be Used Up To
Day of Harvest

ACTIVE INGREDIENTS

Pyrethrins	0.030%
* Pyrethrin Butoxide Technical	0.160%
Rotenone	0.128%
Other Cube Resins	0.238%
Petroleum Distillates	0.120%
INERT INGREDIENTS	99.324%
TOTAL	100.00%

* Equivalent to 0.128% rotenone. Expressions of other
and 0.128% rotenone are based on weight.

KEEP OUT OF REACH OF CHILDREN

WARNING

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY INFORMATION

NET WEIGHT
12 OZ

SOLD BY:
J & L ADKES, INC.
JAMAICA, N.Y. 11423

1 A-1 listed

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

KEEP OUT OF REACH OF CHILDREN. HARMFUL IF SWALLOWED OR INHALED. Avoid breathing
spray or contact with eyes or skin. Wash thoroughly after use and before eating or drinking. Do not eat
tomatoes or vegetables for 12 hours.

Read entire label before using. Use only on plants listed on label. Use only on plants listed on label. Use only on plants listed on label. Use only on plants listed on label.

ENVIRONMENTAL HAZARDS

This product may be harmful to some aquatic organisms. Do not use near water. Do not use near water. Do not use near water. Do not use near water.

PHYSICAL HAZARDS

WARNING: Contains under pressure. Do not use or store near heat or open flame. Exposure to temper-
atures above 130°F may cause bursting. Do not puncture. Never throw container into fire or in
trash.

DIRECTIONS FOR USE

Read a label of Federal law to use this product in a manner consistent with its labeling.

STORAGE: Store in original container in a cool storage area away from other pesticides. Do not use food
and feed bags, cans, or other containers. Do not use or store near heat or open flame. Exposure to tem-
peratures above 130°F may cause bursting.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in
trash.

CONTAINER DISPOSAL: Recycle, if available, and discard container in trash. Do not reuse or puncture.

SHAKE WELL BEFORE USING

GROWELL TOMATO & VEGETABLE SPRAY may be used up to and including the day of picking to
control certain insects on TOMATOES and certain VEGETABLES. Protect plants throughout the grow-
ing season by spraying early in the morning or in the evening when the air is still. Before insects become
numerous. Spray at 7-10 day intervals and after heavy rain. When spraying, cover foliage thoroughly
paying particular attention to the underside of leaves, avoiding excessive over wetting as injury may
occur. Spray with one 1/2 second burst and where possible, spray so as to contact as many insects as
possible.

TO OPERATE: SHAKE CAN WELL BEFORE USING. Remove protective cap. Press button to release
spray. Keep pump away from you. IMPORTANT: HOLD CAN IN AN UPRIGHT POSITION AND
AT LEAST 12 INCHES AWAY FROM PLANT WHILE SPRAYING.

WARRANTY STATEMENT: Federal law requires labeling of use, storage and handling of this material to
be printed on the label of the container. The purchaser assumes all responsibility for the use of this material. No warranty is made by the manufacturer as to the results of its use or
any other aspect of product use.

FOR USE ON

Tomatoes
Eggplants, Peppers, Potatoes
Cucumbers, Squash, Melons
Beans
Green Corn

TO CONTROL

Colorado Potato Beetle, Flea Beetle, Green Pear Aphid,
Hornworm, Sawfly
Black Beetle, Colorado Potato Beetle, Flea Beetle, Green
Pear Aphid, Leafhoppers
Striped & Spotted Cucumber Beetles
Beetles, Mexican, Bean & 1/2 Spotted Cucumber, Whitefly
Japanese Beetle

EPA Reg. No. 5535-81

EPA Est. 5483-MA-1

(B)

Tomato & Veg. Spray No. 51970 - 8/4/83

THURICIDE

32LV
AQUEOUS CONCENTRATE FOR
AERIAL OR GROUND APPLICATION

ACTIVE INGREDIENT:

Bacillus thuringiensis Berliner, potency of 8,000 International Units
(at least 12 million viable spores) per milligram* 1.6%

INERT INGREDIENTS: 98.4%

100.0%

*Equivalent to 8.0 billion International Units per quart.

KEEP OUT OF THE REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans: Avoid inhalation or contact with eyes or open wounds.

LOT NO. _____

NET _____ GALLONS



Sandoz, Inc.
Crop Protection
480 Camino del Rio So.
San Diego, California 92108

EPA EST. NO. 36289-CA-1

EPA REG. NO. 11273-35

BEST DOCUMENT AVAILABLE

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labelling.

Suspension must be shaken or stirred before use. Pour the recommended amount of THURICIDE 32LV into the required volume of water in the spray tank. Agitate as necessary to maintain suspension. Do not allow diluted sprays to remain in the tank for more than 24 hours. THURICIDE 32LV is formulated to provide desirable coverage and stickability on leaf surfaces. Additional adjuvants, spreaders, or stickers may be added but are not essential.

Ground Application:

Use adequate water to obtain good foliar coverage. Wet foliage but do not allow excessive run-off. Apply the recommended per acre rates of THURICIDE 32LV with the following suggested amounts of water:

- 100 gallons with high gallonage hydraulic sprayers
- 10 gallons with mist blower

Aerial Application:

Apply THURICIDE 32LV at recommended rates as a spray mix diluted with water. Use not less than one half gallon of spray mix per acre. Best results are obtained when THURICIDE 32LV is applied to dry foliage with a calibrated aircraft capable of obtaining droplet sized below 300 microns and preferably in the range of 50-150 microns.

Thorough coverage is essential when using THURICIDE 32LV. Use the higher rates against very heavy worm infestations.

BEST DOCUMENT AVAILABLE

← Thorough coverage is essential when using THURICIDE 32LV. Use the lower rate for light to moderate infestations. Use the higher rates for heavier infestations.