

PLANTS FOR USE

AZALEAS Apply at 1/2 lb. per 100 square feet. Water thoroughly after application. Reapply every 14 days until disease control is obtained.

CARNATIONS Apply at 1/2 lb. per 100 square feet. Water thoroughly after application. Reapply every 14 days until disease control is obtained. Do not apply at water temperatures below 50° F.

CHRYSANTHEMUMS Apply at 1/2 lb. per 100 square feet. Water thoroughly after application. Reapply every 14 days until disease control is obtained. Do not apply at water temperatures below 50° F.

ROSES Apply at 1/2 lb. per 100 square feet. Water thoroughly after application. Reapply every 14 days until disease control is obtained. Do not apply at water temperatures below 50° F.

LAWNS TURF (nonpastured areas only) Apply at 1/2 lb. Brown patch, copper spot, dollar spot, dollar blight, and winter blight. St. Augustine grass - use 4 levelspoons per 100 square feet. Water thoroughly after application. Begin when growth starts in early spring. Continue at 14 day intervals or more frequently during periods of favorable weather.

CAUTION To prevent damage to ornamental plants, do not spray on blossoms, flowers, buds, or seed pods.

Do not apply to lawns or turf where there is a chance of water runoff into streams, lakes, or ponds. Do not apply to lawns or turf where treated lawns or lawns treated with other fungicides may be sprayed. Do not apply to lawns or turf where treated lawns or lawns treated with other fungicides may be sprayed.

Do not apply to lawns or turf where there is a chance of water runoff into streams, lakes, or ponds. Do not apply to lawns or turf where treated lawns or lawns treated with other fungicides may be sprayed.

NOTICE Do not apply to lawns or turf where there is a chance of water runoff into streams, lakes, or ponds. Do not apply to lawns or turf where treated lawns or lawns treated with other fungicides may be sprayed.



**AGRI-CHEMICAL
PRODUCTS DIVISION**

Unilever
PRODUCTS

CAPTAN FLOWABLE

A FUNGICIDE FOR PLANT DISEASE CONTROL

Use as a spray for control of certain fungus diseases of ornamental plants. Use as a soil treatment for control of certain soil borne diseases.

ACTIVE INGREDIENTS:

**CAPTAN: N-((Trichloromethyl)thio)-4-cyclohexene-1,
2-dicarboximide**

25%

INERT INGREDIENTS

75%

100%

Not licensed for use in countries where valid patents still exist.

EPA Reg. No. 11287-5

CONTENTS: ONE QUART

Captan may be combined effectively with other fungicides.

GRASS AND TURF — Use 7 pounds per gallon of water. Apply 10 gallons per 1,000 sq. ft. of turf. Begin when grass starts in active growth at 7 to 14 day intervals throughout the season. Do not graze treated areas. Do not allow clippings to reseed.

LEAVES AND FOLIAGE — Leaf spot, Disease of lawn, Damping-off, Gray mold, Cotton boll rot, Tomato blight, and brown spot of impatiens — Use 7 pounds per gallon of water. Apply 10 gallons per 1,000 sq. ft. of turf. Begin when grass starts in active growth at 7 to 14 day intervals throughout the season. Do not graze treated areas. Do not allow clippings to reseed.

CAPTAN Avoid inhalation of dust or spray mist. Avoid contact with the skin. Do not contaminate food and feed. This product is toxic to fish. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty container. Destroy it and bury or discard in a safe place. Store in cool, dry place — protect from excessive heat.

NOTICE: AGRI-CHEMICAL PRODUCTS DIVISION-LTV AEROSPACE CORPORATION, MAKES NO WARRANTY, EXPRESS OR IMPLIED, INCLUDING THE WARRANTIES OF MERCHANTABILITY AND/OR FITNESS FOR ANY PARTICULAR PURPOSE CONCERNING THIS PRODUCT except those which are contained on the Agri-Chemical Products Division label attached to the product container.



AGRI-CHEMICAL PRODUCTS DIVISION
LTV AEROSPACE CORPORATION
HOUSTON, TEXAS 77001

Foamspray™
PRODUCTS

CAPTAN FLOWABLE

A FUNGICIDE FOR PLANT DISEASE CONTROL
Use as a spray for control of certain fungal diseases of plants and trees. Use as a soil treatment for control of certain soil-borne diseases.

CAUTION: KEEP OUT OF REACH OF CHILDREN
See other cautions on back label.

ACCEPTED

8/23/73

100%
ALL