DIRECTIONS FOR USE

APCLES: Apple Maggot, Codling Moth, Green Apple Aphid, Fruit Tree Leafroller, Plum Curculio, Red Banded Leafre, Rosy Apple Aphid — Apply 4 to 6 teaspoons per gallon water as a full cover spray. Use 6 teaspoons per ga , water for Plum Curculio. Repeat as necessary in accordance with insect infestations and local or state s, ay programs. Usually suppresses European Red and Two Spotted Mites when used in a seasonal program. Do not apply within 7 days of harvest.

APRICOTS, NECTARINES, PLUMS, AND PRUNES: Apple Maggot. Oriental Fruit Moth, Peach Twig Borer, Plum Curculio, Red Banded Leafroller — Apply 4 to 6 teaspoons per gallon water. Use 6 teaspoons per gallon water for Plum Curculio. On bearing trees apply as a full cover spray up to bloom. On non-bearing trees apply as required throughout the growing season. Do not graze livestock in treated orchards.

PEACHES: Oriental Fruit Moth, Peach Twig Borer, Plum Curculio — Apply 4 to 6 teaspoons per gallon water as a full cover spray. Use 6 teaspoons per gallon water for Plum Curculio. Repeat as necessary in accordance with insect infestations and local or state spray programs. Usually suppresses European Red and Two Spotted Mites when used in a seasonal program. Do not apply within 14 days of harvest. Do not apply in summer application to peaches which are to be dried.

PEARS: Codling Moth, Fruit Tree Leafroller, Grean Apple Aphid, Pear Psylla, Plum Curculio. Red Banded Leafroller, Rosy Apple Aphid — Apply 4 to 6 teaspoons per gallon water as a full cover spray. Use 6 teaspoons per gallon water for Plum Curculio. Repeat as necessary in accordance with insect infestations and local or state spray programs. Usually suppresses European Red and Two Spotted Mites when used in a seasonal program. Do not apply within 7 days of harvest.

For the orcharc pests. Codling Moth, Apple Maggot, Plum Curculio, apply at petal fall and every 10 to 14 days thereafter until control is achieved.

WARNING: KEEP OUT OF REACH OF CHILDREN. MAY BE HARMFUL IF SWALLOWED, INHALED OR ASSORBED THROUGH THE SKIN. Do not get in eyes or on skin, or on clothing. Do not breathe spray mist. In case of contact, wash immediately with soap and water before re-use.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply where weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

NOTE TO PHYSICIAN: Imidan is an organophosphorus insecticide. Atropine is antidotal.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.

•


NOTICE: AGRI-CHEMICAL PRODUCTS DIVISION, LTV AEROSPACE CORPORATION, MAKES NO WARRANTY, EXPRESS OR IMPLIED, INCLUDING THE WARRANTIES OF MERCHANTABILITY AND/OR FITNESS FOR ANY PARTICULAR PURPOSE CONCERNING THIS PRODUCT except those which are contained on the Agri-Chemical Products Division, LTV Aerospace Corporation label attached to the product container.



DALLAS TEXAS 75235







For Fruit Trees, Shade Trees, and Ornamentals CONTROLS:

Apple Maggot Codling Moth European Red Mites

Fruit Moth Green Aphid Leafrollers

Peach Twig Borer Plum Curculio Two Spotted Mites

ACTIVE INGREDIENT: N-(mercaptomethyl) phthalimide S-(0,0-dimethyl-phosphorodithioate) INERT INGPEDIENTS

12.5% 87.5% 100%

Imidan R Registered Trademark of Stauffer Chemical Company U. S. Patent No. 2,767,194

WARNING: KEEP OUT OF REACH OF CHILDREN See reverse side of label for additional warnings.

EPA Reg. No. 11287-7

CONTENTS ONE QUART