

#### DIRECTIONS FOR USE

Use 3 tablespoons of concentrated Sevin Outdoor Spray for 1 gallon of water and apply thoroughly. Do not store diluted spray. Make new dilution for each use.

**ROSES, ORNAMENTALS, FLOWERS, SHRUBS, SHADE TREES:** Japanese beetles, rose aphids, bagworm, elm leaf beetle, ~~bush hopper~~, and oak leafminers, rose slug, periodical cicada (17-year locust), scale insects, Eastern tent caterpillars, blister beetles, leaf rollers, exposed thrips — Spray when first signs of damage appear and repeat every 7-10 days as necessary. Caution — do not use on Boston ivy.

**APPLES:** Apple maggot, rosy apple aphid, fruit tree leaf rollers, Eastern tent caterpillar, European apple sawfly, apple rust mite.

**PEARS:** Pear psylla, Japanese beetle, green fruitworm, pear leaf blister mite, pear rust mite — Make application between bud break and bloom, and a second at petal fall and then at 3 to 4 weeks intervals for 3 to 4 months as necessary. May be applied within 1 day of harvest.

**PEACHES, APRICOTS AND NECTARINES:** Oriental fruit moth, peach twig borer, plum curculio, codling moth, fruit tree leaf roller, June beetle, Japanese beetle, European earwig — Do not apply within one day of harvest for peaches, and within 3 days of harvest for apricots and nectarines. For peach tree borer, apply 2 or 3 sprays to trunk from ground to scaffold lines timed with moth flight. For codling moth, oriental fruit moth and plum curculio, apply at petal fall and every 10-14 days thereafter until control is achieved.

**LAWN INSECTS:** Sod webworms, fall armyworms, earwigs, grasshoppers, fleas, and ants — Use 2 tablespoons per gallon of water for each 50 sq. ft. of established lawn area covered. For best results use a hose-on type sprayer and apply after mowing and watering area to be treated. DO NOT water for two days after treatment. Repeat in 2 to 3 weeks as necessary.

**TOMATOES, PEPPERS, CUCUMBERS, SUMMER SQUASH:** Pickleworm, fruitworms, melon worms, hornworms, fall armyworms, corn borer — Apply when insects first appear. Repeat at 7 to 10 day intervals. May be applied within 1 day of harvest.

**CAUTION, HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN.** Avoid breathing spray. Do not take internally. Skin contact may be harmful. Wash hands and face before eating. Keep children and animals off treated lawns until these areas are dry.

**Note for Physician:** Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. **AVOID CONTAMINATION OF FOOD.** To avoid possible fruit thinning on apples, do not apply within 15 days after petal fall. Dispose of container properly by burying.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTICE: AGRI-CHEMICAL PRODUCTS DIVISION — LTV AEROSPACE CORPORATION MAKES NO WARRANTY, EXPRESS OR IMPLIED, INCLUDING THE WARRANTIES OF MERCHANTABILITY AND/OR FITNESS FOR ANY PARTICULAR PURPOSE CONCERNING THIS PRODUCT except those which are contained on the Agri-Chemical Products Division, LTV Aerospace Corporation, label attached to the product container.



**AGRI-CHEMICAL  
PRODUCTS DIVISION**

DALLAS, TEXAS 75235

# Foamspray

## PRODUCTS

# GENERAL OUTDOOR SPRAY

With SEVIN\*

For Use on Ornamentals, Flowers, Lawns,  
Fruit Trees, and Vegetables

#### ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate) 23%

#### INERT INGREDIENTS

77%

100%

\*Trademark of Union Carbide Corporation for the active ingredient, 1-naphthyl N-methylcarbamate.

**CAUTION: KEEP OUT OF REACH OF  
CHILDREN  
HARMFUL IF INHALED OR  
SWALLOWED**

See back label for additional  
cautions

EPA REG NO. 11287-9

CONTENTS ONE QUART

