

PM 32

11321-7

8/14/97

Page 1 of 26

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

AUG 14 1997

Ms. Cristina N. Griffin, Agent  
T-Chem Products, Inc.  
C/O Delta Analytical Corp.  
7910 Woodmont Avenue  
Suite #1000  
Bethesda, MD 20814

Dear Ms. Griffin:

Subject: Pure Bright Bleach  
EPA Registration No. 11321-7  
Your Amendment Dated May 20, 1997

This is in response to your label amendment to add all uses from EPA Reg. No. 11321-6 to the subject product.

The labeling referred to above submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended is acceptable subject to the following comments.

1. On the front center panel of page #1, the label must bear directions for each recommended use and level of antimicrobial activity. Since pages 23 and 24 bear label claims as a virucide and as a tuberculocide, the front label must be expanded to reflect the proposed claims. On the front panel, the wordings should be expanded to reflect these terms "Virucide and Tuberculocide."
2. Page 2, expand the use directions for food contact surfaces to read "The necessity for removal of gross food particles and soil by a pre-flush, or pre-scape and, when necessary, pre-soak treatment. Thoroughly wash the surface or object with a good detergent or compatible cleaner, followed by a potable water rinse before application of the sanitizing solution for at least 2 minutes. Let air dry." For guidance refer to item #3 of DID/TSS-17 enclosure.

At the middle of the right panel, revise "and other surfaces: to read "and other hard inaminate surfaces." Also, under disinfecting bathroom, revise scrub with a brush to read "scrub with a brush making sure to get under.


CONCURRENCES							
SYMBOL							
SURNAME							
DATE							

- 3. On page #23, revise the HIV-1 label use directions to read as follows "...Kills HIV-1 on Pre-Cleaned Environmental Hard Surfaces/Objects Previously Soiled with Blood/Body Fluids ....". Refer to the attached enclosure for guidance. Also, revise the cleaning procedures to indicate "The need to clean the specific surfaces/objects, to remove blood and other body fluids before applying disinfectant."
- 4. On page #24, the second paragraph from the top, revise on surfaces to read "on hard nonporous surfaces". Also, in the middle of the panel, delete "Remove visible body fluids" and add/replace with "Remove gross filth or heavy soil from surfaces or objects prior to application of product." For guidance, refer to item #3 of DIS/TSS-15.

Resubmit one copy of the revised final printed label incorporating the above comments.

If you have any questions, please call Marianne Clark at (703) 308-6381.

Sincerely yours,



Robert S. Brennis  
 Acting Product Manager (32)  
 Regulatory Management Branch II  
 Antimicrobial Division (7510W)

Enclosures

**PURE  
BRIGHT™**

Active Ingredient:  
Sodium Hypochlorite ..... 5.5%  
Inert Ingredients: ..... 94.5%  
TOTAL: ..... 100.0%

**BLEACH**

5% STRONGER FOR COMMERCIAL  
AND INSTITUTIONAL USE

AUTHORIZED BY U.S.D.A. FOR USE IN  
FEDERALLY INSPECTED MEAT AND POULTRY PLANTS

FORMULATED FOR FOODSERVICE APPLICATIONS

KEEP OUT OF REACH OF CHILDREN  
**DANGER**

**STATEMENT OF PRACTICAL TREATMENT**

**If in eyes:** Remove contact lenses and rinse with plenty of water  
for at least 15 minutes. Call a physician.

**If swallowed:** Drink large amounts of water. DO NOT induce  
vomiting. Call a physician or poison control center immediately.

**If on skin:** Immediately remove contaminated clothing and wash  
skin thoroughly with water.

See Side Panel for Additional Precautionary Statements

**Commercial/Institutional Use**

**DISINFECTS • SANITIZES • DEODORIZES**

NET 128 FL. OZ. (1 GAL.) 3.79L

Manufactured by:  
T-Chem Products, Inc.  
9028 Dice Road  
Santa Fe Springs, CA, 90670

EPA Reg. No. 11321-7  
EPA Est. No. 11321-CA-1

97  
MAY 21 A9 55

RECD EPA/OPP/DPD1

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

AUG 14

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
amended for the pesticide,  
registered under EPA Reg. No. 11321-7

5/6/97, Page 1 of 24

### PRECAUTIONARY STATEMENTS

#### Hazards to Humans and Domestic animals

**DANGER:** Corrosive, may cause severe skin and eye irritation or chemical burns to broken skin. Causes eye damage. Wear safety glasses and rubber gloves when handling this product. Wash after handling. Avoid breathing vapors. Vacate poorly ventilated areas as soon as possible. Do not return until strong odors have dissipated.

#### Environmental Hazards

This product is toxic to fish and aquatic organisms. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or public waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

#### Physical and Chemical Hazards

Strong oxidizer. Flush drains before and after use. Do not use or mix with other household chemicals, such as toilet bowl cleaners, rust removers, acid or ammonia containing products. To do so will release hazardous gasses. Prolonged contact with metal may cause pitting or discoloration.

### STORAGE AND DISPOSAL

Store in a cool, dry area away from direct sunlight and heat to avoid deterioration. In case of spill, flood area with large quantities of water. Product or rinsates that cannot be used should be diluted with water before disposal in a sanitary sewer. Do not re-use empty container but place in trash collection. Do not contaminate food or feed by storage, disposal or cleaning of equipment.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

One tablespoon (1/2 ounce) of Pure Bright Bleach in a gallon of water is equivalent to 200 parts per million available chlorine.

### LAUNDRY

Use to bleach white and colorfast Acrylics, Cotton, Nylon, Polyester, Rayon (test to be sure). Do not use on Acetate, Leather, Silk, Spandex or Wool.

**BLEACH TEST:** Before using, mix one tablespoon of bleach with 1/4 cup of water in a glass, rubber, porcelain, or plastic container and test a small piece of fabric in a place that doesn't show.

**LAUNDRY:** Before adding clothes, mix 1 cup of bleach with water in top-loading 16 gallon machines or mix 1/2 cup bleach with water in front-loading 8 gallon machines. Add clothes. If sanitization is desired, soak clothes for 5 minutes. Wash and rinse with usual cycles. If clothes are in machine the addition of bleach can cause damage.

**REMOVE STAINS:** Mix 1/4 cup of bleach with a gallon of water. Soak stained area for 5 minutes to remove grass, ink, coffee, tea, scorch, fruit, etc. Rinse thoroughly.

Disinfects, sanitizes, and deodorizes by killing most germs and their odors.

**HAND SANITIZING:** Thoroughly wash hands with soap or detergent and rinse. Mix 1/2 tablespoon of bleach per gallon of water. Submerge hands. Let air dry.

**SANITIZING FOOD CONTACT SURFACES:** Before using this product, remove or carefully protect food. Prewash surface. Mix 1 tablespoon of bleach per gallon of water and cover surface with bleach solution for at least 2 minutes. Air dry.

**DISINFECTING KITCHEN, DISHES, SINKS:** Use 1/4 cup bleach mixed with a quart of water to soak cleaned dishes, teapot, cups, sinks, etc. for 10 minutes. Rinse with a solution of 1 tablespoon of bleach per gallon of water. Do not use on silverware. Bleach solution can be used on porcelain, enamel, etc. surfaces after cleaning. Let air dry.

**DISINFECTING WALLS, FLOORS, AND OTHER SURFACES NOT IN DIRECT CONTACT WITH FOOD:** Prewash surfaces and rinse. Mix 1/2 cup bleach per gallon of water. Spray, rinse, or wipe surface with bleach solution and let stand for 10 minutes. Drain and air dry.

**DISINFECTING BATHROOM:** Prewash toilet and flush. Pour 1/2 cup bleach into toilet bowl, scrub with brush, let stand 10 minutes, flush. Do not use with bowl cleaners or any other household chemicals.

**DEODORIZING GARBAGE CANS:** Wash and rinse. Use 1/2 cup bleach for each gallon of water in can. Empty and let drain.

**EGG SHELL SANITIZING:** Thoroughly clean eggs. Mix 1 tablespoon (1/2 ounce) of bleach per gallon of warm water to produce a 200 ppm available chlorine solution. The sanitizer temperature should not exceed 130°F. Spray the warm sanitizer so that the eggs are thoroughly wetted. Allow the eggs to thoroughly dry before casing or breaking. Do not apply a potable rinse. The solution should not be re-used to sanitize eggs.

APPROVED  
with OMBATTM  
in EPA Letter dated:

g:\labels\chem\5-5%.lbl

5/6/97, Page 2 of 24

AUG 4 1997  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
this product has been found to be  
effective, safe, and of good quality  
registered under FIFRA Reg. No. 11321-7

4  
7  
26

NOTE: This product degrades with age. Use a chlorine test kit and increase dosage, as necessary, to obtain the required level of available chlorine.

### SWIMMING POOL WATER DISINFECTION

For a new pool or spring start-up, superchlorinate with 117 to 233 oz. of product for each 10,000 gallons of water to yield 5 to 10 ppm available chlorine by weight. Check the level of available chlorine with a test kit. Adjust and maintain pool water pH to between 7.2 to 7.6. Adjust and maintain the alkalinity of the pool to between 50 to 100 ppm.

To maintain the pool, add manually or by a feeder device 14 to 24 oz. of this product for each 10,000 gallons of water to yield an available chlorine residual between 0.6 to 1.0 ppm by weight. Stabilized pools should maintain a residual of 1.0 to 1.5 ppm available chlorine. Test the pH, available chlorine residual and alkalinity of the water frequently with appropriate test kits. Frequency of water treatment will depend upon temperature and number of swimmers.

Every 7 days, or as necessary, superchlorinate the pool with 117 to 233 oz. of product for each 10,000 gallons of water to yield 5 to 10 ppm available chlorine by weight. Check the level of available chlorine with a test kit. Do not reenter pool until the chlorine residual is between 1.0 to 3.0 ppm.

At the end of the swimming pool season or when water is to be drained from the pool, chlorine must be allowed to dissipate from treated pool water before discharge. Do not chlorinate the pool within 24 hours prior to discharge.

WINTERIZING POOLS - While water is still clear & clean, apply 7 oz. of product per 1000 gallons, while filter is running, to obtain a 3 ppm available chlorine residual, as determined by a suitable test kit. Cover pool, prepare heater, filter and heater components for winter by following manufacturers' instructions.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

AUG 14 1997

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under EPA Reg. No. 11321-7

SPAS, HOT-TUBS, IMMERSION TANKS, ETC.

SPAS/HOT-TUBS - Apply 12 oz. of product per 1000 gallons of water to obtain a free available chlorine concentration of 5 ppm, as determined by a suitable chlorine test kit. Adjust and maintain pool water pH to between 7.2 and 7.8. Some oils, lotions, fragrances, cleaners, etc. may cause foaming or cloudy water as well as reduce the efficiency of the product.

To maintain the water, apply 12 oz. of product per 1000 gallons of water over the surface to maintain a chlorine concentration of 5 ppm.

After each use, shock treat with 18 oz. of this product per 500 gallons of water to control odor and algae.

During extended periods of disuse, add 7 oz. of product daily per 1000 gallons of water to maintain a 3 ppm chlorine concentration.

-----  
HUBBARD AND IMMERSION TANKS - Add 12 oz. of this product per 200 gallons of water before patient use to obtain a chlorine residual of 25 ppm, as determined by a suitable test kit. Adjust and maintain the water pH to between 7.2 and 7.6. After each use drain the tank. Add 13 oz. to a bucket of water and circulate this solution through the agitator of the tank for 15 minutes and then rinse out the solution. Clean tank thoroughly and dry with clean cloths.

-----  
HYDROTHERAPY TANKS - Add 3 oz. of this product per 1000 gallons of water to obtain a chlorine residual of 1 ppm, as determined by a suitable chlorine test kit. Pool should not be entered until the chlorine residual is below 3 ppm. Adjust and maintain the water pH to between 7.2 and 7.6. Operate pool filter continuously. Drain pool weekly, and clean before refilling.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

AUG 14 1997

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
amended, for the pesticide  
registered under EPA Reg. No. 11321-7

### SANITIZATION OF NONPOROUS NON-FOOD CONTACT SURFACES

**RINSE METHOD** - Prepare a sanitizing solution by thoroughly mixing 5 oz. of this product with 10 gallons of water to provide approximately 200 ppm available chlorine by weight. Clean equipment surfaces in the normal manner. Prior to use, rinse all surfaces thoroughly with the sanitizing solution, maintaining contact with the sanitizer for at least 2 minutes. Do not rinse equipment with water after treatment and do not soak equipment overnight.

**IMMERSION METHOD** - Prepare a sanitizing solution by thoroughly mixing, in an immersion tank, 5 oz. of this product with 10 gallons of water to provide approximately 200 ppm available chlorine by weight. Clean equipment in the normal manner. Prior to use, immerse equipment in the sanitizing solution for at least 2 minutes and allow the sanitizer to drain. Do not rinse equipment with water after treatment.

**SPRAY/FOG METHOD** - Preclean all surfaces after use. Prepare a 200 ppm. available chlorine sanitizing solution of sufficient size by thoroughly mixing the product in a ratio of 5 oz. product with 10 gallons of water. Use spray or fogging equipment which can resist hypochlorite solutions. Prior to using equipment, thoroughly spray or fog all surfaces until wet, allowing excess sanitizer to drain. Vacate area for at least 2 hours.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

AUG 14 1991

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
amended, for the pesticide  
registered under EPA Reg No

11321-7

### SANITIZATION OF POROUS NON-FOOD CONTACT SURFACES

RINSE METHOD - Prepare a sanitizing solution by thoroughly mixing 14 oz. of this product with 10 gallons of water to provide approximately 600 ppm available chlorine by weight. Clean surfaces in the normal manner. Prior to use, rinse all surfaces thoroughly with the sanitizing solution, maintaining contact with the sanitizer for at least 2 minutes. Do not rinse equipment with water after treatment and do not soak equipment overnight.

IMMERSION METHOD - Prepare a sanitizing solution by thoroughly mixing, in an immersion tank, 14 oz. of this product with 10 gallons of water to provide approximately 600 ppm available chlorine by weight. Clean equipment in the normal manner. Prior to use, immerse equipment in the sanitizing solution for at least 2 minutes and allow the sanitizer to drain. Do not rinse equipment with water after treatment.

SPRAY/FOG METHOD - After cleaning, sanitize non-food contact surfaces with 600 ppm available chlorine by thoroughly mixing the product in a ratio of 14 oz. of this product with 10 gallons of water. Use spray or fogging equipment which can resist hypochlorite solutions. Always empty and rinse spray/fog equipment with potable water after use. Prior to using equipment, thoroughly spray or fog all surfaces until wet, allowing excess sanitizer to drain. Vacate area for at least 2 hours.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

MAY 14 1997

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act as  
amended, for the pesticide  
registered under EPA Reg. No. 11321-7



### SEWAGE & WASTEWATER EFFLUENT TREATMENT

The disinfection of sewage effluent must be evaluated by determining the total number of coliform bacteria and/or fecal coliform bacteria, as determined by the Most Probable Number (MPN) procedure, of the chlorinated effluent has been reduced to or below the maximum permitted by the controlling regulatory jurisdiction.

On the average, satisfactory disinfection of secondary wastewater effluent can be obtained when the chlorine residual is 0.5 ppm after 15 minutes contact. Although the chlorine residual is the critical factor in disinfection, the importance of correlating chlorine residual with bacterial kill must be emphasized. The MPN of the effluent, which is directly related to the water quality standards requirements, should be the final and primary standard and the chlorine residual should be considered an operating standard valid only to the extent verified by the coliform quality of the effluent.

The following are critical factors affecting wastewater disinfection.

1. Mixing: It is imperative that the product and the wastewater be instantaneously and completely flash mixed to assure reaction with every chemically active soluble and particulate component of the wastewater.
2. Contacting: Upon flash mixing, the flow through the system must be maintained.
3. Dosaġe/Residual Control: Successful disinfection is extremely dependant on response to fluctuating chlorine demand to maintain a predetermined, desirable chlorine level. Secondary effluent should contain 0.2 to 1.0 ppm chlorine residual after a 15 to 30 minute contact time. A reasonable average of residual chlorine is 0.5 ppm after 15 minutes contact time.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

AUG 14 1997

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act as  
amended, for the pesticide  
registered under EPA Reg. No. 11321-7

### SEWAGE AND WASTEWATER TREATMENT

**EFFLUENT SLIME CONTROL** - Apply a 100 to 1000 ppm available chlorine solution at a location which will allow complete mixing. Prepare this solution by mixing 24 to 233 oz. of this product with 100 gallons of water. Once control is evident, apply a 15 ppm available chlorine solution. Prepare this solution by mixing 4 oz. of this product with 100 gallons of water.

**FILTER BEDS - SLIME CONTROL:** Remove filter from service, drain to a depth of 1 ft. above filter sand, and add 182 oz. of product per 20 sq/ft evenly over the surface. Wait 30 minutes before draining water to a level that is even with the top of the filter. Wait for 4 to 6 hours before completely draining and backwashing filter.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

AUG 14 1997

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act as  
amended, for the pesticide  
registered under EPA Reg. No. 11321-7

DISINFECTION OF DRINKING WATER  
(EMERGENCY/PUBLIC/INDIVIDUAL/SYSTEMS)

PUBLIC SYSTEMS - Mix a ratio of 1 oz. of this product to 100 gallons of water. Begin feeding this solution with a hypochlorinator until a free available chlorine residual of at least 0.2 ppm and no more than 0.6 ppm is attained throughout the distribution system. Check water frequently with a chlorine test kit. Bacteriological sampling must be conducted at a frequency no less than that prescribed by the National Interim Primary Drinking Water Regulations. Contact your local Health Department for further details.

INDIVIDUAL SYSTEMS: DUG WELLS - Upon completion of the casing (lining), wash the interior of the casing (lining) with a 100 ppm available chlorine solution using a stiff brush. This solution can be made by thoroughly mixing 3 oz. of this product into 10 gallons of water. After covering the well, pour the sanitizing solution into the well through both the pipesleeve opening and the pipeline. Wash the exterior of the pump cylinder also with the sanitizing solution. Start pump and pump water until strong odor of chlorine in water is noted. Stop pump and wait at least 24 hours. After 24 hours flush well until all traces of chlorine have been removed from the water. Consult your local Health Department for further details.

INDIVIDUAL WATER SYSTEMS: DRILLED, DRIVEN & BORED WELLS - Run pump until water is as free from turbidity as possible. Pour a 100 ppm available chlorine sanitizing solution into the well. This solution can be made by thoroughly mixing 3 oz. of this product into 10 gallons of water. Add 5 to 10 gallons of clean, chlorinated water to the well in order to force the sanitizer into the rock formation. Wash the exterior of the pump cylinder with the sanitizer. Drop pipeline into the well, start pump and pump water until strong odor of chlorine in water is noted. Stop pump and wait at least 24 hours. After 24 hours, flush well until all traces of chlorine have been removed from the water. Deep wells with high water levels may necessitate the use of special methods for introduction of the sanitizer to the well. Consult your local Health Department for further details.

INDIVIDUAL WATER SYSTEMS: FLOWING ARTESIAN WELLS - Artesian Wells generally do not require disinfection. If analyses indicate persistent contamination, the well should be disinfected. Consult your local Health Department for further details.

EMERGENCY DISINFECTION - When boiling of water for 1 minute is not practical, water can be made potable by using this product. Prior to addition of the sanitizer, remove all suspended material by filtration or by allowing it to settle to the bottom. Decant the clarified, contaminated water to a clean container and add 20 drops of this product to 20 gallons of water. Allow the treated water to stand for 30 minutes. Properly treated water should have a slight chlorine odor, if not, repeat dosage and allow the water to stand an additional 15 minutes. The treated water can then be made palatable by pouring it between clean containers for several times. This process has not been demonstrated to inactivate Cryptosporidium cysts.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

AUG 14 1997

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
amended, for the pesticide  
registered under EPA Reg. No. 11321-7

### PUBLIC WATER SYSTEMS

RESERVOIRS: ALGAE CONTROL - Hypo chlorinate streams feeding the reservoir. Suitable feeding points should be selected on each stream at least 50 yards upstream from the points of entry into the reservoir.

MAINS - Thoroughly flush section to be sanitized by discharging from hydrants. Permit a water flow of at least 2.5 feet per minute to continue under pressure while injecting this product by means of a hypochlorinator. Stop water flow when a chlorine residual test of 50 ppm is obtained at the low pressure end of the new main section after a 24 hour retention time. When chlorination is completed, the system must be flushed free of all heavily chlorinated water.

NEW TANKS, BASINS, ETC. - Remove all physical soil from surfaces. Place 46 oz. of this product for each 5 cubic feet of working capacity (500 ppm available chlorine). Fill to working capacity and allow to stand for at least 4 hours. Drain and flush with potable water and return to surface.

NEW FILTER SAND - Apply 182 oz. of this product for each 150 to 200 cubic feet of sand. The action of the product dissolving as the water passes through the bed will aid in sanitizing the new sand.

NEW WELLS - Flush the casing with a 50 ppm available chlorine solution of water containing 12 oz. of this product for each 100 gallons of water. The solution should be pumped or fed by gravity into the well after thorough mixing with agitation. The well should stand for several hours or overnight under chlorination. It may then be pumped until a representative raw water sample is obtained. Bacterial examination of the water will indicate whether further treatment is necessary.

EXISTING EQUIPMENT - Remove equipment from service, thoroughly clean surfaces of all physical soil. Sanitize by placing 48 oz. of this product for each 5 cubic feet capacity (approximately 500 ppm available chlorine). Fill to working capacity and let stand at least 4 hours. Drain and place in service. If the previous treatment is not practical, surfaces may be sprayed with a solution containing 12 oz. of this product for each 5 gallons of water (approximately 1000 ppm available chlorine). After drying, flush with water and return to service.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

AUG 14 1997

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act as  
amended, for the pesticide  
registered under EPA Reg. No. 11321-7

## EMERGENCY DISINFECTION AFTER FLOODS

WELLS - Thoroughly flush contaminated casing with a 500 ppm available chlorine solution. Prepare this solution by mixing 12 oz. of this product with 10 gallons of water. Backwash the well to increase yield and reduce turbidity, adding sufficient chlorinating solution to the backwash to produce a 10 ppm available chlorine residual, as determined by a chlorine test kit. After the turbidity has been reduced and the casing has been treated, add sufficient chlorinating solution to produce a 50 ppm available chlorine residual. Agitate the well water for several hours and take a representative water sample. Retreat well if water samples are biologically unacceptable.

RESERVOIRS - In case of contamination by overflowing streams, establish hypochlorinating stations upstream of the reservoir. Chlorinate the inlet water until the entire reservoir obtains a 0.2 ppm available chlorine residual, as determined by a suitable chlorine test kit. In case of contamination from surface drainage, apply sufficient product directly to the reservoir to obtain a 0.2 ppm available chlorine residual in all parts of the reservoir.

BASINS, TANKS, FLUMES, ETC. - Thoroughly clean all equipment, then apply 46 oz. of product per 5 cu. ft. of water to obtain 500 ppm available chlorine, as determined by a suitable test kit. After 24 hours, drain, flush, and return to service. If the previous method is not suitable, spray or flush the equipment with a solution containing 12 oz. of this product for each 5 gallons of water (1000 ppm available chlorine). Allow to stand for 2 to 4 hours, flush and return to service.

FILTERS - When the sand filter needs replacement, apply 182 oz. of this product for each 150 to 200 cubic feet of sand. When the filter is severely contaminated, additional product should be distributed over the surface at the rate of 182 oz. per 20 sq.ft. Water should stand at a depth of 1 foot above the surface of the filter bed for 4 to 24 hours. When filter beds can be backwashed of mud and silt, apply 182 oz. of this product per each 50 sq. ft., allowing the water to stand at a depth of 1 foot above the filter sand. After 30 minutes, drain water to the level of the filter. After 4 to 6 hours, drain, and proceed with normal backwashing.

DISTRIBUTION SYSTEM - Flush repaired or replaced section with water. Establish a hypochlorinating station and apply sufficient product until a consistent available chlorine residual of at least 10 ppm remains after 24 hour retention time. Use a chlorine test kit.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

AUG 14 1997

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
amended, for the pesticide  
registered under EPA Reg. No. 11321-7

EMERGENCY DISINFECTION AFTER FIRES

CROSS CONNECTIONS OF EMERGENCY CONNECTIONS - Hypochlorination or gravity feed equipment should be set up near the intake of the untreated water supply. Apply sufficient product to give a chlorine residual of at least 0.1 to 0.2 ppm at the point where the untreated supply enters the regular distribution system. Use a chlorine test kit.

EMERGENCY DISINFECTION AFTER DROUGHTS

SUPPLEMENTARY WATER SUPPLIES - Gravity or mechanical hypochlorite feeders should be set up on a supplementary line to dose the water to a minimum chlorine residual of 0.2 ppm after a 20 minute contact time. Use a chlorine test kit.

WATER SHIPPED IN BY TANKS, TANK CARS, TRUCKS, ETC. - Thoroughly clean all containers and equipment. Spray a 500 ppm available chlorine solution and rinse with potable water after 5 minutes. This solution is made by mixing 12 oz. of this product for each 10 gallons of water. During the filling of the containers, dose with sufficient amounts of this product to provide at least a 0.2 ppm chlorine residual. Use a chlorine test kit.

EMERGENCY DISINFECTION AFTER MAIN BREAKS

MAINS - Before assembly of the repaired section, flush out mud and soil. Permit a water flow of at least 2.5 feet per minute to continue under pressure while injecting this product by means of a hypochlorinator. Stop water flow when a chlorine residual of test of 50 ppm is obtained at the low pressure end of the new main section after a 24 hour retention time. When chlorination is completed, the system must be flushed free of all heavily chlorinated water.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

AUG 1 1997

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act as  
amended, for the pesticide  
registered under EPA Reg. No. 11321-7

### COOLING TOWER/EVAPORATIVE CONDENSER WATER

**SLUG FEED METHOD - Initial Dose:** When system is noticeably fouled, apply 117 to 233 oz. of this product per 10,000 gallons of water in the system to obtain from 5 to 10 ppm available chlorine. Repeat until control is achieved.

**Subsequent Dose:** When microbial control is evident, add 24 oz. of this product per 10,000 gallons of water in the system daily, or as needed to maintain control and keep the chlorine residual at 1 ppm. Badly fouled systems must be cleaned before treatment is begun.

**INTERMITTENT FEED METHOD - Initial Dose:** When system is noticeably fouled, apply 117 to 233 oz. of this product per 10,000 gallons of water in the system to obtain 5 to 10 ppm available chlorine. Apply half (or 1/3, 1/4, or 1/5) of this initial dose when half (or 1/3, 1/4, or 1/5) of the water in the system has been lost by blowdown.

**Subsequent Dose:** When microbial control is evident, add 24 oz. of this product per 10,000 gallons of water in the system to obtain a 1 ppm residual. Apply half (or 1/3, 1/4, or 1/5) of this initial dose when half (or 1/3, 1/4, or 1/5) of the water in the system has been lost by blowdown. Badly fouled systems must be cleaned before treatment is begun.

**CONTINUOUS FEED METHOD - Initial Dose:** When system is noticeably fouled, apply 117 to 233 oz. of this product per 10,000 gallons of water in the system to obtain 5 to 10 ppm available chlorine.

**Subsequent Dose:** Maintain his treatment level by starting a continuous feed of 3 oz. of this product per 1,000 gallons of water lost by blowdown to maintain a 1 ppm residual. Badly fouled systems must be cleaned before treatment is begun.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

AUG 14 1997

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under EPA Reg. No. 11321-7

16 7 26

### FARM PREMISES

Remove all animals, poultry, and feed from premises, vehicles, and enclosures. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities occupied or transverse by animals or poultry. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. To disinfect, saturate all surfaces with a solution of at least 1000 ppm available chlorine for a period of 10 minutes. A 1000 ppm solution can be made by thoroughly mixing 24 oz. of this product with 10 gallons of water. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals or poultry, as well as the cleaned forks, shovels, and scrapers used for removing litter and manure. Ventilate buildings, cars, boats, and other closed spaces. Do not house livestock or poultry or employ equipment until chlorine has been dissipated. All treated feed racks, mangers, troughs, automatic feeders, fountains, and waterers must be rinsed with potable water before reuse.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

AUG 14 1987

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
amended, for the pesticide  
registered under EPA Reg. No. 11321-7



### PULP AND PAPER MILL PROCESS WATER SYSTEMS

**SLUG FEED METHOD - Initial Dose:** When system is noticeably fouled, apply 117 to 233 oz. of this product per 10,000 gallons of water in the system to obtain from 5 to 10 ppm available chlorine. Repeat until control is achieved.

**Subsequent Dose:** When microbial control is evident, add 24 oz. of this product per 10,000 gallons of water in the system daily, or as needed to maintain control and keep the chlorine residual at 1 ppm. Badly fouled systems must be cleaned before treatment is begun.

**INTERMITTENT FEED METHOD - Initial Dose:** When system is noticeably fouled, apply 117 to 233 oz. of this product per 10,000 gallons of water in the system to obtain 5 to 10 ppm available chlorine. Apply half (or 1/3, 1/4, or 1/5) of this initial dose when half (or 1/3, 1/4, or 1/5) of the water in the system has been lost by blowdown.

**Subsequent Dose:** When microbial control is evident, add 24 oz. of this product per 10,000 gallons of water in the system to obtain a 1 ppm residual. Apply half (or 1/3, 1/4, or 1/5) of this initial dose when half (or 1/3, 1/4, or 1/5) of the water in the system has been lost by blowdown. Badly fouled systems must be cleaned before treatment is begun.

**CONTINUOUS FEED METHOD - Initial Dose:** When system is noticeably fouled, apply 117 to 233 oz. of this product per 10,000 gallons of water in the system to obtain 5 to 10 ppm available chlorine.

**Subsequent Dose:** Maintain this treatment level by starting a continuous feed of 3 oz. of this product per 1,000 gallons of water lost by blowdown to maintain a 1 ppm residual. Badly fouled systems must be cleaned before treatment is begun.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act as  
amended, for the pesticide  
registered under EPA Reg. No. 11321-7

AGRICULTURAL USES

POST-HARVEST PROTECTION - Potatoes can be sanitized after cleaning and prior to storage by spraying with a sanitizing solution at a level of 1 gallon of sanitizing solution per ton of potatoes. Thoroughly mix 3 oz. of this product to 2 gallons of water to obtain 500 ppm available chlorine.

Disinfect leafcutting bee cells and bee boards by immersion in a solution containing 1 ppm available chlorine for 3 minutes. Allow cells to drain for 2 minutes and dry for 4 to 5 hours or until no chlorine odor can be detected. This solution is made by thoroughly mixing 2¼ tsp. of this product to 100 gallons of water. The bee domicile is disinfected by spraying with a 0.1 ppm solution until all surfaces are thoroughly wet. Allow the domicile to dry until all chlorine odor has dissipated. [Not for use in California]

FRUIT & VEGETABLE WASHING - Thoroughly clean all fruits and vegetables in a wash tank. Thoroughly mix 12 oz. of this product in 200 gallons of water to make a sanitizing solution of 25 ppm available chlorine. After draining the tank, submerge fruit or vegetables for 2 minutes in a second wash tank containing the recirculating sanitizing solution. Spray rinse vegetables with the sanitizing solution prior to packaging. Rinse fruit with potable water only prior to packaging.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

AUG 14 1997

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act as  
amended, for the pesticide  
registered under EPA Reg. No. 11321-7

### AQUACULTURAL USES

FISH PONDS - Remove fish from ponds prior to treatment. Thoroughly mix 233 oz. of this product to 10,000 gallons of water to obtain 10 ppm available chlorine. Add more product to the water if the available chlorine level is below 1 ppm after 5 minutes. Return fish to pond after the available chlorine level reaches zero.

FISH POND EQUIPMENT - Thoroughly clean all equipment prior to treatment. Thoroughly mix 5 oz. of this product to 10 gallons of water to obtain 200 ppm available chlorine. Porous equipment should soak for one hour.

MAINE LOBSTER PONDS - Remove lobsters, seaweed, etc. from ponds prior to treatment. Drain the pond. Thoroughly mix 13,964 oz. of this product to 10,000 gallons of water to obtain at least 600 ppm available chlorine. Apply so that all barrows, gates, rocks and dams are treated with product. Permit high tide to fill the pond and then close the gates. Allow water to stand for 2 to 3 days until the available chlorine level reaches zero. Open and allow 2 tidal cycles to flush the pond before returning lobsters to the pond.

CONDITIONING LIVE OYSTERS - Thoroughly mix 12 oz. of this product to 10,000 gallons of water at 50 to 70 degrees F to obtain 0.5 ppm available chlorine. Expose oysters to this solution for at least 15 minutes, monitoring the available chlorine level so that it does not fall below 0.05 ppm. Repeat entire process if the available chlorine level drops below 0.05 ppm or the temperature falls below 50 degrees F. [Not for use in California]

CONTROL OF SCAVENGERS IN FISH HATCHERY PONDS - Prepare a solution containing 200 ppm of available chlorine by mixing 5 oz. of product with 10 gallons of water. Pour into drained pond potholes. Repeat if necessary. Do not put desirable fish back into refilled ponds until chlorine residual has dropped to 0 ppm, as determined by a test kit.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

AUG 14 1997  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act as  
amended, for the pesticide  
registered under EPA Reg. No 11321-7

### SANITIZATION OF DIALYSIS MACHINES

Flush equipment thoroughly with water prior to using this product. Thoroughly mix 14 oz. of this product to 10 gallons of water to obtain at least 600 ppm available chlorine. Immediately use this product in the hemodialysate system allowing for a minimum contact time of 15 minutes at 20 degrees C. Drain system of the sanitizing solution and thoroughly rinse with water. Discard and DO NOT reuse the spent sanitizer. Rinsate must be monitored with a suitable test kit to ensure that no available chlorine remains in the system.

This product is recommended for decontaminating single and multipatient hemodialysate systems. This product has been shown to be an effective disinfectant (virucide, fungicide, bactericide, pseudomonicide) when tested by AOAC and EPA test methods. This product may not totally eliminate all vegetative microorganisms in hemodialysate delivery systems due to their construction and/or assembly, but can be relied upon to reduce the number of microorganisms to acceptable levels when used as directed. This product should be used in a disinfectant program which includes bacteriological monitoring of the hemodialysate delivery system. This product is NOT recommended for use in hemodialysate or reverse osmosis (RO) membranes.

Consult the guidelines for hemodialysate systems which are available from the Hepatitis Laboratories, CDC, Phoenix, AR 85021.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act as  
amended, for the pesticide  
registered under EPA Reg. No. 11321-7

### ASPHALT OR WOOD ROOFS AND SIDINGS

To control fungus and mildew, first remove all physical soil by brushing and hosing with clean water, and apply a 5000 ppm available chlorine solution. Mix 12 oz. of this product per gallon of water and brush or spray roof or siding. After 30 minutes, rinse by hosing with clean water. [Not for Use in California]

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act as  
amended, for the pesticide  
registered under EPA Reg. No. 11321-7

### BOAT BOTTOMS

To control slime on boat bottoms; sling a plastic tarp under boat, retaining enough water to cover the fouled bottom area, but not allowing water to enter enclosed area. This envelope should contain approximately 500 gallons of water for a 14 foot boat. Add 41 oz. of this product to this water to obtain a 35 ppm available chlorine concentration. Leave immersed for 8 to 12 hours. Repeat if necessary. Do not discharge the solution until the free chlorine level has dropped to 0 ppm, as determined by a swimming pool test kit. [Not for Use in California]

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
amended, for the pesticide  
registered under EPA Reg. No. 11321-7

23 7 26

### ARTIFICIAL SAND BEACHES

To sanitize the sand, spray a 500 ppm available chlorine solution containing 12 oz. of this product per 10 gal. of water at frequent intervals. Small areas can be sprinkled with a watering can. [Not for Use in California]

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

AUG 1 9

Under the Federal Insecticide  
Fungicide, and Rodenticide Act as  
amended, for the pesticide  
registered under EPA Reg. 1: 11321-7

### WAREWASHING

FOR SANITIZING TABLEWARE IN LOW TEMPERATURE DISHWASHING MACHINE -  
Dispense Pure Bright Institutional Sanitizer into final rinse water at 100 ppm available chlorine. Do not allow concentration to fall below 50 ppm. Air dry. Dispenser should be set to deliver 7cc of sanitizing solution per gallon of water to give 100 ppm of available chlorine. Only a qualified service representative should set or adjust dispenser on the machine.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

AUG 4 1997

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
amended, for the pesticide  
registered under EPA Reg. No. 11321-9



**SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV (AIDS VIRUS) OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.**

Kills HIV-1 (AIDS virus) on surfaces soiled with blood/body fluids in healthcare settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of HIV.

**Personal Protection:** Disposable latex or vinyl gloves, gowns, masks, and/or eye coverings as appropriate must be worn during all cleaning and decontamination procedures of blood and other body fluids.

**Cleaning Procedures:** Remove visible blood and other body fluids before applying disinfectant.

**Disinfectant Use and Contact Time:** Effective against HIV-1 (AIDS virus) on hard non-porous surfaces in the presence of a moderate amount of organic soil (e.g. 5% blood serum). Prepare disinfectant by mixing 24 oz. of this product per gallon of water to provide 10,000 ppm of available chlorine. Leave surfaces wet for 5 minutes. Drain and let air dry.

**Disposal of Infectious Materials:** Blood and other body fluids should be autoclaved and disposed of according to Federal, State, and local regulations for infectious waste disposal.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

AUG 14 1997  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act as  
amended, for the pesticide  
registered under EPA Reg. No. 11321-7

**SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST MYCOBACTERIUM TUBERCULOSIS (TB) OF SURFACES/OBJECTS SOILED WITH BODY FLUIDS.**

Kills Mycobacterium tuberculosis (TB) on surfaces soiled with body fluids in healthcare settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with body fluids, and in which the surfaces/objects likely to be soiled with body fluids can be associated with the potential for transmission of Mycobacterium tuberculosis.

**Personal Protection:** Disposable latex or vinyl gloves, gowns, masks, and/or eye coverings as appropriate must be worn during all cleaning and decontamination procedures of body fluids.

**Cleaning Procedures:** Remove visible body fluids before applying disinfectant.

**Disinfectant Use and Contact Time:** Effective against Mycobacterium tuberculosis on hard non-porous surfaces in the presence of a moderate amount of organic soil (e.g. sputum). Prepare disinfectant by mixing 24 oz. of this product per gallon of water to provide 10,000 ppm of available chlorine. Leave surfaces wet for 5 minutes. Drain and let air dry.

**Disposal of Infectious Materials:** Blood and other body fluids should be autoclaved and disposed of according to Federal, State, and local regulations for infectious waste disposal.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

AUG 14 1997

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act as  
amended, for the pesticide  
registered under EPA Reg. No. 11321-7