

Cold Disinfection

For disinfection of previously cleaned surgical instruments, barber tools and dental equipment, simply submerge instruments into solution containing 1-1/2 ounces RID ALL DISINFECTANT DEODORANT SANITIZER per gallon of water for 10 minutes.

Restaurant and Bar Rinse

Dishes, glassware, silverware, cooking utensils, wash with soap or synthetic detergent; rinse thoroughly and immerse in sanitizing solution containing 1 ounce RID ALL DISINFECTANT DEODORANT SANITIZER to 4 gallons of water.

Potable Water Rinse Required.

DANGER

Keep Out of Reach of Children. Corrosive. Causes eye damage and severe skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling concentrate. Harmful if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

Rinse empty container thoroughly with water and discard.

ACCEPTED
OCT 12 1973
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 11327-6 SUBJECT
TO ATTACHED COMMENTS.

CHEMCO

RID ALL DISINFECTANT DEODORANT SANITIZER

ACTIVE INGREDIENTS:

n-Alkyl (60% C14, 30% C16, 5% C12, 5% C18)
dimethyl benzyl ammonium chlorides.....5%

n-Alkyl (68% C12, 32% C14)
dimethyl ethylbenzyl ammonium chlorides...5%

INERT INGREDIENTS:90%
100%

RID ALL DISINFECTANT DEODORANT SANITIZER is a concentrate and should be diluted before using.

NET CONTENTS: _____ EPA Reg. No. 11327-6

DANGER

Keep Out of Reach of Children. See other precautionary and first aid statements on left side panel.



INDUSTRIES INCORPORATED

Manufacturing Chemists

PHONE (813) 626-2178 P. O. BOX 2635

7501 ADAMO DRIVE TAMPA, FLORIDA 33601

Garbage Pails, Telephone Restrooms, use RID ALL DISINFECTANT DEODORANT SANITIZER at 3-1/2 ounces per 5 gallons of water. Apply with cloth, mop or sponge. At this level RID ALL DISINFECTANT DEODORANT SANITIZER will be effective against Pseudomonas aeruginosa PRD-10.

Hospitals

For floors, walls and other hard surfaces. In Nursing Homes, Institutions, as well as Sink Tops, Garbage Pails, Telephones and Restrooms, use RID ALL DISINFECTANT DEODORANT SANITIZER at 3-1/2 ounces per 5 gallons of water. Apply with cloth, mop or sponge. At this level RID ALL DISINFECTANT DEODORANT SANITIZER will be effective against Pseudomonas aeruginosa PRD-10.

Food Processing Equipment

For sanitization of previously cleaned food processing equipment and food utensils dilute 1 ounce RID ALL DISINFECTANT DEODORANT SANITIZER per 4 gallons of water to provide 200 ppm of active quaternaries. At this level, No Potable Water Rinse is required.

Dairies

To sanitize dairy equipment such as tanks, lines, pails and milk cans, first clean and rinse the equipment thoroughly. Then apply sanitizing solution containing 1 ounce RID ALL DISINFECTANT DEODORANT SANITIZER to 4 gallons of water (200ppm). At this level, no Potable Water Rinse is required. Follow recommendations of local Health Board.

ACCEPTED
OCT 12, 1973
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 11327-6 SUBJECT
TO ATTACHED COMMENTS.

