

JUMBO CHLORINE TABLETS

DANGER!

STRONG OXIDIZING AGENT

DANGER

KEEP OUT OF REACH OF CHILDREN

Corrosive, causes eye damage. May be fatal if swallowed. Do not get in eyes, on skin or on clothing. Irritating to nose and throat. Avoid breathing dust. May cause burns to broken skin. Wash hands after handling.

FIRST AID:

If swallowed feed gruel, cooked cereal, or bread soaked in milk, followed by olive oil or cooking oil. In case of personal contact, remove excess material and flush skin or eyes with cold water for at least 15 minutes. For eyes get medical attention. Remove and wash contaminated clothing before reuse.

Mix only with water. Use clean dry utensils. Contamination by moisture, organic matter, or other chemicals may liberate hazardous gases. Store in a cool, dry well ventilated area away from heat or open flame. Decomposes at 350° - 475°F with liberation of harmful gases. In case of decomposition, if possible, isolate container in open air. Flood with large amounts of water. Keep container tightly closed when not in use. Do not reuse empty container. Rinse empty container thoroughly with water to dissolve all material before discarding.

ENVIRONMENTAL HAZARDS:

This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds, or public waters. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label.

DIRECTIONS FOR USE

When starting up a new pool, super-chlorinate to satisfy chlorine demand and to establish a chlorine residual of 1.0 ppm to 2.0 ppm available chlorine. To accomplish this, add 2 oz. SUPER CLEARON Concentrated Pool Chlorine per 1000 gallons of pool water. The next morning, add cyanuric acid at the rate of 4 oz. per 1000 gallons to stabilize pool. For pools with stabilizer the pool should be maintained between 1.0 - 1.5 ppm.

A shock treatment should be made every week during hot weather or after heavy rains. Less frequent shock treatments may be made during cool weather. For the shock treatment, add 2 oz. of SUPER CLEARON Concentrated Pool Chlorine per 1000 gallons of pool water. Swimmers should not be permitted in the pool after the shock treatment until the chlorine has reached the 1.0 ppm to 2.0 ppm range.

Keep in mind that a slight flexibility must be maintained in the event of heavy accumulations of dust and dirt caused by storms or super heavy bathing loads which require additional dosages as needed to maintain the chlorine residual between 1.0 ppm and 2.0 ppm.

JUMBO should be at the rate to 10,000 maintain times of mined by should and 7.6.

These t and are high pH Water a eyes an scale for ing syst irritation effective

Seller m or impl this pro the labe use and, when su contrary

E.P.A. R

PACKAGI

ALA
Hig
R

ACCEPTED
11329-9

CHLORINE TABLETS

DANGER!

STRONG OXIDIZING AGENT

Mix only with water. Use clean dry utensils. Contamination by moisture, organic matter, or other chemicals may liberate hazardous gases. Store in a cool, dry well ventilated area away from heat or open flame. Decomposes at 350° - 475°F with liberation of harmful gases. In case of decomposition, if possible, isolate container in open air. Flood with large amounts of water. Keep container tightly closed when not in use. Do not reuse empty container. Rinse empty container thoroughly with water to dissolve all material before discarding.

ENVIRONMENTAL HAZARDS:

This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds, or public waters. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label.

DIRECTIONS FOR USE

When starting up a new pool, super-chlorinate to satisfy chlorine demand and to establish a chlorine residual of 1.0 ppm to 2.0 ppm available chlorine. To accomplish this, add 2 oz. SUPER CLEARON Concentrated Pool Chlorine per 1000 gallons of pool water. The next morning, add cyanuric acid at the rate of 4 oz. per 1000 gallons to stabilize pool. For pools with stabilizer the pool should be maintained between 1.0 - 1.5 ppm.

A shock treatment should be made every week during hot weather or after heavy rains. Less frequent shock treatments may be made during cool weather. For the shock treatment, add 2 oz. of SUPER CLEARON Concentrated Pool Chlorine per 1000 gallons of pool water. Swimmers should not be permitted in the pool after the shock treatment until the chlorine has reached the 1.0 ppm to 2.0 ppm range.

Keep in mind that a slight flexibility must be maintained in the event of heavy accumulations of dust and dirt caused by storms or super heavy bathing loads which require additional dosages as needed to maintain the chlorine residual between 1.0 ppm and 2.0 ppm.

JUMBO CHLORINE TABLETS

should be added daily or as needed at the rate of 1 (one) tablet per 5,000 to 10,000 gallons of pool water to maintain a chlorine residual at all times of 1.0 ppm to 2.0 ppm as determined by a suitable test kit. The pH should be maintained between 7.2 and 7.6.

These tablets are slow dissolving and are acidic, and will balance the high pH of pool water in most areas. Water at pH below 7.0 will burn the eyes and may cause corrosion and scale formation to the pool or filtering system. At pH above 8.0 eye irritation also occurs and chlorine effectiveness rapidly diminishes.

NOTICE

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of the material when such use and/or handling is contrary to label instructions.

E.P.A. REG. 11329-9

PACKAGED BY:

ALADDIN CHEMICAL CO.

Highway 152 - P.O. Box 244
Rogers, Minnesota 55374

