

NET WT. 20 LBS.

GREENWAY WEED AND FEED PLUS IRON

KILLS WEEDS AS IT FEEDS YOUR LAWN

DIRECTIONS

Apply anytime during growing season—Spring, Summer
or Fall. Always use a mechanical spreader.

SPREADER SETTINGS—On standard models—Set all makes at
about 1/5 of total opening.

Supergreen Models 1622 B, 1622 C.....	5½
Supergreen Models 1622 E, 1622 F.....	6
Central, Greenfield, Cross Country.....	5½
Scotts-Standard Models Only.....	4

These settings are furnished as a guide **ONLY**—Age, condition of spreader, speed
of operation and roughness of terrain can cause variations in rates applied.

HIGHLY EFFECTIVE AGAINST

WILD CARROT	CHICKWEED	WILD ONION	(MORNING GLORY)
CURLED DOCK	BINDWEED	DANDELION	and many other
RAGWEED	PLANTAIN	WILD HONEYSUCKLE	similar broadleaf
		WILD MUSTARD	weeds.

The plant food supplies the quick pick-up needed to fill out bare spots
left by dead weeds and help prevent re-establishment of weeds by
encouraging growth and formation of thick, resistant turf.

BE CAREFUL

- 1) Select a time when rain is not expected. (Rain washes weed killer
off weeds—reducing kill.)
- 2) For best results apply when grass is damp—dew or after watering.
- 3) Apply evenly—avoid drift to and on plants, flowers and shrubs—
it can kill them.
- 4) Chickweeds—Best kill when temperature is low (spring and fall).
A repeat application may be required in two weeks.
- 5) Water 2-3 days after application—this will start the feeding.
Weed killer has started to kill weeds the first day.
- 6) New lawns should have 2-3 months growth before using this
product—DO NOT use on bent grass, St. Augustine Grass, Dichondra, Clover
- 7) Will not burn when applied as recommended.
- 8) Do not store near fertilizers, seeds, insecticides, or fungicides.
- 9) Clean spreader thoroughly after use.
- 10) Do not contaminate water used for irrigation or domestic purposes.

CAUTION

May be harmful if swallowed. Avoid contact with skin or eyes.
Avoid breathing dust. Avoid contamination of foodstuffs or
feeds. Wash thoroughly with soap and water after using.

This product is toxic to fish. Keep out of lakes, streams, or
ponds. Do not contaminate water by cleaning of equipment or
disposal of wastes. Do not reuse empty bag. Dispose of
through regular refuse collection system.

ANTIDOTE—EXTERNAL—In case of spillage on skin, wash
promptly with soap and water.

INTERNAL—Induce vomiting promptly. CALL A PHYSICIAN

U.S. PAT. NO. 3,281,060

WEED AND FEED PLUS IRON

KILLS WEEDS AS IT FEEDS YOUR LAWN

DIRECTIONS

Apply anytime during growing season—Spring, Summer
or Fall. Always use a mechanical spreader.

SPREADER SETTINGS—On standard models—Set all makes at
about 1/5 of total opening.

Supergreen Models 1622 B, 1622 C.....	5½
Supergreen Models 1622 E, 1622 F.....	6
Central, Greenfield, Cross Country.....	5½
Scotts-Standard Models Only.....	4

These settings are furnished as a guide **ONLY**—Age, condition of spreader, speed
of operation and roughness of terrain can cause variations in rates applied.

HIGHLY EFFECTIVE AGAINST

WILD CARROT	CHICKWEED	WILD ONION	(MORNING GLORY)
CURLED DOCK	BINDWEED	DANDELION	and many other
RAGWEED	PLANTAIN	WILD HONEYSUCKLE	similar broadleaf
		WILD MUSTARD	weeds.

The plant food supplies the quick pick-up needed to fill out bare spots
left by dead weeds and help prevent re-establishment of weeds by
encouraging growth and formation of thick, resistant turf.

BE CAREFUL

- 1) Select a time when rain is not expected. (Rain washes weed killer
off weeds—reducing kill.)
- 2) For best results apply when grass is damp—dew or after watering.
- 3) Apply evenly—avoid drift to and on plants, flowers and shrubs—
it can kill them.
- 4) Chickweeds—Best kill when temperature is low (spring and fall).
A repeat application may be required in two weeks.
- 5) Water 2-3 days after application—this will start the feeding.
Weed killer has started to kill weeds the first day.
- 6) New lawns should have 2-3 months growth before using this
product—DO NOT use on bent grass, St. Augustine Grass, Dichondra, Clover
- 7) Will not burn when applied as recommended.
- 8) Do not store near fertilizers, seeds, insecticides, or fungicides.
- 9) Clean spreader thoroughly after use.
- 10) Do not contaminate water used for irrigation or domestic purposes.

CAUTION

May be harmful if swallowed. Avoid contact with skin or eyes.
Avoid breathing dust. Avoid contamination of foodstuffs or
feeds. Wash thoroughly with soap and water after using.

This product is toxic to fish. Keep out of lakes, streams, or
ponds. Do not contaminate water by cleaning of equipment or
disposal of wastes. Do not reuse empty bag. Dispose of
through regular refuse collection system.

ANTIDOTE—EXTERNAL—In case of spillage on skin, wash
promptly with soap and water.

INTERNAL—Induce vomiting promptly. CALL A PHYSICIAN

The manufacturer makes no warranties, express or implied,
concerning this product or its use, which extend beyond the
description on the face hereof. All statements concerning this
product apply only when used as directed.

Manufactured for GREENWAY PRODUCTS

**By DOLD FEED CO., INC.
BUFFALO, N. Y. 14210**

U.S. PAT. NO. 3,281,060

E. P. 6-74

TREATS & FEEDS 5,000 SQ. FT. OF LAWN AREA

795

NET WT. 20 LBS

SUPERGREEN

WEED

AND

FEED

PLUS



113453

SUPERGREEN WEED AND FEED PLUS

UNION

U



WEED AND FEED PLUS

SUPERGREEN WEED AND FEED PLUS

ACTIVE INGREDIENTS

2,4-Dichlorophenoxyacetic Acid 1.28%

INERT INGREDIENTS 98.72%

(RELATIVE TO HERBICIDAL CONTENT)

100% Wettable Acidity Equivalent to approximately 250 lbs. Calcium Carbonate Per Ton

GUARANTEED ANALYSIS

TOTAL NITROGEN (N) 10.0%

Available Phosphoric Acid (P₂O₅) 4.0%

Soluble Potash (K₂O) 6.0%

Iron (Fe) 0.7%

CAUTION KEEP OUT OF REACH OF CHILDREN

SEE BACK PANEL FOR ADDITIONAL CAUTIONS

EPA ESTAB. REG. # 6665-NY-01

EPA PROD. # 11345-3

