

Reg # 11345-3

PM-23

Kilo

196

United States Environmental Protection Agency
Washington, D.C. 20460

Office of
Prevention, Pesticides and
Toxic Substances

OCT 15 1993

John L. White
DOLD FEED COMPANY INC
717 ELK STREET
BUFFALO NY 14210


Subject: Label Amendment Submission of 06/21/93
in Compliance with the PR Notice 93-7
EPA Reg Nr. 11345-3
SUPERGREEN WEED AND FEED PLUS

Dear Registrant:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, REMOVES THE PRODUCT FROM THE SCOPE OF THE REQUIREMENTS CONTAINED IN 40 CFR PART 156, SUBPART K. Five copies of the finished labeling must be submitted prior to releasing the product for shipment.

The Agency is accepting only those changes to your labeling submitted in connection with your amendment application dated 06/21/93 which are necessary to comply with the requirements of 40 CFR part 156, subpart K. Any proposed changes to the labeling submitted in connection with this amendment application that were not directly related to compliance with the requirements of PR Notice 93-7 were not considered and have not been accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing such changes. If your product is currently suspended, the acceptance of this labeling amendment does not affect in any way the suspension.

Sincerely,



Jim Tompkins, Deputy Chief
Registration Support Branch

DOLD FEED COMPANY INC
SUPERGREEN WEED AND FEED PLUS
11345-3 06/21/93
Original Submission

29 6

The exclusionary statement must conform to the EPA-approved language specified in PRM 93-11.

396

(FRONT PANEL)

8/93

FEEDS 5,000 SQ. FT. OF LAWN AREA

SUPER GREEN

NET WT. 17 LBS. (7.8 kg)

**WEED
and
FEED
13-3-3**

ACCEPTED
with COMMENTS
in EPA Letter Dated

OCT 15 1993

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
11345-3

GUARANTEED ANALYSIS

TOTAL NITROGEN (N).....13.0%
1.0% AMMONIACAL NITROGEN
12.0% UREA NITROGEN
AVAILABLE PHOSPHORIC ACID (P₂O₅) 3.0%
SOLUBLE POT ASH (K₂O)..... 3.0%

ACTIVE INGREDIENTS

2,4-DICHLOROPHOXYACETIC ACID..... 1.26%
INERT INGREDIENTS..... 98.74%
TOTAL 100.00%

PRIMARY PLANT FOOD DERIVED FROM UREA, DIAMMONIUM PHOSPHATE AND MURIATE OF POTASH.

POTENTIAL ACIDITY EQUIVALENT TO APPROXIMATELY 250 LBS. CALCIUM CARBONATE PER TON

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANELS FOR ADDITIONAL PRECAUTIONARY STATEMENTS
AND STATEMENT OF PRACTICAL TREATMENT

EPA REG. # 11345-3

EPA ESTAB.. - 11345-NY-01

BEST AVAILABLE COPY

496

(REAR PANEL)

**Feeds 5,000 sq. ft.
(50' x 100')**

**Net Wt 17 lbs.
(7.8 kg)**

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Avoid contact with skin, eyes or clothing. Avoid breathing dust. Wash thoroughly after handling. Causes moderate eye injury.

When using this product, wear long sleeved shirt, long pants, socks and shoes. After using this product, remove clothing and launder separately before reuse and promptly and thoroughly wash hands and exposed skin with soap and water. The maximum number of Broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or Pets on treatment area. During application, Do Not enter treatment area until dust has settled.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED	GET MEDICAL ATTENTION.
IF ON SKIN	Wash Skin with Plenty of Soap and water.
IF IN EYES:	Immediately Flush Eyes with Water for 15 minutes. If Irritation develops, Get Medical Attention.

ENVIRONMENTAL HAZARD

This product is toxic to Aquatic invertebrates. Drift or runoff may adversely effect Aquatic invertebrates and non target plants. DO NOT Apply directly to water or to areas where surface water is present or to intertidal areas below the mean high watermark. DO NOT contaminate water when disposing of equipment washwaters.

BEST AVAILABLE COPY

596

(Rear Panel Cont)

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read entire label before using this product. Use strictly in accordance with label precautionary statement and directions.

It is for use on Home Lawns, Turf, (Excluding Pastures, Sod Farms, and turf grown to produce seed) and Golf Fairways, aprons, Tees and Roughs, Parks, Cemeteries and Similar non-crop areas. This product may be used on Blue Grass, Fescue, Rye, Bermuda, Bahia, Zoysia, Centipede and St. Augustine grass. DO NOT USE on Dichondra and Bentgrass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to promote Lush green growth and control many broadleaf weeds in lawns. Some of these weeds are:

Black Medic, Buckhorn, Burdock, Chicory, Chickweed, Clover, Dock, Dandelion, English Daisy, Ground Ivy, Healall, Lamb's -quarter, Mallow, Morning Glory, Pigweed, Plantain, Poison Ivy, Poison Oak, Pursland, Ragweed, Red Clover, Red Sorrel, Stitchwort, Speedwell, Spurge, White Clover, Wild Carrot, Wild Garlic, Wild Onion, Wild Lettuce, Yarrow and other broadleaf weeds.

WHEN TO APPLY; Best results are obtained when weeds are actively growing in Spring or early Fall.

IMPORTANT: Apply when grass is moist (after a rain, sprinkling or dew). DO NOT water lawn after material is applied. For most effective broadleaf weed killing action, avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material. Avoid contact with valuable shrubs, plants or flowers. Do not over-apply. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded lawn until after 3 mowings

NOTE: When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, overapplication may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of the grass.

(Rear panel contd)

BEST AVAILABLE COPY

696

(Rear panel contd)

HOW TO APPLY: This 17 pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, crisscrossing the area at half the rate each time is also acceptable. Lawns that are heavily overrun by weeds and that are in poor condition may require more than one application of the product. A suggested second application should be made in approximately 30 days.

The following settings are approximate for applying contents of this package at a suggested rate of 3.4 pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 sq. ft. and adjust to apply at the rate of .85 lb. per 250 sq. ft. Clean spreader thoroughly after use.

SPREADER SETTINGS - On Standard Models - Set all makes at about 1/5 of total opening.

- Central, Greenfield, Cross Country6
- Scotts-Standard Models Only 5

These settings are furnished as a guide ONLY - Age, condition of spreader, speed of operation and roughness of terrain can cause variations in rates applied.

Sweep all fertilizer materials from concrete surface immediately after application. Do not wash material from concrete surfaces because fertilizers which contain iron may cause discoloration upon oxidation.

STORAGE: Always store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near foodstuffs or feed. To prevent cross-contamination, do not store near other fertilizers or near herbicides, insecticides or fungicides.

CONTAINER DISPOSAL: Do not reuse empty bag.. Fold empty bag and discard in trash.

The Manufacturer and Seller makes no warranties, express or implied, concerning this product or its use which extend beyond the description on the face hereof. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

MANUFACTURED FOR SUPERGREEN PRODUCTS

BY DOLD FEED CO., INC. BUFFALO, N.Y. 14210

(ERP,WPS,WET) (0802393 JLW)

BEST AVAILABLE COPY