

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

NOV 29 1993

Mr. John L. White
Gro Green Products, Inc.
717 Elk Street
Buffalo, New York 14210

Dear Mr. White:

SUBJECT: Super Green Weed and Feed
EPA Registration No. 11345-3
Label Amendment
Your Resubmission Dated October 28, 1993

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended is acceptable with the following provisions:

1. Under the Caution Statement, correct the typographical errors located in the last two sentences of the paragraph.
2. Under the Environmental Hazards Section, change the third sentence to read: "Do not apply directly to water."
3. In the Directions for Use, under the Container Disposal Statement, please change the sentence to read as follows:
"Do not reuse empty bag. Securely wrap original container in several layers of newspaper and discard in trash."

A stamped copy is enclosed for your records. Please submit five (5) final printed copies for the referenced label, incorporating the above changes, before releasing the product for shipment.

Sincerely yours,

Joanne I. Miller
Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (7505C)

CONCURRENCES

SYMBOL	7505C						
SURNAME	J. Miller						
DATE	11/24/93						

(FRONT PANEL)

16# 10/93

FEEDS 5,000 SQ FT OF LAWN AREA

SUPER GREEN

NET WT 16 LBS. (7.25 kg)

**WEED
and
FEED
13-3-3**

GUARANTEED ANALYSIS

TOTAL NITROGEN (N)	13.0%
1.0% AMMONIACAL NITROGEN	
12.0% UREA NITROGEN	
AVAILABLE PHOSPHORIC ACID (P ₂ O ₅)	3.0%
SOLUBLE POTASH (K ₂ O)	3.0%

ACTIVE INGREDIENTS

2,4-DICHLOROPHOXYACETIC ACID	1.26%
INERT INGREDIENTS	98.74%
TOTAL	100.00%

PRIMARY PLANT FOOD DERIVED FROM UREA, DIAMMONIUM PHOSPHATE AND MURIATE OF POTASH.

POTENTIAL ACIDITY EQUIVALENT TO APPROXIMATELY 250 LBS. CALCIUM CARBONATE PER TON

KEEP OUT OF REACH OF CHILDREN


CAUTION

**SEE BACK PANELS FOR ADDITIONAL PRECAUTIONARY STATEMENTS
AND STATEMENT OF PRACTICAL TREATMENT**

EPA REG. # 11345-3

EPA ESTAB. # 11345-NY-01

~~REGISTERED~~
 2,4-DICHLOROPHOXYACETIC ACID
 50% SOLUTION
 29 NOV 1993



11345-3

(REAR PANEL)

#2

Feeds 5,000 sq. ft
(50' x 100')

Net Wt 16 lbs.
(7.25 kg)

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION


Avoid contact with skin, eyes or clothing. Avoid breathing dust. Wash thoroughly after handling. Causes moderate eye injury. When using this product, wear long sleeved shirt, long pants, socks and shoes. After using this product, remove clothing and launder separately before reuse and promptly and thoroughly wash hands and exposed skin with soap and water. The maximum number of Broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or Pets on treatment area. During application, Do Not enter treatment area until dust has settled.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED GET MEDICAL ATTENTION.
IF ON SKIN Wash Skin with Plenty of Soap and water.
IF IN EYES: Immediately Flush Eyes with Water for 15 minutes. If Irritation develops, Get Medical Attention.

ENVIRONMENTAL HAZARD

This product is toxic to Aquatic invertebrates. Drift or runoff may adversely effect Aquatic invertebrates and non target plants. DO NOT Apply directly to water or to areas where surface water is present or to intertidal areas below the mean high watermark. DO NOT contaminate water when disposing of equipment washwaters.



BEST AVAILABLE COPY

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read entire label before using this product. Use strictly in accordance with label precautionary statement and directions.

It is for use on Home Lawns, Turf, (Excluding Pastures, Sod Farms). This product may be used on Blue Grass, Fescue, Rye, Bermuda, Bahia, Zoysia, Centipede and St. Augustine grass. DO NOT USE on Dichondra and Bentgrass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.


This product is designed to promote Lush green growth and control many broadleaf weeds in lawns. Some of these weeds are:

- Black Medic, Buckhorn, Burdock, Chicory, Chickweed, Clover, Dock, Dandelion, English Daisy, Ground Ivy, Healall, Lamb's -quarter, Mallow, Morning Glory, Pigweed, Plantain, Poison Ivy, Poison Oak, Pursland, Ragweed, Red Clover, Red Sorrel, Stitchwort, Speedwell, Spurge, White Clover, Wild Carrot, Wild Garlic, Wild Onion, Wild Lettuce, Yarrow and other broadleaf weeds.

WHEN TO APPLY: Best results are obtained when weeds are actively growing in Spring or early Fall.

IMPORTANT: Apply when grass is moist (after a rain, sprinkling or dew) DO NOT water lawn after material is applied. For most effective broadleaf weed killing action, avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material. Avoid contact with valuable shrubs, plants or flowers. Do not over-apply. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded lawn until after 3 mowings

NOTE: When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, overapplication may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of the grass.



BEST AVAILABLE COPY

(Rear panel contd)

4

HOW TO APPLY: This 16 pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, crisscrossing the area at half the rate each time is also acceptable. Lawns that are heavily overrun by weeds and that are in poor condition may require more than one application of the product. A suggested second application should be made in approximately 30 days.

The following settings are approximate for applying contents of this package at a suggested rate of 3.2 pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 sq. ft. and adjust to apply at the rate of .08 lb per 250 sq. ft. Clean spreader thoroughly after use.

SPREADER SETTINGS - On Standard Models - Set all makes at about 1/5 of total opening.

- Central, Greenfield, Cross Country 6
- Scotts-Standard Models Only 5



These settings are furnished as a guide ONLY - Age, condition of spreader, speed of operation and roughness of terrain can cause variations in rates applied.

Sweep all fertilizer materials from concrete surface immediately after application. Do not wash material from concrete surfaces because fertilizers which contain iron may cause discoloration upon oxidation.

STORAGE: Always store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near foodstuffs or feed. To prevent cross-contamination, do not store near other fertilizers or near herbicides, insecticides or fungicides.

CONTAINER DISPOSAL: Do not reuse empty bag. Fold empty bag and discard in trash.

The Manufacturer and Seller makes no warranties, express or implied, concerning this product or its use which extend beyond the description on the face hereof. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.



6 of 6

MANUFACTURED FOR SUPERGREEN PRODUCTS
BY DOLD FEED CO., INC. BUFFALO, N.Y. 14210

#5

(ERP,WPS,WET,16#)DK (10/93 JW)

95
50