

USE DIRECTIONS FOR SAN-I-TIZE

San-I-Tize is a powerful disinfectant and sanitizer. It is effective against a wide range of bacteria, viruses, and fungi. It is safe for use on most hard surfaces and is non-toxic to humans and animals when used as directed.

For general disinfection and sanitization, use a solution of 1 oz. San-I-Tize in 4 gallons of water.

Apply to sink tops, countertops, refrigerated storage and display equipment and other stationary hard surfaces by cloth or brush.

Dishes, silverware, glasses, cooking utensils and other similar size food processing equipment can be sanitized by immersion in a 1 oz. 4 gallon dilution of San-I-Tize.

For Potable water rinse is required.

A 1 oz. 4 gallons San-I-Tize has a hard water tolerance of 500FPM of hardness calculated as CaCO₃ when evaluated by the AOAC Germicidal and Detergent Sanitizer Method against Escherichia coli and Staphylococcus aureus.

The udders, flanks, and teats of dairy cows can be sanitized by washing with a solution of 1 oz. San-I-Tize in 4 gallons of warm water.

Use a fresh towel for each cow. Avoid contamination of sanitizing solution by dirt and soil. Do not dip used towel back into sanitizing solution. When solution becomes visibly dirty, discard and provide fresh solution.

DANGER:

Keep Out of Reach of Children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

FIRST AID:

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution; or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

Note To Physician: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

SAN-I-TIZE

DISINFECTANT-SANITIZER-DEGREASER
 Effective Against Bacteria, Viruses, and Fungi
 For Industrial, Institutional, Veterinary and Other Commercial and Home Use

BLUE LINE

GAC JANITORIAL SUPPLIES, INC.

AOAC PHENOL COEFFICIENTS

Staphylococcus aureus, ATCC #6538	79.2
Salmonella typhosa, ATC #6539	78.1

ACTIVE INGREDIENTS

Alkyl (C ₁₂ , 50%; C ₁₀ , 40%; C ₈ , 10%)	
Dimethyl Benzyl Ammonium Chloride	10.0%
Ethanol	2.0%
	12.0%
Inert Ingredients Water	88.0%
	100.0%

EPA Registration No. 11371-1

DANGER:

KEEP OUT OF REACH OF CHILDREN. SEE FIRST AID AND OTHER PRECAUTIONS ON LEFT PANEL.

2501 Commercial St.,
 San Diego, CA 92113
 233-7318

ACCEPTED

SAN-I-TIZE

DISINFECTANT, SANITIZER, DEODORIZER

Commercial Division, The University of California, Institute
of Industrial Hygiene, University of California, and Home Products

BLUE LINE

GAC JANITORIAL SUPPLIES, INC.

AOAC PHENOL COEFFICIENTS

Staphylococcus aureus, ATCC #6538	79.2
Salmonella typhosa, ATC #6539	78.1

ACTIVE INGREDIENTS

Alkyl (C ₁₂ , 50%; C ₁₀ , 40%; C ₈ , 10%)	
Dimethyl Benzyl Ammonium Chloride	10.0%
Ethanol	2.0%
	12.0%
Inert Ingredients Water	88.0%
	100.0%

EPA Registration No. 11371-1

DANGER:

KEEP OUT OF REACH OF CHILDREN. SEE
FIRST AID AND OTHER PRECAUTIONS ON
LEFT PANEL.

2501 Commercial St.,
San Diego, CA 92113
233-7313

San-I-Tize is a powerful disinfectant, sanitizer, and deodorizer.

It is effective against a wide range of bacteria, viruses, and fungi. It is also effective against odors and is safe for use on most hard surfaces.

It is recommended for use in hospitals, schools, homes, and other public buildings.

For disinfection of floors, walls, bed-rines, countertops, tables, chairs, garden paths, bathroom fixtures and other hard surfaces.

Add 3 oz. of San-I-Tize to 5.5 gallons of water. Apply to previously cleaned hard surface with mop or cloth.

At 3 oz. per gallon use level, San-I-Tize is effective against Staphylococcus aureus and Salmonella choleraesuis.

Disinfection of Barber Tools

Barber tools (such as combs, brushes, razors, and scissors) can be disinfected by immersing in a 1/2 oz. per gallon solution of San-I-Tize.

Disinfection of Poultry Equipment, Animal Quarters and Kennels.

Poultry brooders, watering fountains, feeding equipment and other animal quarters (such as stalls and kennel areas) can be disinfected after thorough cleaning by applying a solution of 4 1/4 oz. San-I-Tize to 4 gallons of water with a mop, cloth or brush. Small utensils should be immersed in this solution.

Prior to disinfection, all poultry, other animals and their feeds must be removed from the premises. This includes emptying all troughs, racks and other feeding and watering appliances. Remove all litter and droppings from floors, walls and other surfaces occupied or traversed by poultry or other animals.

After disinfection, ventilate buildings, coops and other closed spaces. Do not house poultry, or other animals, or employ equipment until treatment has been absorbed, set or dried.

All treated equipment that will contact feed or drinking water must be rinsed with potable water before reuse.

San-I-Tize can be removed thoroughly with water and dried if necessary.

1 GALLON = 128 FL. OZ.

ACCEPTED WITH COMMENTS

ACCEPTED
21 1965

...of the...
...of the...
...of the...

...of the...
...of the...
...of the...

...of the...
...of the...
...of the...

DANGER:

Causes severe eye and skin damage. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when mixing. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID:

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution; or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

Physician: For more information, contact your local health department or the manufacturer. For more information, contact your local health department or the manufacturer.

JEROME

...of the...
...of the...
...of the...

BLUE LINE

GAC JANITORIAL SUPPLIES, INC.

AGAC PHENOL COEFFICIENTS

Staph. aureus (ATCC No. 6538)	70
Salmonella typhosa (ATCC No. 6539)	50

ACTIVE INGREDIENTS

Octyl Decyl Dimethyl Ammonium Chloride	1.50%
Dioctyl Dimethyl Ammonium Chloride	2.70%
Didecyl Dimethyl Ammonium Chloride	2.25%
Tetrasodium Ethylenediamine Tetraacetate	2.10%
Isopropyl Alcohol	2.70%

INERT INGREDIENTS

63.00%
100.00%

EPA Registration No. 115-17

DANGER

...of the...
...of the...
...of the...

2501 Commercial St.,
San Diego, CA 92113
233-7318

ACCEPTED
AUG 27 1975
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND ROBERTICIDE ACT