

LANE'S PRE-EMERGENCE HERBICIDE

**READ ALL DIRECTIONS CAREFULLY BEFORE APPLYING.
WEEDS CONTROLLED**

Annual Grasses

Large Crabgrass	Crowfootgrass	Sandbur (sandspur)
Smooth Crabgrass	Barnyardgrass (watergrass)	Stinkgrass
Yellow Foxtail	Johnsongrass (from seed)	Annual Bluegrass
Giant Foxtail	Witchgrass (ticklegrass)	(poa annua)
Goosegrass	Fall Panicum (panic grass)	Ryegrass
		Cheat

Annual Broadleaf Weeds

Pigweed (carelessweed)
Spry Amaranth
(stickerweed)
Lambsquarters
Smartweed

Purslane
Common Chickweed
Carpetweed
Knotweed

Winter Germinating Weeds

Red Sorrel
(from seed)
Pepperweed
Shepherd's Purse
Common Chick-
weed

GENERAL DIRECTIONS

Lane's Pre-Emergence Herbicide is a selective, pre-emergence herbicide which provides long-lasting weed control. Pre-Emergence Herbicide controls weeds by killing their seeds as they germinate. It does not control weeds once they have emerged. Pre-Emergence Herbicide is recommended for use on a wide variety of ornamental plants as well as established dichondra and ice plant. The ornamental species on which Pre-Emergence Herbicide can be used at the recommended rates include those listed on the side panels.

APPLICATION DIRECTIONS

Use a granular applicator which will uniformly apply Pre-Emergence Herbicide granules at the recommended rate. Both drop-type and rotary-type spreaders may be used. Calibrate the applicator prior to use and check frequently during application to be sure the equipment is working properly and distributing the granules uniformly. Apply Pre-Emergence Herbicide granules to a smooth soil surface to assure a uniform distribution.

Rainfall or overhead irrigation following Pre-Emergence Herbicide application will improve its weed control activity. Excessive moisture may leach Pre-Emergence Herbicide from the weed seed germination zone. Pre-Emergence Herbicide does not need to be mixed into the soil in order to be effective. Shallow cultivation will not reduce the weed control activity of Pre-Emergence Herbicide.

In areas of furrow irrigation incorporate Pre-Emergence Herbicide to a depth of 2 to 4 inches with P.T.O. driven equipment (tillers, cultivators, hoes), or double discs. Use only discs with positive depth control; incorporation of less than 2 inches or more than 4 inches may result in erratic weed control.


USDA Reg. No. 1140 1

A Selective Pre-Emergence Herbicide for the Control of Annual Grasses and Broadleaf Weed in Established Flowers and Dichondra, Ice Plant, Nursery Stock, Roses, Ornamental Trees and Wood Shrubs.

Active Ingredients

Diphenamid (N, N-Dimethyl-1-2, 2 diphenylacetamide) 5%
Inert Ingredients 95%

Caution: KEEP OUT OF REACH OF CHILDREN.



NET CONTENTS: [REDACTED] lbs.

ACCEPTED
Sep 27 - 1971

FOR [REDACTED]
USE [REDACTED]

ACCEP

NURSERY STOCK, ORNAMENTALS AND ICE PLANT

Apply Pre-Imergence Herbicide anytime throughout the year, prior to weed seed germination, for weed control. Pre-Imergence Herbicide may be applied overtop of plants. On large plants the granules should be directed to the soil beneath the plants. Do not apply to wet foliage.

Broadcast Application Rates

Light soil	Medium Soil	Heavy Soil
80 lbs. per acre or 2 lbs. per 1,100 sq. ft. (4 lbs. active)	100 lbs. per acre or 2½ lbs. per 1,100 sq. ft. (5 lbs. active)	120 lbs. per acre or 3 lbs. per 1,100 sq. ft. (6 lbs. active)

The above rates are for a broadcast (overall) application. For a band treatment, use the following formula to figure rates:

$\frac{\text{band width in inches}}{\text{row width in inches}} \times \text{recommended broadcast rate} = \text{amount to apply per acre on band}$

DICHONDRA & ICE PLANTS

Apply Pre-Imergence Herbicide overtop of established dichondra prior to weed seed germination. Overhead irrigation, following Pre-Imergence Herbicide application, is recommended for most consistent weed control results. To insure continuous weed control, repeat the application every six months.

Application Rates

All Soils

200 lbs. per acre or 5 lbs. per 1,100 sq. ft. (10 lbs. active)

To convert lawns of dichondra and turtgrass mixtures to pure dichondra, irrigate the lawn thoroughly the day before application. Apply the recommended rate of Pre-Imergence Herbicide the following day and irrigate. The established grass turf will discolor and die slowly within a period of 3 to 4 weeks.

PRECAUTIONS: Do not use Pre-Imergence Herbicide in direct combination with fertilizers, insecticides, or fungicides. Do not graze livestock on treated areas.

CAUTION: Harmful if swallowed. Avoid prolonged breathing of dust or repeated contact with skin. Wash thoroughly after using. Avoid contamination of feed or foodstuffs.

The manufacturer makes no warranties, express or implied, concerning this product or its use, which extend beyond the description on the label. All statements made concerning this product apply only when used as directed.

Destroy empty container. Do not reuse

ORNAMENTAL WOODY SHRUBS

<i>Abutilon</i>	<i>Abutilon</i>
<i>Adiantum</i>	<i>Adiantum</i>
<i>Azalea</i>	<i>Azalea</i>
<i>Begonia</i>	<i>Begonia</i>
<i>Camellia</i>	<i>Camellia</i>
<i>Clematis</i>	<i>Clematis</i>
<i>Cydonia</i>	<i>Cydonia</i>
<i>Forsythia</i>	<i>Forsythia</i>
<i>Gardenia</i>	<i>Gardenia</i>
<i>Hibiscus</i>	<i>Hibiscus</i>
<i>Hydrangea</i>	<i>Hydrangea</i>
<i>Jasminum</i>	<i>Jasminum</i>
<i>Philadelphus</i>	<i>Philadelphus</i>
<i>Rhododendron</i>	<i>Rhododendron</i>
<i>Syringa</i>	<i>Syringa</i>
<i>Taxus</i>	<i>Taxus</i>
<i>Yucca</i>	<i>Yucca</i>
<i>... ..</i>	<i>... ..</i>

<i>Abutilon</i>	<i>Abutilon</i>
<i>Adiantum</i>	<i>Adiantum</i>
<i>Azalea</i>	<i>Azalea</i>
<i>Begonia</i>	<i>Begonia</i>
<i>Camellia</i>	<i>Camellia</i>
<i>Clematis</i>	<i>Clematis</i>
<i>Cydonia</i>	<i>Cydonia</i>
<i>Forsythia</i>	<i>Forsythia</i>
<i>Gardenia</i>	<i>Gardenia</i>
<i>Hibiscus</i>	<i>Hibiscus</i>
<i>Hydrangea</i>	<i>Hydrangea</i>
<i>Jasminum</i>	<i>Jasminum</i>
<i>Philadelphus</i>	<i>Philadelphus</i>
<i>Rhododendron</i>	<i>Rhododendron</i>
<i>Syringa</i>	<i>Syringa</i>
<i>Taxus</i>	<i>Taxus</i>
<i>Yucca</i>	<i>Yucca</i>

ORNAMENTAL TREES

<i>Abutilon</i>	<i>Abutilon</i>
<i>Adiantum</i>	<i>Adiantum</i>
<i>Azalea</i>	<i>Azalea</i>
<i>Begonia</i>	<i>Begonia</i>
<i>Camellia</i>	<i>Camellia</i>
<i>Clematis</i>	<i>Clematis</i>
<i>Cydonia</i>	<i>Cydonia</i>
<i>Forsythia</i>	<i>Forsythia</i>
<i>Gardenia</i>	<i>Gardenia</i>
<i>Hibiscus</i>	<i>Hibiscus</i>
<i>Hydrangea</i>	<i>Hydrangea</i>
<i>Jasminum</i>	<i>Jasminum</i>
<i>Philadelphus</i>	<i>Philadelphus</i>
<i>Rhododendron</i>	<i>Rhododendron</i>
<i>Syringa</i>	<i>Syringa</i>
<i>Taxus</i>	<i>Taxus</i>
<i>Yucca</i>	<i>Yucca</i>

ROSE ESTABL

<i>Abutilon</i>	<i>Abutilon</i>
<i>Adiantum</i>	<i>Adiantum</i>
<i>Azalea</i>	<i>Azalea</i>
<i>Begonia</i>	<i>Begonia</i>
<i>Camellia</i>	<i>Camellia</i>
<i>Clematis</i>	<i>Clematis</i>
<i>Cydonia</i>	<i>Cydonia</i>
<i>Forsythia</i>	<i>Forsythia</i>
<i>Gardenia</i>	<i>Gardenia</i>
<i>Hibiscus</i>	<i>Hibiscus</i>
<i>Hydrangea</i>	<i>Hydrangea</i>
<i>Jasminum</i>	<i>Jasminum</i>
<i>Philadelphus</i>	<i>Philadelphus</i>
<i>Rhododendron</i>	<i>Rhododendron</i>
<i>Syringa</i>	<i>Syringa</i>
<i>Taxus</i>	<i>Taxus</i>
<i>Yucca</i>	<i>Yucca</i>