

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
AND DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed. Avoid contact with skin or clothing. In case of contact, immediately flush skin and eyes with plenty of water. Get medical attention if irritation persists. Remove pets and cover fish aquariums before spraying.

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: - Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions give herewith.

SUN-BUGGER # 4

Insecticide Spray With S.B.P. 1382
For Bakeries, Food Plants, Restaurants, Schools,
Hotels, Motels, Office Buildings, Hospitals and Food
Handling Plants

ACTIVE INGREDIENTS:	% By Weight
*Resmethrin	0.050%
Related compounds	0.007%
INERT INGREDIENTS	99.943%
	TOTAL 100.00%

*Cis/trans isomers ratio: max. 30%(+)cis and min. 70%(+)trans
+ U.S. Patent No. 3,465,007 and 3,542,928

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Side Panel for Additional Precautionary Statements

EPA Reg. No. 11474-9

EPA Est. No. 11474CA01

MANUFACTURED BY:
SUNGRO CHEMICALS, INC.
P.O. BOX 24632 Los Angeles, CA. 90024

DIRECTIONS FOR USE.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Use any type of hand sprayer which produces a cloud of fog like mist. Apply at a rate of 5 fluid ounces per 1000 cubic feet of surface.

For flies, Mosquitoes, and Gnats: Close doors and windows. Spray freely toward the ceiling in all directions until room is filled with mist, then leave the room. Keep doors closed for at least 15 minutes. Ventilate the room when treatment is completed.

Wasps: Application should be made in the late evenings when the insects are at rest. Spray liberally into hiding and breeding places, contacting as many insects as possible. Repeat as necessary.

Avoid contamination of feed and foodstuffs. Use in food processing plants only when plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered or thoroughly cleaned before using. In hospitals, patients should be removed from room prior to treatment. Room should be ventilated for 2 hours after spraying. Do not return patients to room until after ventilation.

NET CONTENTS _____ GALLONS

011690

Reviewed. Registrant claims to be in accordance with registration of Pesticide Product Guidance for Phase 2 Reference, Page 2.12, Active to Inert Change in Status.