

DIRECTIONS FOR USE

Shake well before each use. Remove protective cap. Aim spray opening away from people. Push button to spray. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using the product in these areas, apply only when the facility is not in operation.

Domestic **INDOORS — FOR HOUSEFLIES, MOSQUITOES, AND GNATS:**

Close all doors and windows. Point nozzle upward, direct the spray mist to all parts of the room, especially windows and other light sources which attract these insects. Fill the room with mist, then leave the treated area. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed.

Domestic **INDOORS — FOR CRAWLING INSECTS:**

Roaches, earwigs, grain mites, sowbugs, spiders and centipedes: Spray thoroughly into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards, and storage areas. Spray directly on insects where possible.

Ants: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

Fleas: Protect dogs and cats from fleas by spraying their resting areas. Hold spray nozzle about 18 inches from area to be treated. Apply spray at the rate of two seconds per linear foot. Direct spray on resting areas and floor around these areas. Treat bedding the same way. Also spray entire inside surface of dog houses. Repeat as necessary.

Also treat dogs and cats with a registered product before they reenter the area just treated with this product.

Domestic **OUTDOORS — TO KILL FLYING INSECTS:**

Spray areas infested with mosquitoes, gnats (midges), houseflies, biting flies and flying moths (millers). Follow precautions listed above. Hold container 18 to 24 inches from shrubs, bushes and grassy areas. Spray these areas with a slow sweeping motion at the rate of from 1 to 3 seconds per cubic yard of space while moving away from the treated areas. To reduce annoyance from these flying insects, treat open porches and patios in a similar manner. Spray only when the air is still. Allow spray to

SUN-BUGGER#1

Active Ingredients:

*+(5-Benzyl-3-furyl) methyl 2, 2-dimethyl-3-(2-Methylpropenyl) Cyclopropanecarboxylate	0.250% w/w
Related compounds	0.034%
Aromatic petroleum hydrocarbons	0.332%
Inert Ingredients:	99.384%
	100.000%

*Cis/trans isomers ratio: max. 30% (+)cis and min. 70% (+)trans.

+U.S. Patent No. 3,465, 007

SHAKE WELL BEFORE USING

CAUTION

KEEP OUT OF REACH
OF CHILDREN.

Harmful if swallowed. Avoid contamination of feed and foodstuffs. Do not use on edible crop. Remove pets, birds and cover fish aquariums before spraying.

WARNING:

Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting. Never throw container into fire or incinerator.

Manufactured by:

SUNGRO CHEMICAL, INC.

P.O. Box 24632

Los Angeles, California 90024

E.P.A. Reg. No. 11474-11

Net contents _____ gals.

dissipate before occupying the treated area. Repeat as necessary.

Spray peonies, iris, zinnia, snapdragons, ageratums, chrysanthemums, nasturtiums, daisy, geraniums, marigolds, daphne, fatshedera, ivy, tobira, queen sago and broad leaf acacia for control of aphids, exposed thrips, adult white flies, leafhoppers, plant bugs, spittlebugs and Japanese beetles. Spray as directed above.

Ornamental Quince, Juniper, Cypress, Ornamental Mulberry, Ash, Cottonaster, Rose, Camellia, Azalea, Pyracantha and Caenothus may also be treated to control the above pests, but use with extreme caution. Certain varieties have shown some leaf burn under certain conditions of use such as heavy dosage. Repeat as necessary to obtain control.

(INDOORS) — TO PROTECT HOUSE PLANTS:

Spray infested African violets, philodendron, geraniums, begonias, coleus for control of aphids, exposed thrips and adult white flies. Repeat as necessary.

OUTDOORS — TO CONTROL NUISANCE PESTS:

Ants: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

Spiders: Direct short burst of spray on spiders and repeat as necessary.

Wasps: Application should be made in late evening when insects are at rest. Spray liberally into hiding and breeding places contacting as many insects as possible. Repeat as necessary.

OUTDOORS — TO PROTECT ORNAMENTAL PLANTS:

To control aphids, exposed thrip, adult white flies, leafhoppers, plant bugs, spittlebugs and Japanese beetle on small rhodendrons, pines, spruce, hemlock, spirea, dogwood, palm, mimosa, toyan, viburnum euonymus, oaks, maples, locust, forsythia and althea; and mimosa webworm larvae on mimosa, toyan, viburnum euonymus, oaks, maples, locust, forsythia and althea; and mimosa webworm larvae on mimosa, spray directly on insects. Hold container 18 inches from the infested plants. Spray in short bursts of one to three seconds of spray at the rate of one linear foot of growth per second. Direct spray to all sides covering foliage until slightly moist. Avoid heavy wetting. Repeat as necessary to obtain control.

