

Food Processing Equipment

Clean and rinse equipment thoroughly, then apply sanitizing solution containing one ounce SANITIZ-IT to four gallons of water. No potable water rinse required.

Dairies

Clean and rinse equipment thoroughly, then apply sanitizing solution containing one ounce SANITIZ-IT to four gallons of water (195ppm). No potable water rinse required. Follow recommendations of Local Health Board.

Restaurant and Bar Rinse

Dishes, glassware, silverware cooking utensils, wash with soap or synthetic detergent, rinse thoroughly and immerse in a solution containing one ounce SANITIZ-IT to four gallons of water (195ppm). No terminal potable water rinse required.

RINSE EMPTY CONTAINER THOROUGHLY WITH WATER AND DISCARD.

Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and bury in a safe place.

Seller makes no warranty expressed or implied concerning the use of this product other than indicated on the label. Buyer assumes all risks of use, storage, and handling of this material not in strict accordance with directions given herein.

SANITIZ-IT

DISINFECTANT - SANITIZER - DEODORIZER

ACTIVE INGREDIENTS:

n Alkyl (60%C₁₄, 30%C₁₆, 5%C₁₂, 5%C₁₈)
dimethyl benzyl ammonium chlorides 5%
n Alkyl (68%C₁₂, 32%C₁₄) dimethyl ethyl-
benzyl ammonium chlorides 5%

INERT INGREDIENTS 90%
TOTAL 100%

WARNING: KEEP OUT OF REACH OF CHILDREN!

Causes severe eye irritation; causes skin irritation. Do not get in eyes, on skin or clothing. Harmful if swallowed. Avoid contamination of food.

FIRST AID: In cases of contact, immediately flush eyes or skin with water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution, or if not available, drink large quantities of water. Avoid alcohol.

E.P.A. NO. 11474 - 2

NET CONTENTS _____ GALLONS

Manufactured by:

SUNGRO CHEMICALS INCORPORATED
P.O. Box 24632 Los Angeles, California 90024

SANITIZ-IT is a concentrated bactericide solution useful as a disinfectant, sanitizer and deodorant. It should be diluted according to instructions prior to use.

DIRECTIONS FOR USE

Cold Disinfection


For disinfection of previously cleaned surgical instruments, barber tools and dental equipment, simply submerge instruments into solution containing 1½ oz. SANITIZ-IT per gallon of water for ten minutes.

Household

For sanitizing and disinfecting floors, walls and inanimate hard surfaces. For Schools, Homes, Locker Rooms, Garbage Pails, Sink Tops, Corridors, Classrooms, Offices, Shower Stalls and Toilet Seats. Apply solution with mop or cloth. Sanitize with 1 oz. SANITIZ-IT to 4 gallons of water. Disinfect with 3 ounces SANITIZ-IT to 5 gallons of water. Kills Staphylococcus aureus and Salmonella choleraesuis.

Hospitals

For floors, walls and other hard surfaces. In Nursing Homes, Institutions, as well as Sink Tops Garbage Pails, Telephones and Restrooms, use 3½ ounces SANITIZ-IT per 5 gallons of water. At this level SANITIZ-IT will be effective against Pseudomonas aeruginosa PRD-10.



PURGE, WATER TREATMENT MICROBIOCIDE will control algae and bacterial slimes found in recirculating cooling tower waters. PURGE, WATER TREATMENT MICROBIOCIDE helps clean and loosen slime debris from cooling system surfaces. When used in slug doses, no other microbiocide is required.

PURGE WATER TREATMENT MICROBIOCIDE is economical to use because it is concentrated. It should be handled with care.

DANGER:

Keep Out of Reach of Children. Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID:

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution; or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

PURGE WATER TREATMENT MICROBIOCIDE

Twin-Chain Quaternary
Ammonium Compound Concentrate

Water Treatment Microbiocide
for Building and Industrial
Cooling Towers

Active Ingredients

Didecyl dimethyl ammonium chloride	50%
Isopropyl alcohol	20%

Inert Ingredients

30%
100%

EPA Registration No.

11474-14

Net Weight

400 lbs.

Net Volume

55 gals.

SUNGRO CHEMICALS, INC
P.O. BOX 24632 LOS ANGELES, CAL. 90024

DANGER

KEEP OUT OF REACH OF CHILDREN. SEE
LEFT PANEL FOR ADDITIONAL PRECAU-
TIONARY STATEMENTS.

USE DIRECTIONS

To control algae and bacterial slimes, use PURGE WATER TREATMENT MICROBIOCIDE as directed. For best results, slug feed. The frequency of addition of microbiocide needed depends on many factors. To optimize your use of PURGE WATER TREATMENT MICROBIOCIDE, follow this procedure.

1. Initially use 6 fluid ounces per 1000 gallons of water to be treated (20 ppm active quaternary).

Should the above dosage not give satisfactory results, use 9 fluid ounces per 1000 gallons of water.

Repeat the initial dose every seven days or increase the frequency if needed.

2. When the above treatment level is successful, use 2 to 3 fluid ounces per 1000 gallons of water to maximize efficiency. Repeat weekly as needed.

Should slime develop again, go back to initial dosage.

Cooling tower waters that are inherently low in algae growth and bacteria count may be adequately controlled by the lower range of these dosages; slug fed every seven days.

Dilute the appropriate amount of PURGE WATER TREATMENT MICROBIOCIDE in 1 or 2 gallons of water then add to the sump of the tower. Note, this product weighs 7.49 lbs per gallon (at 20°C).

This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds, or public water.

Should tower be heavily fouled, a pre-cleaning is required.

Do not reuse empty drum. Return to drum reconditioner or rinse well with soap solution and discard. Do not contaminate water by disposal of waste.

5/72

ACCEPTED

FEB 20 1974

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 11474-14 SUBJECT
TO ATTACHED COMMENTS.