

BEST DOCUMENT AVAILABLE

00 3

#### CAUTIONARY STATEMENTS FOR HUMANS AND DOMESTIC ANIMALS

##### CAUTION

Get prompt medical aid. May be harmful if swallowed, or absorbed through the skin. Do not use on clothing. In case of contact, wash with soap and warm water. Wash hands, arms, face, and body with soap and water before eating or drinking. Wash contaminated clothing with soap and hot water.

Do not contact treated areas until surfaces are dry. Remove pets and cover fish bowls before spraying.

Get prompt medical aid.

Pine sulfate is antidotal.

#### ENVIRONMENTAL HAZARDS

Use spray. Provide adequate ventilation of area. Do not store near feed or food products. Do not use on feed, food, food processing equipment, or surfaces. Do not apply to animals.

**PACKING & FOOD PROCESSING PLANTS:** Use in product areas of meat packing plants, poultry processing plants, or other food processing plants. Any exposed food, or packaging materials must be removed before spraying. Rinse all equipment with water before spraying. Treat food processing plants early in the morning, when plant is not in operation. Docks and outside areas where insects are present in preventing re-infestation. Good sanitation of wastes is essential in any effective program.

#### FLAMMABLE OR CHEMICAL HAZARDS

Do not spill or store near heat or open flame.

#### USE

Follow federal law to use this product in a manner consistent with its labeling.

# SUNGRO RESIDUAL SPRAY

CONTAINS BAYGON®

#### For Industrial and Institutional Use

Residual Spray is a highly effective ready to use formulation for control of the insects listed below, including those resistant to certain chlorinated hydrocarbon and phosphate. This product gives rapid kill and long residual activity.

##### ACTIVE INGREDIENTS:

2-(1-Methylethoxy) phenol methylcarbamate*	1.0%
Petroleum Distillate	84.0%
INERT INGREDIENTS	15.0%

\*U.S. Patent No. 3,111,539

\*BAYGON is a Reg. TM of the Parent Company of Farbenfabriken BAYER GmbH, Leverkusen.

**KILLS:** Ants, Brown Dog Ticks, Cockroaches, Clover Mites, Crickets, Earwigs, Fleas, Flies, Millipedes, Mosquitos, Scorpions, Silverfish, Sawbugs and Spiders, Saw-toothed Grain Beetles, Wasps, and Hornets.

## CAUTION:

**KEEP OUT OF REACH OF CHILDREN**  
(SEE SIDE PANEL FOR ADDITIONAL CAUTIONS)


MANUFACTURED BY:

**SUNGRO CHEMICALS INCORPORATED**

P.O. BOX 24632 LOS ANGELES, CALIFORNIA 90024

E.P.A. Reg. No. 11474-20 E.P.A. Est. No. 11474CA01

NET CONTENTS:.....GAL.



Apply with a low-pressure sprayer equipped with a coarse spray nozzle. Spray surfaces only until wet. Avoid excessive run-off. Apply to all insect breeding sites and harborage. Also treat adjacent exposed surfaces where insects may crawl when they come out of hiding. Repeat treatment as necessary.

**Cockroaches:** Thoroughly treat the infested area. Spray baseboards, cracks and crevices; around window and door frames, the under surfaces of shelves, drawers and work tables behind and beneath cabinets, refrigerators, sinks and stoves, in and around waste containers, switch panels, junction boxes, floor drains and other places where cockroaches hide or nest.

**Ants:** Apply to trails and places where ants enter premises such as window and door sills. Spray hills and runways outdoors.

**Sawtoothed Grain Beetles (exposed stages):** Thoroughly treat floors and the walls of the infested area with attention to cracks, crevices, and corners and perimeters. Sanitation and dust control are of primary importance. Fumigate or destroy infested products.

**Silverfish:** Apply to localized areas infested by these insects.

**Clover Mites, Crickets, Earwigs, Millipedes, Sawbugs, Scorpions and Spiders:** Thoroughly spray the interior and exterior perimeter of the infested premises. For Clover Mites, give special attention to window sills.

**Fleas, Flies, Mosquitoes:** Spray infested areas thoroughly including outside surfaces of screens, doors, window frames, foundations, patios and other areas where insects may enter.

**Insect Control in Research Facilities:** The use of Residual Spray in and around research animal quarters for insect control will not induce changes in liver microsomal enzyme activity in these animals.

#### STORAGE AND DISPOSAL

**PROHIBITIONS:** Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

**PESTICIDE DISPOSAL:** Pesticide, spray, mixture, or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

**CONTAINER DISPOSAL:** (a) Containers of 30 gallons or more—reseal container and offer for reconditioning, or; (b) All containers triple rinse (or equivalent) and offer for recycling, reconditioning, or disposal in approved landfill, or bury in a safe place.

BEST 1.

...LABLE

71-4

71