# PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION.

In case of poisoning, get prompt medical aid. May be harmful if swallowed, inhaled, or absorbed through the skin. Do not get in eyes, on skin, or on clothing. In case of contact, wash immediately with soap and warm water. Wash hands, arms, and face thoroughly with soap and water before eating gesmoking. Wash all contaminated clothing with soap and lost water before re-use.

Do not allow children to contact treated areas until surfaces are thoroughly dry. Remove pets and cover fish bowls before spraying.

Should illness occur, get prompt medical aid.

TO PHYSICIAN: Atropine sulfate is antidotal.

#### ENVIRONMENTAL HAZARDS

Do not use as a space spray. Provide adequate ventilation of area being treated. Do not store near feed or food products. Avoid contamination of feed, food, food processing equipment, and food handling surfaces. Do not apply to animals.

FOR USE IN MEAT PACKING & FOUR PROCESSING PLANTS:
I've only in the inscible product areas of seat packing plants. Any exposed food products, requirement, or packaging materials out be resound our covered before spraying, wash all equipment outless with potable mater before resuming operations with potable mater before resuming operations before canning or other processing begins. Repeate applications, if necessary, when plant is not in operation. Spraying unloading docks and outside areas where insects congregate will aid in preventing reinfestation. Sood sanitation and proper disposal of mastes is essential in any effective insect control program.

#### PHYSICAL OR CHEMICAL HAZARDS

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.



## CONTAINS BAYGON®

### For Industrial and Institutional Use

Residual Spray is a highly effective ready to use formulation for control of the insects listed below, including those resistant to certain chlorinated hydrocarbon and phosphate. This product gives rapid kill and long residual activity.

ACTIVE INGREDIENTS:

3-(1-Methylethoxy) phenol methylcarbamate 1.0% Petroleum Distillate 84.0% INERT-INGREDIENTS 15.0%

BAYGON Is a Reg. TM of BAYER! AG. Leverkusen.

KILLS: Ants. Brown Dog Ticks, Cockrosches, Stever Miles, Crickets, Earwigs, Fleas, Flies, Millinedes, Meaguithe, Scorpions, Silverfish, Sawhugs and Spiders, Say toolhed Grain Beetles, Wasps, and Hornets.

## CAUTION

KEEP, OUT: OF: REACH(OF CHILDREN (SEE), SIDE: PANEL FOR ADDITIONAL CAUTIONS

SUNGRO CHEMICALS INCORPORATED
P.D. BOX 24632 LOS ANGELES CALIFORNIA 90024

NET CONTENTS GAL



It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Apply with L tow-p. es. urb scrayer on Lapad with a coarse spray nozzle. Sp. ay surfaces only until wet. Avoid excessive run-off. Apply to all insect breeding sites and harborages. Also treat adjacent exposed surfaces where insects may craw when they come out of hiding. Repeat treatment as necessary.

Cockroaches: Thoroughly treat the infested area. Spray baseboards, cracks and crevices; around window and door frames; the under surfaces of shelves, drawers and work tables behind and beneath cabinets, refrigerators, sinks and stoves; in and around waste containers, switch panels, junction boxes, floor drains and other places where cockroaches hide or nest.

Ants: Apply to trails and places where ants enter premises such as window and door sills. Spray hills and runways outdoors.

Sawtoothed Grain Beetles (exposed stages): Thoroughly treet floors and the walls of the intested area with attention to exacts, crevices, and comers and perimeters. Sanitation and agust control are of primary importance. Fumigate or destroy injested products.

Silverfish: Apply to localized areas infested by these insects.

Clover Mites, Crickets, Earwigs, Millipedes, Sowbugs, Scorpions and Spiders: Thoroughly spray the interior and exterior perimter of the infested premises, For Clover Mites, give special attention to window sills.

Fleas, Flies, Mosquitoes: Spray infested areas thoroughly including outside surfaces of screens, doors, window frames, foundations, patios and other areas where insects may enter.

Insect Control in Research Facilities: The use of Residual Spray in and around research animal quarters for insect control will not induce changes in liver microsomal enzyme activity in these animals.

## STORAGE & DISPOSAL

Do not contaminate water, 'food or feed by storage or disposal. '

PESTICIDE DISPOSAL:

Mastes resulting from the use of this product any be disposed of on site or at an approved maste disposal facility.

CONTRINER DISPOSAL:

Triple rinse for equivalentl. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfull, or by other procedures approved by state and local authorities.