

P. # 11474-20

PM-12

191

**Precautionary Statements**  
**Hazards to Humans and Domestic Animals**  
**Caution**

In case of poisoning, get prompt medical aid. May be harmful if swallowed, inhaled, or absorbed through the skin. Do not get in eyes, on skin, or on clothing. In case of contact, wash immediately with soap and warm water. Wash hands, arms, and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use.

Do not allow children to contact treated areas until surfaces are dry. Remove pets and cover fish bowls before spraying. Should illness occur, get prompt medical aid.

TO PHYSICIAN: Atropine sulfate is antidotal.

**ENVIRONMENTAL HAZARDS**

Do not use as a space spray. Provide adequate ventilation of area being treated. Do not store near feed or food products. Avoid contamination of feed, food, food processing equipment and food handling surfaces. Do not apply to animals.

**PHYSICAL OR CHEMICAL HAZARDS**

Do not use, pour, spill or store near heat or open flame.

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Apply with a low-pressure sprayer equipped with a coarse spray nozzle. Spray surfaces only until wet. Avoid excessive run-off. Apply to all insect breeding sites and harborages. Also treat adjacent exposed surfaces where insects may crawl when they come out of hiding. Repeat treatment as necessary.

**FOR USE IN MEAT PACKAGING & FOOD PROCESSING PLANTS:** Use only in the inedible areas of meat packaging plants. Any exposed food products, equipments or packaging materials must be removed or covered before spraying, wash all equipment with an effective cleaning compound followed by a rinse with potable water before resuming operations. Treat food processing plants early in the season before canning or other processing begins. Repeat application if necessary, when plant is not in operation. Spraying unloading docks and outside areas where insect congregate will aid in preventing reinfestation. Good sanitation and proper disposal of wastes is essential in any effective insect control program.

# RESIDUAL SPRAY

**CONTAINS PROPOXUR**  
**FOR INDUSTRIAL AND INSTITUTIONAL USE**

Residual Spray is a highly effective ready to use formulation for control of the insects listed below, including those resistant to certain chlorinated hydrocarbon and phosphate. This product gives rapid kill and long residual activity.

**KILLS:** Ants, Brown Dog Ticks, Cockroaches, Clover Mites, Crickets, Earwigs, Fleas, Flies, Millipedes, Mosquitoes, Scorpions, Silverfish, Sawbugs, and Spiders, Saw-toothed Grain Beetles, Wasps and Hornets.

Active Ingredients	by wt.
2-(1-methylethoxy)phenol methylcarbamate	1.00%
INERT INGREDIENTS	99.00%
TOTAL	100.00%

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
 (See Side Panel for Additional Cautions)

E.P.A. Reg. No. 11474-20 E.P.A. Est. No. 11474CA01

Manufactured by:  
**SUNGRO CHEMICALS, INC.**  
 P.O. BOX 24632  
 LOS ANGELES, CA. 90024

NET CONTENTS.....GAL

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Apply with a low-pressure sprayer equipped with a coarse spray nozzle. Spray surfaces only until wet. Avoid excessive run-off. Apply to all insect breeding sites and harborages. Also treat adjacent exposed surfaces where insects may crawl when they come out of hiding. Repeat treatment as necessary.

**Cockroaches:** Thoroughly treat the infested area. Spray baseboards, cracks and crevices around window and door frames; the under surfaces of shelves, drawers and work tables behind and beneath cabinets, refrigerators, sinks and stoves; in and around waste containers, switch panels, junction boxes, floor drains and other places where cockroaches hide or nest.

**Ants:** Apply to trails and places where ants enter premises such as window and door sills. Spray hills and runways outdoors.

**Sawtoothed Grain Beetles (exposed stages):** Thoroughly treat floors and walls of the infested area with attention to cracks, crevices, and corners and perimeters. Sanitation and dust control are of primary importance. Fumigate or destroy infested products.

**Silverfish:** Apply to localized areas infested by these insects.

**Clover Mites, Crickets, Earwigs, Millipedes, Sawbugs, Scorpions and Spiders:** Thoroughly spray the interior and exterior perimeter of the infested premises. For Clover Mites, give special attention to window sills.

**Fleas, Flies, Mosquitoes:** Spray infested areas thoroughly including outside surfaces of screens, doors, window frames, foundations, patios and other areas where insects may enter.

**Insect Control in Research Facilities:** The use of Residual Spray in and around research animal quarters for insect control will not induce changes in liver microsomal enzyme activity in these animals.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

011690

Reviewed. Registrant claims to be in accordance with  
 Registration of Pesticide Product Guidance for Phase 2  
 Rule, Page 2.12, Active to Inert Change in Status.