

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
AND DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed. Foods should be removed or covered during treatment. Any food processing surfaces should be covered during treatment or thoroughly cleaned before using. Do not apply while food is being prepared. Remove pets, birds, and cover fish aquariums before spraying.

ENVIRONMENTAL HAZARDS

This product may be toxic to fish. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not add directly to water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with it's labeling.

INDOOR APPLICATION

Vacate treated areas and ventilate before reoccupying.

For maximum effectiveness, a combination of spot surface treatment and fogging is recommended.

SURFACE SPRAYING: For rapid control of Roaches, Ants, Spiders, Crickets, Silverfish, Centipedes, Cheese Mites, Granary Weevils, Rice Weevils, Confused Flour Beetles, Rust Red Flour Beetles, Mites (clover, cheese, grain), Meal Worms, Grain Beetles, (confused flour beetles, saw toothed grain beetles), Spider Beetles, Cigarette Beetles, Drug-store beetles, meal worms, Weevils (Granary Weevils, Rice Weevils) and Cadelles; apply in a good spray adjusted to deliver a coarse wet spray at the rate of 1 gallon to 750 square feet of surface. Direct spray into locations where these insects are found: garbage chutes and cans, into cracks & crevices, dark corners, behind and beneath sinks, stoves and refrigerators, and cabinets, along baseboards, and around door and window frames. For silverfish spray bookcases. Spray ant trails, nests and point of entry. Spray insects directly when possible.

Kitchen and Can Insecticide

SPECIAL LOW OIL FORMULA

**FOR USE IN HOMES, INSTITUTIONS,
RESTAURANTS, FOOD PROCESSING PLANTS,
DAIRIES, LIVESTOCK, AND OUTDOORS
FOR MOSQUITO CONTROL**

**ALSO FOR USE IN INDUSTRIAL, AND
RESIDENTIAL GARBAGE COMPACTORS,
GARBAGE CHUTES, KITCHEN GARBAGE
CANS AND OTHER KITCHEN LOCATIONS.**

A PYRENONE® Insecticide-leaves no objectional residues.

ACTIVE INGREDIENTS*	% BY WEIGHT
Pyrethrins.....	0.12
*Piperonyl Butoxide, Technical.....	1.20
Petroleum Distillate.....	0.48
INERT INGREDIENTS.....	98.20
	100.00

*Equivalent to 0.96% (butylcarbityl) (6-propylpiperonyl) ether and 0.24% related compounds

PYRENONE-Registered trademark of Fairfield American Corp.

CAUTION
KEEP OUT OF REACH OF CHILDREN
See side panel for additional precautions

EPA REG. No. 11474-33 EPA EST. No. 11474CA01
Manufactured By:
SUNGRO CHEMICALS, INC.
P.O. Box 24632 Los Angeles, Calif. 90024

NET CONTENTS

GALLONS

To control Carpet Beetles: Spray edge of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, closets and infested areas of clothing. Repeat treatment as needed.

To control Bedbugs: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray to all joints. Treat baseboards, moldings and floors. Repeat treatment as needed.

To control fleas, and brown dog ticks: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.

To control clothes moths: Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining.

In case of infested foods in cabinets, dispose of the infected material. Remove all contents and clean thoroughly. After spray has dried cover shelves with paper or otherwise protect foods from direct contact with sprayed surfaces. Repeat treatment if reinfestation occurs.

SPACE SPRAYING: To kill flies, including Fruit flies, Gnats, Mosquitoes, Flying moths, Hornets, skipper flies, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, and tobacco moths. Close windows and doors and shut off ventilating system. Apply as a fine mist using a non-thermal sprayer. Direct spray at an upward angle distributing it uniformly throughout the entire area at a rate of 1-6 ounces per 1000 cubic feet of space.

FOGGING: To kill exposed or accessible stages of the above named crawling insects, use at a rate of one gallon to 20,000 to 40,000 cubic feet of space. Contact insects directly whenever possible.

ON ANIMALS: To protect cattle and horses from horn flies, mosquitoes and gnats, apply a light mist sufficient to wet the surface of the hair. To control stable flies, horse flies, and deer flies, apply at the rate of 2 ounces per adult cow or horse, sufficient to wet hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

To control blood sucking lice, apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every two to three weeks if required.

OUTDOOR ADULT MOSQUITO CONTROL: Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where mosquitoes may rest, low marshy places, stagnant pools or ponds and shore lines. Apply at the rate of 1 1/2 gallons per acre (4.5 ounces per 1000 square feet).

Using truck-mounted conventional misting equipment such as Rotomist Models 51C, 91 or 100 HT traveling at 5 m.p.h. to treat a 300 foot swath calibrate equipment to deliver 4.5 gallons per minute.

STORAGE AND DISPOSAL

STORAGE: Store in a cool dry place. Keep container closed.
One Gallon or Under:
Pesticide Disposal: Securely wrap original container in several layers of newspaper and discard in trash.
Container Disposal: Do not reuse container. Rinse thoroughly before discarding in trash.
Over 1 Gallon:
Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste pesticide disposal facility.
Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in sanitary landfill, or by other procedure approved by state and local authorities.

ACCEPTED

SEP 23 1985

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 11474-33