PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS STATEMENT OF PRACTICAL TREATMENT

Camtion: Harmful if swallowed.

IF SMALLOWED - Call a physician or poison control center immediately. Gastric 1 avage is indicated if material is taken internally. DO NOT INDUCE VOMITING. Vomiting may cause aspiration pneumonia. IF INHALED - Remove victim to fresh air. Apply respiration if indicated. IF ON SKIN - Remove contaminated clothing and wash area with soap and water. IF IN EYES - Flush eyes with plenty of water and get prompt medical attention.

ENVIRONMENTAL HAZARDS

Avoid contamination of feed and foodstuffs.

This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply only as specified on the label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. Stored food products which are infested should be fumigated or treated by other effective methods using an approved product intended for that purpose.

SPACE SPRAY: For control of accessible, exposed stages of flies, including fruit flies, mosquitoes, small flying moths, gnats, wasps, hornets, cheese skipp ers, and these common stored product pests: granary weevils, rice weevils, confused flour beetles,, saw toothed grain beetles, soider beetles, cigarette beetles, Drugstore beetles, Angoumous grain moths, Hediterranean - flour moths, Indian meal moths, Tobacco moths, meal worms, and grain mites: Close windows and doors and shut off ventilating systems. Use a fogger or vaporizer adjusted to deliver an aerosol spray ino droplets over 50 microns in diameter and 80% less than 30 microns) Apply at the rate of 1



SUNBUGGER SUPER FOG FOOD PLANT FOGGING INSECTICIDE

DESIGNED FOR USE IN FOOD PROCESSING PLANTS

ACTIVE INGREDIENTS

Pyrethrins 0.50% Priperonyl Butoxide, Technical 5.00% Petroleum Distillate 94.50%

* Equivalent to 4.0%(butylcarbityl)(6-propylpiperonyl) ether and to 1.0% related compounds

CAUTION

Keep Out Of Reach Of Children
See rear panel for further precautions.

E.F. A Reg. No. 11474-45

E.P.A. Est. No 11474CA01

SUNGRO CHEMICALS INC.

PO Box 24632

Los Angeles, Catil 90024

MET CONTENTS_____GALLON

MAY 2 1985

ounce per 1000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area and behind obstructions. For best control of wasps and horsets, apply disectly to mests in the evening when the linguists have returned to the nests. Keep the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary.

SPACE AND CONTACT SPRAY: For control of accessible emposed stages of silverfish, trickets, roaches, spiders, scorpions, clo ver mites, cheese mites, earwigs, and ants, first using equipment which can deliver a pin spray, direct spray into all cracks and crevices in woodwork, walls, floors, underneath sinks, behind pipes and in all places where these insects are known to seek shelter. Contact as many insects as possible, spray directly all ant trails. Do not allow spray to contact stored food. Then shutting all doors, windows and shutting off all ventilating systems, proceed with space spraying directions as given above. Use a dosage rate of 1 ounce per 1000 cubic feet of space. Keep the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary.

IN MUSHROOM PRODUCTION AND PROCESSING: To control austroom flies or fungus gnats in growing houses, apply at a rate of 1 ounce per 1000 cubic feet of space. Fogger should be operated 8 to 10 feet away from beds and directed slightly upward over them. Circulating fams should be left in operation, but for best results, refrigeration should be turned off. In packing and canning plants, use at a rate of 1 ounce to 2000 cubic feet of space.

Apply with Equipment Designed and Adjusted to Deliver a True Aerosol Mist.

STORAGE AND DISPOSAL

PESTICIDE DISPOSAL: Festicide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to applicable federal, state, or local procedures.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with the directions given herewith.

51 - 11.411

las 1

110