

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed. Avoid contamination of food and feed stuffs.
The use of this product in food processing plants should be confined to time periods when the plant is not in operation. Foods should be removed or covered before treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.
Remove pets, birds, and cover fish aquaria before spraying.
ENVIRONMENTAL HAZARDS
This product is toxic to fish. Keep out of lakes, streams, or ponds.
Do not contaminate water by cleaning of equipment or disposal of wastes.
Apply this product only as directed on the label.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE: Store in a cool dry place. Keep container closed.
PESTICIDE DISPOSAL: Pesticide or waste that cannot be used or chemically processed should be disposed of in a landfill approved for pesticide or buried in a safe place away from water supplies.

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Do not reuse container (bottle, can, jars). Rinse thoroughly before discarding in trash.



SUNGRO PYRETH

Effective, Fast Acting Insecticide. For Use in Homes, Institutions, Restaurants and Food Processing Plants.

Kills on contact exposed or accessible stages of:

Roaches	Gnats	Rice Weevil	Granary Weevil
Silverfish	Wasp	Spider Beetles	Red Flour Beetle
Crickets	Hornets	Cheese Mites	Clothes Moths
Spiders	Sand Flies	Carpet Beetles	
Ants	Mosquitoes	Angoumois Grain Moth	
Flies	Bedbugs	Confused Flour Beetle	
Caddisflies	Meal Worms	Saw Toothed Grain Beetle	
Flies	Meal Moths	Mediterranean Flour Moth	
Shippers	Grain Mites	Brown Dog Ticks in Buildings	

ACTIVE INGREDIENT:

Pyrethrins	0.1%
Piperonyl Butoxide, Technical*	0.9%
Petroleum Distillate	99.4%
INERT INGREDIENT:	0.0%
TOTAL	100.0%

*Equivalent to 0.4% of (butylcarbitol)(6-propylpiperonyl) ether and to 0.1% of related compounds.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See side panel for additional precautionary statements

EPA Reg No 11474-32

EPA Est No 11474CAD1

Manufactured by
SUNGRO CHEMICALS, INC.

Los Angeles, Calif. 90024

NET CONTENTS _____ GALLONS

ACCEPTED

AUG 23 1984

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under
EPA Reg. No. 11474-32

SPACE SPRAY: For use in homes, institutions, restaurants, and food processing plants. Using a good nozzle spray toward the insects and away from people. Vacuum the area after spraying.

SURFACE APPLICATION: For use in homes, institutions, restaurants, and food processing plants. Apply to surfaces, including window frames, and into cracks and crevices as necessary.

Against an infestation, spray the surfaces of every room. In controlling flies, application toward carpets and floors and bedding should be made.

To control bedbugs, spray sides and bottom of boxes and folds. Spray into all cracks and crevices, floors, closets with particular attention to these places.

For most effective, remove infested or if possible, air out. Thoroughly inspect every corner of the house to be treated. Infested articles at least 3 feet away from the source to avoid contact. Carpets, surfaces of drawers are made of rubber, covering. Repeat treated in each night.

To control infested leather goods, leather shoes, leather bags, leather furniture, etc. Apply to the surface of the leather goods. Repeat treated in each night.

FOG OR VAPOR: For use in homes, institutions, restaurants, and food processing plants. Apply to the surfaces of the room. Repeat treated in each night.

INSTRUCTIONS FOR USE ON DOMESTIC ANIMALS

If feed and feed stuffs, plants should be confined in aeration. Foods should be stored in processing surfaces should be cleaned before using. before spraying.

also, streams or ponds, prevent the disposal of wastes, label.

marker inconsistent with the

to container closed not cannot be used used of in a landfill be place away from

use of this f on site or osal facility.

nt). Then conditioning, f in a other procedures al authorities. ttle, can, jars). discarding in



SUNGRO PYRETH

Effective, Fast Acting Insecticide. For Use in Homes, Institutions, Restaurants and Food Processing Plants.

Kills on contact exposed or accessible stages of:

Roaches	Gnats	Rice Weevil	Granary Weevil
Silverfish	Wasps	Spider Beetles	Red Flour Beetle
Crickets	Hornets	Cheese Mites	Clothes Moths
Spiders	Sand Flies	Carpet Beetles	
Ants	Mosquitoes	Angoumois Grain Moth	
Fleas	Bedbugs	Confused Flour Beetle	
Cadellies	Meal Worms	Saw Toothed Grain Beetle	
Flies	Meal Moths	Mediterranean Flour Moth	
Stinkbugs	Grain Mites	Brown Dog Ticks in Buildings	

ACTIVE INGREDIENT:

Pyrethrins	0.1%
Piperonyl Butoxide, Technical*	0.5%
Petroleum Distillate	99.4%
INERT INGREDIENT:	0.0%
TOTAL	100.0%

*Equivalent to 0.4% of (butylcarbityl)(6-propylpiperonyl) ether and to 0.1% of related compounds.

KEEP OUT OF REACH OF CHILDREN CAUTION

See side panel for additional precautionary statements

EPA Reg No 11474-32

EPA Est No 11474CA01

Manufactured By

SUNGRO CHEMICALS, INC.

Los Angeles, Calif. 90024

NET CONTENTS _____ GALLONS

ACCEPTED

AUG 23 1984

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under

EPA Reg. No. 11474-32

DIRECTIONS FOR USE

SPACE SPRAY: For rapid control of flies, gnats, mosquitoes, and other flying insects. Apply as a coarse misting spray directed into the room to fill the entire area with mist, using 1 to 2 ounces per 1000 cubic feet of space. Ventilate the treated area and vacate before reoccupying. Keep the area closed for 24 hours after spraying. Repeat application when reinfestation occurs.

SURFACE APPLICATION: To control roaches, silverfish, crickets, and spiders. Apply as a coarse misting spray directed into locations where these insects may hide, behind baseboards, moldings, cabinets, shelves, under door and window frames, around plumbing, drains and in dark corners and into cracks and crevices, hitting insects with the spray whenever possible. Repeat treatment as necessary.

Against ant infestations, treat shelves, cabinets and storage areas; spray out runs, walls, baseboards, thresholds and other places of entry, and treat nests liberally. Repeat treatment as necessary.

In controlling flies and Brown Dog ticks in houses, direct the application toward the lower wall portions, baseboards, rugs, carpets and floors of infested areas. Animal quarters, kennels and bedding should also be sprayed.

To control bedbugs effectively, apply liberally to top, ends, sides and bottom of mattresses, spraying thoroughly all beds, rooms and folds. Spray springs and frame, directing the liquid into all cracks and joint joints. Treat walls, baseboards, moldings, floors, closets and bedroom furniture, picture frames, with particular attention to protected bedding hiding and breeding places.

For most effective control of clothes moths and carpet beetles, remove infested articles from storage, brush thoroughly and, if possible, air several hours in direct sunlight. Apply spray liberally in empty closets, closets, bureaus and storage, directing the liquid into cracks, joints and crevices. After airing, infested articles may be treated by holding the spray nozzle at least 3 feet away from the fabric and applying the liquid lightly to the seams to avoid the possibility of staining. Treat both sides of rugs, carpets and pads, and all exposed and accessible surfaces of overstuffed furniture. Do not overspray if products are made of rubber or are rubber coated, to prevent expansion or swelling. Repeat treatment monthly unless the articles are stored in moth-tight containers.

To control infestations of granary weevil, rice weevil, saw-toothed grain beetle, confused flour beetle, red flour beetle, etc. Apply spray liberally to all areas where these insects may be found, including grain bins, silos, and storage areas. If possible, clean out and destroy any infested food products; otherwise, fumigation or other effective treatment should be made using an approved product intended for this purpose. Apply this spray thoroughly to all surfaces of shelves, cabinets, bins, pantries, and similar food storage areas with particular attention to cracks, crevices, niches, and other hiding places.

FOG OR VAPOR SPRAY: To kill larvae and exposed stages of insects named on this label (excepting clothes moths, bedbugs, and carpet beetles). Apply using conventional nebulizer, thermal or compressed air equipment. Fill the entire area with mist using 3 to 6 ounces per 1000 cubic feet of space. Repeat treatment as needed.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed. Avoid contamination of feed and food stuffs. The use of this product in food processing plants should be confined to time periods when the plant is not in operation. Foods should be removed or covered before treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

Remove pets, birds, and cover fish aquaria before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as directed on the label.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE Store in a cool dry place. Keep container closed.
PESTICIDE DISPOSAL Pesticide or rinsate that cannot be used or chemically processed should be disposed of in a landfill approved for pesticide or buried in a safe place away from water supplies.

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Do not reuse container (bottle, can, jars). Rinse thoroughly before discarding in trash.



SUNGRO PYRETH

Effective, Fast Acting Insecticide. For Use in Homes, Institutions, Restaurants and Food Processing Plants.

Kills on contact exposed or accessible stages of:

Roaches	Gnats	Rice Weevil	Granary Weevil
Silverfish	Wasps	Spider Beetles	Red Flour Beetle
Crickets	Hornets	Cheese Mites	Clothes Moths
Spiders	Sand Flies	Carpet Beetles	
Ants	Mosquitoes	Angoumois Grain Moth	
Fleas	Bedbugs	Confused Flour Beetle	
Cadellids	Meal Worms	Saw Toothed Grain Beetle	
Flies	Meal Moths	Mediterranean Flour Moth	
Skippers	Grain Mites	Brown Dog Ticks in Buildings	

ACTIVE INGREDIENT:

Pyrethrins	0.1%
Piperonyl Butoxide, Technical	0.5%
Petroleum Distillate	99.4%
INERT INGREDIENT:	0.0%
TOTAL	100.0%

*Equivalent to 0.4% of (butylcarbityl)(5-propylpiperonyl) ether and to 0.1% of related compounds.

KEEP OUT OF REACH OF CHILDREN CAUTION

See side panel for additional precautionary statements


EPA Reg No 11474-32

EPA Est No 11474CA01

Manufactured By
SUNGRO CHEMICALS, INC.

Los Angeles, Calif 90024

NET CONTENTS _____ GALLONS



SPACE SPRAY. For flies, gnats, mosquitoes, and other flying insects. Using a good nozzle spray toward the insects while misting the space. Vacate the area. Keep the area well ventilated.

SURFACE APPLICATION. Apply to cracks and crevices, under baseboards, around window frames, and into cracks and holes wherever possible.

Against ant infestation, spray all nooks and crannies, on necessary. In controlling fleas, application toward carpets and floors and bedding should be made.

To control bedbugs, sides and bottom of mattress and folds into all cracks and crevices, floors, close with particular attention to places.

For most effective, remove infested or if possible, air out liberally in empty rooming the liquid in infested articles at least 3 feet away to the room to avoid of rash, carpet surfaces or overalls are made of rubber, swelling. Repeat stored in moth-tight.

To control infested, toothed grain beetle and flour beetle, make grain with 1/2 inch in food storage any infested food effective treatment intended for this purpose of shelves storage areas with sacks, and other.

FOG OR VAPOR stages of insects in bedding and carpet (chemical) thermal or even with mist and space. Repeat use.

ENTS
TIC ANIMALS

of feed and food stuffs.
plants should be confined
eration. Foods should be
ed processing surfaces should
eashed before using.
before spraying.

akes, streams or ponds.
peest or disposal of wastes.
label.

anner inconsistent with its

Al
o container closed
not cannot be used
posed of in a landfill
le place away from

use of this
f on site or
osal facility.

nt). Then
conditioning,
f in a
other procedures
al authorities.
tle, can, jars).
discarding in



SUNGRO PYRETH

Effective, Fast Acting Insecticide. For
Use in Homes, Institutions, Restaurants
and Food Processing Plants.

Kills on contact exposed or accessible stages of:

Roaches	Gnats	Rice Weevil	Granary Weevil
Silverfish	Wasps	Spider Beetles	Red Flour Beetle
Crickets	Hornets	Cheese Mites	Clothes Moths
Spiders	Sand Flies	Carpet Beetles	
Ants	Mosquitoes	Angoumois Grain Moth	
Fleas	Bedbugs	Confused Flour Beetle	
Cadellies	Meal Worms	Saw Toothed Grain Beetle	
Flies	Meal Moths	Mediterranean Flour Moth	
Shippers	Grain Mites	Brown Dog Ticks in Buildings	

ACTIVE INGREDIENT:

Pyrethrins	0.1%
Piperonyl Butoxide, Technical*	0.5%
Petroleum Distillate	99.4%
INERT INGREDIENT:	0.0%
TOTAL	100.0%

*Equivalent to 0.4% of (butylcarbityl)((6-propylpiperonyl)
ether and to 0.1% of related compounds.

KEEP OUT OF REACH OF CHILDREN CAUTION

See side panel for additional precautionary statements

EPA Reg No 11474-32 EPA Est No 11474CA01

Manufactured By
SUNGRO CHEMICALS, INC.
Los Angeles, Calif 90024

NET CONTENTS _____ GALLONS

ACCEPTED

AUG 23 1984

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg No 11474-32

DIRECTIONS FOR USE

SPACE SPRAY. For rapid control of flies, fruit flies, house flies, gnats, flying beetles and flies, wasps and hornets. Close doors and windows and shut off ventilating systems. Apply using a good machine adjusted to deliver a fine mist. Direct spray toward the upper portions of the room to fill the entire area with mist, using 1 to 2 ounces per 1000 cubic feet of space. Vacate the treated area and ventilate before reoccupying. Keep the area closed for 24 hours after treating. Repeat applications when reinfestation occurs.

SURFACE APPLICATION. To control roaches, silverfish, crickets and spiders. Apply as a coarse wetting spray directed into locations where these insects may hide behind baseboards, moldings, cabinets, shelves, eaves, door and window frames around plumbing drains and in dark corners and into cracks and crevices, hitting insects with the spray whenever possible. Repeat treatment as necessary.

Against ant infestations, treat shelves, cabinets and storage areas, spray ear runs, trails, baseboards, thresholds and other places of entry, and treat nests liberally. Repeat treatment as necessary.

In controlling fleas and Brown Dog ticks in houses, direct the application toward the lower wall portions, baseboards, rugs, carpets and floors of infested areas. Animal quarters, kennels and bedding should also be sprayed.

To control bedbugs effectively, apply liberally to top, ends, sides and bottom of mattresses, spraying thoroughly all tufts, seams and folds. Spray springs and frame, directing the liquid into all cracks and lock joints. Treat walls, baseboards, moldings, floors, closets and bedroom furniture, picture frames, with particular attention to protected bedbug hiding and breeding places.

For most effective control of clothes moths and carpet beetles, remove infested articles from storage, brush thoroughly and, if possible, air several hours in direct sunlight. Apply spray liberally in empty chests, closets, bureaus and storage, directing the liquid into cracks, joints and crevices. After airing, infested articles may be treated by holding the spray nozzle at least 3 feet away from the fabric and applying the liquid lightly to the seams to avoid the possibility of staining. Treat both sides of rugs, carpets and pads, and all exposed and accessible surfaces of overstuffed furniture. Do not overspray if products are made of rubber or are rubber coated, to prevent expansion or swelling. Repeat treatments monthly unless the articles are stored in airtight containers.

To control infestations of granary weevil, rice weevil, saw-toothed grain beetle, confused flour beetle, rusty grain beetle, red flour beetle, cadellies, meal worms, grain mites, Angoumois grain moth, Indian meal moth and Mediterranean flour moth in food storage areas: If possible, clean out and destroy any infested food products; otherwise, fumigation or other effective treatment should be made using an approved product intended for this purpose. Apply this spray thoroughly to all surfaces of shelves, cabinets, bins, partitions, and similar food storage areas with particular attention to cracks, crevices, niches, and other hiding places.

FOG OR VAPOR SPRAY: To kill accessible or exposed stages of insects named on this label (excepting clothes moths, bedbugs and carpet beetles). Apply using conventional or chemical thermal or compressed air equipment. Fill the entire area with mist using 3 to 6 ounces per 1000 cubic feet of space. Repeat treatment as needed.