

SUNGRO

PYRETH

Effective, Fast Acting Insecticide.
For Use in Homes, Institutions,
Restaurants, Food Processing
Plants, Dairy rooms, Milk Rooms and Dairies.

KILLS ON CONTACT EXPOSED OR
ACCESSIBLE STAGES OF:

Roaches	Gnats	Rice Weevil	Granary Weevil
Silverfish	Wasps	Spider Beetle	Red Flour Beetle
Crickets	Hornets	Cheese Mites	Clothes Moth
Spiders	Sand Flies	Ants	Carpet Beetles
Mosquitoes	Meal Moths	Grain Mites	Angoumois Grain Moth
Fleas	Bedbugs	Skippers	Confused Flour Beetle
Cadellies	Meal Worms	Houseflies	Sawtooth Grain Beetle
Stable Flies	Horse Flies	Horn Flies	Lice
Mediterranean Flour Moth		Brown Dog Ticks in Buildings	

ACTIVE INGREDIENTS

Pyrethrins	0.1%
Piperonyl Butoxide, Technical*	0.5%
INERT INGREDIENTS	99.4%
TOTAL	100.0%

*Equivalent to 0.4% of (butylcarbyl)(5-propylpiperonyl) ether and to 0.1% of related compounds

CONTAINS PETROLEUM DISTILLATE

KEEP OUT OF REACH OF CHILDREN
CAUTION

See Side Panel for Additional Precautionary
Statements

EPA Reg. No. 11474-50 EPA Est. No. 11474CA01

MANUFACTURED BY:
SUNGRO CHEMICALS, INC.
P.O. BOX 24632 LOS ANGELES, CA. 90024

NET CONTENTS _____ GALLONS

PRECAUTIONARY STATEMENTS HAZARDS TO HUMAN & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid contamination of feed and food stuffs

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately
IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists
IF ON SKIN OR CLOTHING: Remove contaminated clothing and wash before reuse. Wash skin with soap and warm water. Get medical attention if irritation persists
IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated
The use of this product in food processing plants should be confined to time periods when the plant is not in operation. Foods should be removed or covered before treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using
Remove pets, birds, and cover fish aquaria before spraying

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as directed on the label

STORAGE AND DISPOSAL

STORAGE: Store in a cool dry place. Keep container closed
PESTICIDE DISPOSAL: Pesticide or insecticide that cannot be used or chemically processed should be disposed of in a landfill approved for pesticide or buried in a safe place away from water supplies.
CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Do not reuse container (bottles, can, jars). Rinse thoroughly before discarding in trash

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling
SPACE SPRAY: For rapid control of flies, fruit flies, mosquitoes, gnats, flying moths, and flies, wasps and hornets. Close doors and windows and shut off ventilating systems. Apply using a good machine adjusted to deliver a fine mist. Direct spray toward the upper portions of the room to fill the entire area with mist, using 1 to 2 ounces per 1000 cubic feet of space. Vacate the treated area and ventilate before reoccupying. Keep the area closed for ten minutes after treating. Repeat application when reinfestation occurs
SURFACE APPLICATION: To control roaches, silverfish, crickets, and spiders. Apply as a coarse wetting spray directed into locations where these insects may hide behind baseboards, mouldings, cabinets, shelves, sinks, door and window frames, around plumbing drains and in dark corners and into cracks and crevices, hitting insects with the spray whenever possible. Repeat treatment as necessary
Against ant infestation, treat shelves, cabinets and storage areas, spray ant runs, trails, baseboards, thresholds, and other places of entry, and treat nests liberally. Repeat treatment as necessary

Page 1 of 1

In controlling fleas and Brown Dog Ticks in houses, direct the application toward the lower wall portions, baseboards, rug, carpets and floors of infested areas. Animal quarters kennels and bedding should also be sprayed
To control bedbugs effectively, apply liberally to top, ends, sides and bottom of mattresses, spraying thoroughly all tufts, seams and folds. Spray springs and frame, directing the liquid into all cracks and lock joints. Treat walls, baseboards, moldings, floors, closets and bedroom furniture, picture frames, with particular attention to protected bedbug hiding and breeding places

For most effective control of clothes moths and carpet beetles, remove infested articles from storage, brush thoroughly and, if possible, air several hours in direct sunlight. Apply spray liberally in empty chests, closets, bureaus and storages, directing the liquid into cracks, joints and crevices. After airing, infested articles may be treated by holding the spray nozzle at least 3 feet away from the fabric and applying the liquid lightly to the seams to avoid the possibility of staining. Treat both sides of rugs, carpet and pads, and all exposed and accessible surfaces of overstuffed furniture. Do not overspray if products are made of rubber or are rubber coated, to prevent expansion or swelling. Repeat treatment monthly unless the articles are stored in moth-tight containers

To control infestations of granary weevil, rice weevil, sawtooth grain beetle, confused flour beetle, rusty grain beetle, red flour beetle, cadellies, meal worms, grain mites, Angoumois grain moth, Indian meal moth and Mediterranean flour moth in food storage areas: If possible, clean out and destroy any infested food products, otherwise, fumigation or other effective treatment should be made using an approved product intended for this purpose. Apply this spray thoroughly to all surfaces of shelves, cabinets, bins, pantries, and similar food storage areas with particular attention to cracks, crevices, niches, and other hiding places


FOG OR VAPOR SPRAY: To kill accessible or exposed stages of insects named on this label (excepting clothes moths, bedbugs, and carpet beetles). Apply using conventional mechanical thermal or compressed air equipment. Fill the entire area with mist using 3 to 6 ounces per 1000 cubic feet of space. Repeat treatment as needed

ON ANIMALS:

To protect animals from horn flies, house flies, mosquitoes and gnats, apply a light mist sufficient to wet the tips of the hair. To control stable flies, horse flies, deer flies, apply 2 ounces per adult animal, sufficient to wet the hair, but not to soak the hide. Repeat treatment once or twice daily or at intervals necessary to give continued protection

To control bloodsucking lice, apply the spray lightly to infected areas, using a stiff brush to get to the base of the hair. Use 2 ounces or less per animal. Repeat every two or three weeks as required

BARN, MILK ROOMS, DAIRY PLANTS: To control flies, mosquitoes and gnats apply as a fog or fine mist, directing the spray toward the ceiling and upper corners of the room until the area is filled with mist, using about 2 ounces per 1000 cubic feet of space. For best results, close windows and doors before spraying and keep closed for ten to fifteen minutes. Vacate the area and ventilate before reoccupying. Repeat application as necessary. 74.50 12-93nq



BEST AVAILABLE COPY