

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Harmful if swallowed. Avoid contamination of feed and foodstuffs. Remove pets and birds, and cover aquaria before spraying. Do not get in eyes. In case of contact, immediately flush eyes with plenty of water. Get medical attention if irritation persists.

**STATEMENT OF PRACTICAL TREATMENT**

IF SWALLOWED: Call a physician or poison control center immediately. Drink one or two glasses of water and induce vomiting by touch back of throat with finger. DO NOT INDUCE VOMITTING or give anything by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated.

IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and water.

IF IN EYES: Flush eyes with water for 10-15 minutes. Contact a physician if irritation persists.

**ENVIRONMENTAL HAZARDS**

This product may be toxic to fish. Keep out of lakes, ponds, streams, tidal marshes and estuaries. Do not contaminate water by cleaning of equipment or disposal of wastes.

**DIRECTIONS FOR USE**

It is a violation of Federal law use this product in a manner inconsistent with it's labeling.

**INDOOR APPLICATION**

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. Vacate treated areas and ventilate before reoccupying. For maximum effectiveness, a combination of spot surface treatment and fogging is recommended.

**SURFACE SPRAYING:** To control of roaches, ants, silverfish, spiders, crickets, clover mites, Cheese Mites, granary weevils, rice weevils, confused flour beetles, red flour beetles, saw-toothed grain Beetles, spider beetles, cigarette beetles, drugstore beetles, meal worms, grain mites and caddisflies. Use in a good sprayer adjusted to a coarse wet spray. Direct spray into hiding places, crack & crevices under pallets, around containers of stored foods, around the base of machinery and behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests and points of entry. If surface application only is to be used, spray floors and other surfaces, applying at a rate of 1 gallon to 750 square feet of surface.

To control carpet beetles: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, closets and infested areas of shelving. Repeat treatment as needed.

To control bedbugs: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, mouldings and floors. Repeat treatment as needed.

To control fleas and brown dog ticks: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along behind baseboards, mouldings, window and door frames and entire areas of floor and floor covering.

# SUNGRO #214 WATER-BASE MULTI-PURPOSE INSECTICIDE

A Concentrated PYRENONE<sup>®</sup> Micro-Emulsion  
Effective, Fast-Acting Insecticide  
For Use in Homes, Institutions, Restaurants,  
Food Processing Plants, Dairies, Livestock  
and Outdoors for Mosquito Control  
Apply with Conventional Mechanical or  
Compressed Air Equipment

**ACTIVE INGREDIENTS:**

Pyrenones 0.1%

\*\*Piperonyl Butoxide, Technical 1.0%

**INERT INGREDIENTS**

98.90%

Total 100.00%

\*\*Equivalent to 0.2% [(butylcarbitol) o-propyl(piperonyl)] ether and 0.2% related compounds.

PYRENONE<sup>®</sup> is a Registered trademark of Fairfield American Corporation.


## CAUTION KEEP OUT OF REACH OF CHILDREN

(SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONS)

Do Not Transport or Store Under 32° F.  
EPA Reg. No. 11474-51 EPA Est. No. 11474CA01

Buyer assumes all risks of use storage or handling of this product not in strict accordance with directions given herewith.

Manufactured By:  
SUNGRO CHEMICALS, INC.  
P.O. Box 24632 Los Angeles, CA 90024  
NET CONTENTS \_\_\_\_\_ GALLONS



Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.

To control clothes moths: Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight if possible. Apply spray liberally to empty chests, closets, bureaus and other storages, directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining.

**SPACE SPRAYING:** Close doors and windows and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, Angoumois grain moths, Mediterranean flour moths, Indian meal moths and tobacco moths: Direct spray at an upward angle, distributing it uniformly throughout the entire area at a rate of 3-6 ounces per 1000 cubic feet of space.

**FOGGING:** To kill exposed or accessible stages of the above named crawling insects, use at a rate of one gallon to 20,000 to 40,000 cubic feet of space. Contact insects directly whenever possible.

**ON ANIMALS:** To protect cattle and horses from horn flies, mosquitoes and gnats, apply at the rate of 2 ounces per adult cow or horse, sufficient to wet hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

**OUTDOOR APPLICATION**

**OUTDOOR ADULT MOSQUITO CONTROL:** Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where mosquitoes may rest, low marshy places, stagnant pools or ponds and shore lines. Apply at the rate of 1 1/2 gallons per acre (4 1/2 ounces per 1000 square feet). Using truck-mounted conventional misting equipment such as Rotomist Models 51c, 91 or 100 HT, traveling at 5 m.p.h. to treat a 300 foot swath calibrate equipment to deliver 4.5 gallons per minute.

Not for use in USDA Meat and Poultry Processing Plants.

**STORAGE AND DISPOSAL**

**STORAGE:** Store in a cool, dry place. Keep container closed. One gallon or under:

**PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash.

**CONTAINER DISPOSAL:** Do not reuse container. Rinse thoroughly before discarding in trash.

Over one gallon:

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.