

| | | | | | | | |
|--|---|---|--|------------------|--|---|--|
| U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (TS-767) WASHINGTON, DC 20460 | <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%; padding: 2px;">EPA REGISTRATION NO. 11474-67</td> <td style="width: 50%; padding: 2px;">DATE OF ISSUANCE MAR 30 1987</td> </tr> <tr> <td colspan="2" style="padding: 2px;">TERM OF ISSUANCE</td> </tr> <tr> <td colspan="2" style="padding: 2px;">NAME OF PESTICIDE PRODUCT Sungro Pernith With Permanone</td> </tr> </table> | EPA REGISTRATION NO. 11474-67 | DATE OF ISSUANCE MAR 30 1987 | TERM OF ISSUANCE | | NAME OF PESTICIDE PRODUCT Sungro Pernith With Permanone | |
| EPA REGISTRATION NO. 11474-67 | DATE OF ISSUANCE MAR 30 1987 | | | | | | |
| TERM OF ISSUANCE | | | | | | | |
| NAME OF PESTICIDE PRODUCT Sungro Pernith With Permanone | | | | | | | |
| NOTICE OF PESTICIDE: <input type="checkbox"/> REGISTRATION <input type="checkbox"/> REREGISTRATION (Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended) | | | | | | | |
| NAME AND ADDRESS OF REGISTRANT (Include ZIP code) | | | | | | | |
| <div style="display: flex; justify-content: space-between;"> <div style="width: 80%;"> <p style="margin: 0;">Sunagro Chemical, Inc.</p> <p style="margin: 0;">P.O. Box 24632</p> <p style="margin: 0;">Los Angeles, CA 90024</p> </div> <div style="width: 15%; text-align: right; font-size: 1.2em;"> 161 / 189054 16 / 1 </div> </div> | | | | | | | |
| NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number. | | | | | | | |
| On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act. | | | | | | | |
| A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith. | | | | | | | |
| Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others. | | | | | | | |
| This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you: | | | | | | | |
| <ol style="list-style-type: none"> 1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data. 2. Make the labeling changes listed below before you release the product for shipment: <ol style="list-style-type: none"> a. Add the phrase "EPA Registration No. 11474-69." b. Add the statement, "Recommended for Commercial/Industrial Use" to the front panel of the labeling. c. Delete the statement beginning, "All food processing surfaces should be covered . . . etc." The product is <u>not</u> for use in edible product areas of food/feed processing plants. This was brought to your attention in our last correspondence. | | | | | | | |
| <div style="border: 2px solid black; padding: 5px; display: inline-block; font-weight: bold; font-size: 1.1em;">BEST AVAILABLE COPY</div> | | | | | | | |
| <input type="checkbox"/> ATTACHMENT IS APPLICABLE | | | | | | | |
| SIGNATURE OF APPROVING OFFICIAL | DATE 3/30/87 | | | | | | |

- d. Add the following precaution under the "Precautionary Statements" paragraph.

Wear full-faced gas mask with canister type recommended for general insecticide protection when applying indoors as a space spray or fog.

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

George T. LaRocca
Product Manager (15)
Insecticide-Rodenticide Branch
Registration Division (TS-767C)

Enclosures

LUNGRO PERMITH

WITH PERMANONE

A Concentrated PERMANONE® Stable Emulsion
Effective, Fast Acting Insecticide

For Use Outdoors, and in Homes, and Nonfood Areas
of Kennels, Commerical Buildings, Institutions,
Warehouses, Theaters, Office Buildings, Schools,
Motels, Hotels, Restaurants, and Food/Feed Processing
Establishments.

For Control of Fleas, Brown Dog Ticks and Lice
On Premises and On Pets

Apply With Conventional Mechanical Or
Compressed Air Equipment

ACTIVE INGREDIENTS:

| | |
|--------------------|---------|
| *Permethrin | 8.50% |
| INERT INGREDIENTS: | 99.50% |
| | 100.00% |

2-(1-phenoxyphenyl) methyl (2) cis/trans 3-(2,2-dichloroethenyl)-
2,2-dimethylcyclopropanecarboxylate. Cis/trans ratio: min.
35% (2) cis and max. 65% (2) trans.

**KEEP OUT OF REACH OF CHILDREN.
CAUTION**

See rear panel for use cautions.

EPA Reg. No. 11474

EPA Est. No. 11474CA01

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a man-
ner inconsistent with its labeling.

SHAKE WELL BEFORE USE

INDOOR APPLICATIONS

Do not use in the edible product areas of food/feed
processing plants, restaurants, or other areas
where food/feed is commercially prepared or
processed. Do not use in areas while food is
exposed.

All food processing surfaces should be covered during
treatment and thoroughly cleaned and rinsed with potable wa-
ter, before using.

Nonfood/Feed Areas includes garbage rooms, lavatories, floor
drains (to sewers), entries and vestibules, offices, locker rooms,
machine rooms, boiler rooms, garages, mop closets, etc.

Vacate treated areas and ventilate before reoccupying. For
maximum effectiveness, a combination of spot, surface treat-
ment and fogging is recommended.

SURFACE SPRAYING: To control co-
burs, scorpions, millipedes, cen-
turs, silverfish, firebrats, and
cheese mites, granary weevils, r-
beetles, rust red flour beetles,
der beetles, cigarette beetles,
grain mites and cadellies: Use a
coarse wet spray. Direct the sp-
and crevices, under pallets, and
around the base of machinery, ar-
For silverfish, spray bookcases,
nests and points of entry. If
be used, spray floors and other
of 1 gallon to 750 square feet

To control carpet beetles: Spr-
carpeting and rugs. Make local
baseboards. Spray directly into
areas of shelving. Repeat treat-

To control bedbugs: Spray mat-
around tufts and seams. Take b-
joints. Treat baseboards, moul-
ment as needed.

To control fleas and ticks (adu-
spray infested areas, pet beds,
cracks and crevices, along and
window and door frames, and ent-
covering. Fresh bedding should
following treatment. Repeat tr-

To control clothes moths: Remo-
storage, brush thoroughly, and
light, if possible. Apply spra-
closets, bureaus and other stor-
cracks, joints and crevices. A
may be treated lightly, keeping
from fabric to avoid staining.

To kill lice and louse eggs: S-
to rest for possible staining o-
after drying, then proceed to s-
Spray from a distance of 5 to 1
garments and parts of bedding,
ture that cannot be either laun-
all treated articles to dry tho-

SPACE SPRAYING: Close doors and
tine systems. Apply with a fog-
a fine mist or fog. For rapid
flies, gnats, mosquitoes, skip-
black flies, Anomalous train a-
Indian meal moths and tobacco
ward angle, distributing it un-
area at a rate of 1 ounce per
area closed for 2 hours. Vent-

CONTINUED

MAR 30 1987

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
As amended, for the pesticide
registered under EPA Reg. No.
11474-607

BES

FOGGING: To kill closed or accessible stages of the above named crawling insects, use a rate of 1 gallon to 20,000 to 40,000 cubic feet of space. Contact insects directly when possible. Keep area closed for 2 hours. Ventilate before re-occupying.

ON LIVESTOCK: To protect cattle (beef and dairy), goats, sheep, hogs and horses from horn flies, face flies, house flies, mosquitoes and gnats: Apply a light mist sufficient to wet the surface of the hair. For face flies, spray face and head, but do not spray into eyes.

To control stable flies, horse flies and deer flies: Apply at a rate of 2 ounces per adult animal, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

To control blood-sucking lice: Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

To control poultry lice: Spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

To control bedbugs and mites in poultry houses: Spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

ON PETS: To control fleas, ticks and lice: Start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and underbody. While spraying, fluff the hair so that the spray will penetrate to the skin. Make sure spray wets thoroughly. Do not spray into eyes or face. Avoid contact with genitalia. Repeat as necessary.

NOTE: Not recommended for use on kittens or puppies less than four weeks old.

OUTDOOR APPLICATION

FOR CONTROL OF GARDEN INSECTS: For use on roses, dahlias, asters and other ornamentals to kill Japanese beetles, aphids, exposed thrips, spiders, red mites and leafminers.

May be used on plants, as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, asters, azaleas, begonias, cactillias, carnations, chrysanthemums, dahlias, delphiniums, dogwoods, English ivy, amononymus, fuschia geraniums, crassula, grape vines, Kentia palms, laurels, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering Jews and zinnias.

Effective when used as directed against whiteflies, aphids, spider and red mites, exposed thrips, lace bugs, Japanese beetles, pavement ants, clover mites and army worms.

When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. Repeat treatment as infestations occur.

ADULT MOSQUITO CONTROL: Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where mosquitoes may rest (low, marshy places). Apply at a rate of 2.5

pints per acre (0.9 ounce per 1000 square feet). Use mounted conventional misting equipment such as 3000 51C, 91 or 100 HT traveling at 5 m.p.h. to treat a 3/4 swath, calibrate equipment to deliver 0.91 gallon per acre. Under no conditions should open water be treated with duct.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE AND SPILL PROCEDURES: Store upright. Do not expose to freezing temperatures. Do not expose to extreme temperatures. In case of spill, soak up with an absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste.

PESTICIDE DISPOSAL: Pesticide, spray mixture or residue that cannot be used according to label instructions should be disposed of at or by an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent) then for recycling or reconditioning, or puncture and discard in a sanitary landfill, or by other approved State and Federal procedures.

CONTAINERS ONE GALLON AND SMALLER: Do not transport below 32°F. Do not reuse container. Wrap containers in layers of newspaper and discard in trash.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid contamination of feed and other stuffs. Remove pets and birds, and cover fish aquariums before spraying or surface applications.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink one to two glasses of water and vomit by touching the back of the throat with finger until vomit fluid is clear. Call a physician if necessary. Do not induce vomiting or give anything by mouth to unconscious person.

IF INHALED: Remove affected person to fresh air. Give artificial respiration if indicated.

IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and water.

IF IN EYES: Flush eyes with plenty of water. Call physician if irritation persists.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply direct to ponds, streams, tidal marshes or estuaries. Do not contaminate water by cleaning of equipment or disposal of waste.

Buyer assumes all risks of use, storage or handling of product not in strict accordance with directions given with.

Manufactured by:

SUNGRO CHEMICALS, INC.
P.O. BOX 24632 LOS ANGELES, CA 90024

NET CONTENTS:

BEST AVAILABLE COPY