

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Do not pour or spill near heat or open flame. Remove pets, birds and cover fish aquaria before spraying. Do not breathe spray mist. Wear mask or respirator of a type recommended by the Mine Safety and Health Administration to give adequate protection against this material. Wear protective clothing. Wash exposed skin surfaces with soap and water.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Never give anything by mouth to an unconscious person.

IF INHALED: Remove affected person to fresh air. Apply artificial respiration if indicated.

IF IN EYES: Flush eyes with water for 10-15 minutes. Contact a physician if irritation persists.

IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and water. Contact a physician if irritation persists. Wash clothing before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

STORAGE AND DISPOSAL

For containers larger than one gallon:

Do not contaminate water, food or feed by storage or disposal. PESTICIDE STORAGE AND SPILL PROCEDURES: Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with an absorbent material such as sand, sawdust, earth, fuller's earth etc. Dispose of with chemical waste.

PESTICIDE DISPOSAL: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of at a by an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent) then offer for recycling, reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved State and local procedures.

For containers one gallon and smaller:

STORAGE: Store in original container in a cool dry area, inaccessible to children and pets and away from heat and direct sunlight. Protect from freezing.

DISPOSAL: Wrap container in several layers of newspaper and discard in trash. Do not reuse empty container.

BEST AVAILABLE COPY

SUNGRO WATER BASE FOOD PLANT FOGGING INSECTICIDE

A CONCENTRATED PYRENONE[®] MICRO EMULSION

- CLEAR
- STABLE
- LEAVES NO OILY OR OBJECTIONABLE RESIDUE

DESIGNED FOR USE IN EDIBLE PRODUCT AREAS OF OFFICIAL ESTABLISHMENTS OPERATING UNDER THE MEAT POULTRY, SHELL EGG GRADING AND EGG PRODUCTS INSPECTION PROGRAMS AND IN OTHER FOOD PROCESSING PLANTS.

APPLY WITH A CONVENTIONAL MECHANICAL OR COMPRESSED AIR EQUIPMENT DESIGNED AND ADJUSTED TO DELIVER A TRUE AEROSOL MIST.

ACTIVE INGREDIENTS:

Pyrethrins 0.5%
*Piperonyl Butoxide, Technical 5.0%

INERT INGREDIENTS: 94.5%
TOTAL 100.0%

*Equivalent to 4.0% (butylcarbityl) (6-propylpiperonyl) ether and 1.0% related compounds.

PYRENONE[®] - Registered Trademark of Fairfield American Corporation.

KEEP OUT OF REACH OF CHILDREN CAUTION

See Side Panel For Additional Precautions

DO NOT TRANSPORT OR STORE UNDER 32 DEG. F

E.P.A. Est.No. 11474CA01 E.P.A. Reg.No. 11474-78

NET CONTENTS: _____ GALLONS

Manufactured By:
SUNGRO CHEMICALS, INC.
P.O. BOX 24632
LOS ANGELES, CA 90024

DIRECTION FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment and thoroughly cleaned with an effective cleaning compound followed by a potable water rinse before using.

Close windows and doors and shut off ventilating systems. Use a non-thermal fogger or vaporizer adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 30 microns). Apply at the rate of 1 ounce per 1000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area and behind obstructions. the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary.

IN BARN, POULTRY HOUSES AND STORAGE ROOMS: To control HOUSE FLIES, STABLE FLIES and HORN FLIES in barns, poultry houses, warehouses, storage bins and storage rooms: dilute 1 part of material with 4 parts of water. Apply as a fine mist over the livestock in stanchions or holding pens. Use a fogger or mist generator to direct spray above the animals and to the center of the area where animals are housed. Apply 1 ounce per 1000 cubic feet uniformly throughout the entire area sprayed. Do not ventilate the area for at least 10 minutes after spray has been applied. If used in dairy barns, ventilate the barn thoroughly before milking. Repeat daily in the morning and afternoon, or as required.

The above described application will kill accessible, exposed stages of FLIES, including FRUIT FLIES, MOSQUITOES, SILVERFISH, FLYING MOTHS, GNATS, CRICKETS, COCKROACHES, SPIDERS, SCORPIONS, WASPS, HORNETS, CLOVER MITES, CHEESE MITES, CHL SKIPPES and these common stored product pests: GRAY-NARY WEEVILS, RICE WEEVILS, CONFUSED FLOUR BEETLES, SAW-TOOTHED GRAIN BEETLES, SPIDER BEETLES, CIGAR' RTE BEETLES, DRUGSTORE BEETLES, ANGOUMOIS GRAIN MOTHS, MEDITER-RANEAN FLOUR MOTHS, INDIAN MEAL MOTHS, TOBACCO MOTHS, MEAL MOTHS AND GRAIN MITES.

IN MUSHROOM PRODUCTION AND PROCESSING: To control adult MUSHROOM FLIES, FUNGUS GNATS and CECID FLIES in growing houses, apply at a rate of 1 ounce per 1000 cubic feet of space. Fogger should be operated 8 to 10 feet away from beds and directed slightly upward over them. Circulating fans should be left in operation, but for best results refrigeration should be turned off. In packing and canning plants use at a rate of 1 ounce to 2000 cubic feet of space.