

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS & DOMESTIC ANIMALS  
CAUTION**

Harmful if swallowed. Do not pour or spill near heat or open flame. Remove pets, birds and cover fish aquaria before spraying. Do not breathe spray mist. Wear mask or respirator of a type recommended by the Mine Safety and Health Administration to give adequate protection against this material. Wear protective clothing. Wash exposed skin surfaces with soap and water.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Do not apply directly to lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

**FIRST AID:**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses if present after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 9-1-1 or an ambulance then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

Questions? 1-800-463-9179 or call poison control center.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in a dry place no lower in temperature than 32°F or higher than 120°F.

**DISPOSAL INSTRUCTIONS:**

If Empty: Do not reuse this container. Place in trash or offer for recycling if available.

If Partly Filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Buyer assumes all risks of use storage or handling of this product not in strict accordance with directions given herewith.

**SUNGRO  
WATER BASE  
FOOD PLANT  
FOGGING  
INSECTICIDE**

A CONCENTRATED PYRENONE® MICRO EMULSION

**CLEAR  
STABLE  
LEAVES NO OILY OR  
OBJECTIONABLE RESIDUE**

DESIGNED FOR USE IN EDIBLE PRODUCT AREAS OF OFFICIAL ESTABLISHMENTS OPERATING UNDER THE MEAT POULTRY, SHELL EGG GRADING AND EGG PRODUCTS INSPECTION PROGRAMS AND IN OTHER FOOD PROCESSING PLANTS.

APPLY WITH A CONVENTIONAL MECHANICAL OR COMPRESSED AIR EQUIPMENT DESIGNED AND ADJUSTED TO DELIVER A TRUE AEROSOL MIST.

| ACTIVE INGREDIENTS:            | BY WEIGHT |
|--------------------------------|-----------|
| Pyrethrins                     | 0.5%      |
| *Piperonyl Butoxide, Technical | 5.0%      |
| INERT INGREDIENTS:             | 94.5%     |
| TOTAL                          | 100.0%    |

\*Equivalent to 4.0% (butylcarbitol) (6-propylpiperonyl) ether and 1.0% related compounds.

PYRENONE® - Registered Trademark of The Aventis Group.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See Side Panel For Additional Precautions  
DO NOT TRANSPORT OR STORE UNDER 32°F.

E.P.A. Reg.No. 11474-78

E.P.A. Est.No. 11474CA01

Manufactured By:

SUNGRO CHEMICALS, INC.

P.O. BOX 24632 LOS ANGELES, CA 90024

NET CONTENTS: \_\_\_\_\_ GALLONS ( \_\_\_\_\_ LITERS)

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment and thoroughly cleaned with an effective cleaning compound followed by a potable water rinse before using.


Close windows and doors and shut off ventilating systems. Use a non-thermal fogger or vaporizer adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 30 microns). Apply at the rate of 1 ounce per 1000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area and behind obstructions. Keep the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary.

IN BARN, POULTRY HOUSES AND STORAGE ROOMS: To control HOUSE FLIES, STABLE FLIES and HORN FLIES in barns, poultry houses, warehouses, storage bins and storage rooms: dilute 1 part of material with 4 parts of water. Apply as a fine mist over the livestock in stanchions or holding pens. Use a fogger or mist generator to direct spray above the animals and to the center of the area where animals are housed. Apply 1 ounce per 1000 cubic feet uniformly throughout the entire area sprayed. Do not ventilate the area for at least 10 minutes after spray has been applied. If used in dairy barns, ventilate the barn thoroughly before milking. Repeat daily in the morning and afternoon, or as required.

The above described application will kill accessible, exposed stages of FLIES, including FRUIT FLIES, MOSQUITOES, SILVERFISH, FLYING MOTHS, GNATS, CRICKETS, COCKROACHES, SPIDERS, SCORPIONS, WASPS, HORNETS, CLOVER MITES, CHEESE MITES, CHEESE SKIPPERS and these common stored product pests: GRANARY WEEVILS, RICE WEEVILS, CONFUSED FLOUR BEETLES, SAW-TOOTHED GRAIN BEETLES, SPIDER BEETLES, CIGARETTE BEETLES, DRUGSTORE BEETLES, ANGOUMOIS GRAIN MO, MEDITERRANEAN FLOUR MOTHS, INDIAN MEAL MC TOBACCO MOTHS, MEAL MOTHS AND GRAIN MITES.

IN MUSHROOM PRODUCTION AND PROCESSING: To control adult MUSHROOM FLIES, FUNGUS GNATS and CEDIC FLIES in growing houses, apply at a rate of 1 ounce per 1000 cubic feet of space. Fogger should be operated 8 to 10 feet away from beds and directed slightly upward over them. Circulating fans should be left in operation, but for best results refrigeration should be turned off. In packing and canning plants use at a rate of 1 ounce to 2000 cubic feet of space.

013103lam



11474-78

4/15/2003

Page 181