

 <p>U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (H7505C) 401 "M" St., S.W. Washington, D.C. 20460</p> <p><b>NOTICE OF PESTICIDE:</b>  <u>  X  </u> Registration  <u>     </u> Reregistration</p> <p>(under FIFRA, as amended)</p>	EPA Reg. Number:  <b>11474-95</b>	Date of Issuance:  <b>AUG 20 2003</b>
	Term of Issuance:  <b>Conditional</b>	
	Name of Pesticide Product:  <b>Sunbugger Flea &amp; Mite Spray</b>	
Name and Address of Registrant (include ZIP Code): <b>Sungro Chemicals, Inc. P.O. Box 24632 Los Angeles, CA 90024</b>		
<p><b>Note:</b> Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</p>		
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.</p> <p>Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p><b>This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:</b></p> <ol style="list-style-type: none"> <li>1. Submit and/or cite all data required for registration/reregistration of your product when the Agency requires all registrants of similar products to submit data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.</li> <li>2. Make the labeling changes listed below before you release the product for shipment:             <ol style="list-style-type: none"> <li>a. Add the phrase "EPA Registration No. 11474-95".</li> <li>b. Insert the following statement at the top rear or side panel in bold, all caps. <b>READ ENTIRE LABEL BEFORE EACH USE.</b></li> <li>c. Move the paragraph beginning <b>Non Food areas are</b> xxx etc., beneath the paragraph beginning, <b>Do not use in food handling</b> xxx etc.</li> </ol> </li> </ol>		
Signature of Approving Official:		Date:

page 2

EPA Reg. No. 11474-95

d. Make a separate paragraph for the statement beginning,  
We recommend that those with asthma xxx etc.

e. Under first sentence of house dust mite directions add  
on treated surfaces between the words mites and that.

3. Submit two (2) copies of your final printed labeling before  
you release the product for shipment.

If these conditions are not complied with, the registration  
will be subject to cancellation in accordance with FIFRA section  
6(e). Your release for shipment of the product constitutes  
acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Enclosure

# SUNBUGGER FLEA & MITE SPRAY

**Controls Residential, Industrial and Garden Insects.  
For Use on Dogs and on Premises to Control Fleas and Ticks.  
Controls Fleas and Ticks on Dogs for 14 Days.**

**A Water-Based Aerosol.**

**Controls Pests for up to Four Weeks.**

**Kills Lice and Their Eggs on mattresses, furniture, car interior  
and other non-washable items.**

**Kills House Dust Mites in Carpets, Curtains and on Upholstery.**

**Controls House Dust Mites (Dermatophogoides spp) for up to 8  
weeks in Carpets Curtains and on Upholstery**

**Controls House Dust Mites that may cause Asthma, Hay Fever,  
Rhinitis, Watery Eyes, and Sneezing.**

**Kills Screw Worms on Beef Cattle, Horses and Sheep.  
Not For Use On Humans.**

ACTIVE INGREDIENT:	% By Weight
Permethrin*	0.50%
OTHER INGREDIENTS	<u>99.50%</u>
Total	100.00%

\*(3-phenoxyphenyl)methyl +/- cis/trans 3-(2,2-dichloroethenyl)2,2 dimethyl-cyclopropanecarboxylate cis/trans ratio: min. 35% +/- cis and max. 65% +/- trans.

PRECAUCION AL CONSUMIDOR: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

(To THE USER: If you cannot read English, do not use this product until the label has been fully explained to you.)

**KEEP OUT OF REACH OF CHILDREN**

## CAUTION

See Side Panel for Additional Precautions

EPA Reg. No. 11474-

EPA Est. No. 11474CA01

NET WEIGHT: \_\_\_\_\_ OZ. ( g)

## PRECAUTIONARY STATEMENTS Hazards To Humans and Domestic Animals CAUTION

Harmful if absorbed through skin. Avoid inhalation of spray mist. Causes eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish aquaria before space spraying or surface applications. This product is not for use on humans. Vacate room after treatment and ventilate before reoccupying. Do not allow children or pets to contact treated areas until surfaces are dry.

### Environmental Hazards

This product is toxic to fish. Do not apply directly to any body of water. Do not contaminate water when disposing of equipment washwaters.

### Physical Or Chemical Hazards

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F (54°C) may cause bursting.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Do not use in food areas of food handling establishments, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as dining rooms, but excluding areas where foods may be prepared or held.

In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed. We recommend that those with asthma or severe allergies consult their doctor before using this product and have someone else apply this product.

Nonfood areas are areas such as garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

ACCEPTED  
with COMMENTS  
in EPA Letter Dated

AUG 20 2003

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

11474-95

**SHAKE WELL BEFORE USING**

Remove protective cap, hold container upright and spray from a distance of 12 to 15 inches. Remove birds and cover fish aquariums before spraying.

**Houseflies, Fruit Flies, Mosquitoes, Gnats, Black Flies, Skipper Flies, Blow Flies and Small Flying Moths:** Close all doors and windows and spray room at the rate of 2 to 3 seconds per 1000 cubic feet. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Retreat if infestation occurs, but not more than once every two weeks.

**Cockroaches, Waterbugs, Palmetto Bugs, Silverfish, Firebrats, Crickets, Scorpions, Spiders, Millipedes, Centipedes, Sowbugs and Pillbugs:**

Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray when ever possible. Spray to the point of dampness but not run-off. Retreat if infestation occurs, but not more than once every two weeks

**Ants:** Spray trails, nests and points of entry. Spray on ants where possible. Retreat if infestation occurs but not more than once every two weeks.

**Indian Meal Moths, Angoumois Grain Moths:** Close doors and windows, direct spray into all parts of room especially around stored food product containers, pallets and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep area closed for 15 minutes following treatment. Spray to the point of dampness but not run off. Retreat if infestation occurs, but not more than once every two weeks.

**Exposed stages of Confused Flour Beetles, Saw-Toothed Grain Beetles, Red Flour Beetles, Rice Weevils, Granary Weevils, Grain Mites:** Spray into cracks and crevices, around cartons or containers or other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Spray to the point of dampness but not run-off. Retreat if infestation occurs, but not more than once every two weeks. Infested stored food products in the home should be disposed of.

**Carpet Beetles:** Spray directly on beetles, floors, baseboards, shelves, and carpets. Spray to the point of dampness but not run-off. Retreat if infestation occurs, but not more than once every two weeks.

**Bedbugs:** Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, moldings and floors. If lice infestations should occur on humans, use a product labeled for use on humans. Spray to the point of dampness but not run-off. After mattress is dry cover with mattress cover and sheet. Do not use mattress without cover. Retreat if infestation occurs, but not more than once every two weeks.

**To control Clothes Moths and Carpet Beetles on wool and other proteinaceous fibers:** Remove infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. Spray to the point of dampness but not run-off. After airing, infested articles may be treated by spraying lightly, keeping nozzle at least three feet from fabric to avoid staining.

**To Control House Dust Mites (Dermatophagoides spp.):** Sunbugger Flea & Mite Spray controls house dust mites that may cause asthma, hay fever, rhinitis, watery eyes and sneezing. One treatment lasts for up to 8 weeks. On Upholstered Furniture - test fabric in an inconspicuous area to insure dyes will not bleed. Hold container upright with nozzle away from you. Depress valve and spray from a distance of 8 to 10 inches. Spray all fabric surfaces until damp. Spray to the point of dampness but not run-off. Do not use on delicate fabrics. Allow furniture to dry thoroughly before using. On Carpets - Spray over entire carpet until damp. Spray to the point of dampness but not run-off. Hold the nozzle approximately 12 to 18 inches from the carpet surface and spray in a sweeping motion. Use an overlapping pattern to insure total coverage. After application, brush the surface of the carpet with a broom or brush to insure deep penetration. Allow carpet to dry thoroughly before reentering the treated area. Drying time can be accelerated by using a fan. On Mattresses - spray the top, bottom and sides of mattresses until

damp, paying particular attention to tufts and seams. Allow to dry thoroughly before using. After mattress is dry cover and sheet. Do not use mattress without cover. Vacuum entire area once dry. Since vacuuming stirs dust up in the air those with allergies should have another person vacuum or should wear a dust mask and use vacuum with a high efficiency particulate (HEPA) filter.

**Fleas and Ticks (Adults and Larvae)** Remove and thoroughly clean or destroy pet bedding. Spray pet sleeping quarters bed entire floor and floor covering with special attention to cracks and crevices around baseboards and window and door frames. Infested upholstered furniture may also be treated with this product. Spray to the point of dampness but not run-off. A test application should be made to upholstery or drapery fabrics in an inconspicuous place before use.

**To Kill Lice and Louse Eggs:** Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Hold container upright with nozzle away from you. Depress valve and spray from a distance of 8 to 10 inches. Spray each square foot for three seconds. Spray to the point of dampness but not run-off. Allow all sprayed articles to dry thoroughly before use.

Spray only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Do not use on sheets or pillowcases.

**READ ENTIRE LABEL BEFORE EACH USE  
USE ONLY ON DOGS**

**To control Fleas, Ticks and Lice on dogs.** Starting at the tail, stroke against the lay of the hair, spraying the parted fur in short bursts, directly behind the hand to insure penetrating the coat. Dampen the fur up to the head. Retreat if infestation occurs, but not more than once every two weeks. Do not spray directly in/on eyes, mouth or genitalia of pets. Do not use on dogs under twelve (12) weeks old. Consult a veterinarian before using this product on medicated, debilitated, aged, pregnant or nursing animals. Sensitivities may occur after using any pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

**IN ANIMAL QUARTERS AND MILK ROOMS:** For rapid control of House Flies, Mosquitoes and Small Flying Moths, close windows and doors and spray at the rate of 2 seconds per 1000 cubic feet of space. For best results, keep area closed for 10 minutes after spraying. Do not remain in treated area and ventilate after treatment is completed.

**TO TREAT LIVESTOCK:** To protect beef cattle, dairy cattle, goats, sheep, hogs and horses from attacks of Stable Flies, Horse Flies, Deer Flies, Face Flies, House Flies, Horn Flies, Mosquitoes and Gnats: Spray about 3 seconds on each side being careful to spray back, withers and forelegs thoroughly. To protect from Face Flies, spray the face and head, but do not spray into eyes. Repeat treatment when flies are troublesome.

**To kill Screw Worms** infesting beef cattle, horses, and sheep, hold can about 4 to 8 inches from area to be treated with nozzle pointed at of such factors as weather conditions, presence of other materials, or wound. Spray the area until it is completely wet - three to ten seconds are usually enough. The area surrounding the wound should also be sprayed. This product is not recommended for use in deep wounds. Some strains of screw worms may develop resistance to pyrethroids. This will be observed by lack of response to treatment. Where pyrethroid resistance is known or suspected, this product should not be used.

**FOR SWINE:** Do not ship animals for slaughter within 5 days of last treatment.

**To control Blood Sucking Lice:** Apply to the infested areas of the animal using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

**To control Poultry Lice:** Spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist. For control of Bedbugs and Mites in poultry houses, spray

crevices of roost poles, cracks in walls and cracks in nests where the bedbugs and mites hide Spray to the point of dampness but not run off.

**FOR THE CONTROL OF GARDEN INSECTS:** For use on roses dahlias, asters and other ornamentals to kill Japanese Beetles hit by spray, as well as Whiteflies, Aphids exposed Thrips, Spider Mites, Red Mites and Leafminers.

May be used on plants as well as a multi purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, asters, azaleas, begonias, carnations, chrysanthemums, dahlias, delphiniums, dogwood, English Ivy, euonymus, fuchsia, geraniums, cress, grape vine, Kentia palm, laurel, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering Jews and zinnias.

Effective when used as directed against Whiteflies, Aphids, Spider and Red Mites, exposed Thrips, Lace Bugs, Japanese Beetles, Pavement Ants, Clover Mites and Army Worms. When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Spray in one second bursts. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. (32°C). Contact insects directly whenever possible. Retreat if infestation occurs, but not more than once every two weeks.

#### IMPORTANT: READ BEFORE USE

#### STORAGE AND DISPOSAL

**Storage:** Store in a cool, dry area. Do not transport or store below 32°F (0°C.)

**CONTAINER DISPOSAL:** Do not Puncture or Incinerate.

If empty: Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions.

#### FIRST AID

Call a poison control center or a doctor immediately for treatment or advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

**QUESTIONS? Call 1-800-463-9179.**

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes.

Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

**IF INHALED:** Move person to fresh air.

If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.

**IF ON SKIN:** Take off contaminated clothing.

**OR CLOTHING:** Rinse skin immediately with plenty of water for 15-20 minutes.

**Purchaser assumes all responsibility for safety and use not in accordance with directions.**

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following conditions, disclaimer of warranties and limitations of liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Sungro Chemicals, Inc. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** Sungro Chemicals, Inc. MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Sungro Chemicals, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. Sungro Chemicals, Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

**LIMITATIONS OF LIABILITY THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT SUNGRO CHEMICALS, INC. ELECTION, THE REPLACEMENT OF PRODUCT.**

Sungro Chemicals, Inc.  
P.O. Box 24632  
Los Angeles, CA 90024

030703lam