

SHALCO  
**AMINE**  
 SELECTIVE WEED KILLER

**ACCEPTED**  
 OCT 1 1975  
 Under the Federal Insecticide,  
 Fungicide, and Plant Disease Act,  
 as amended, for the pesticide  
 registered under  
 EPA Reg. No. 11511-23

*Comments*

**CAUTIONS:**

May cause skin irritation.  
 Harmful if swallowed.  
 Avoid contact with eyes, skin or clothing.  
 Avoid inhaling spray mist.  
 Do not use on light sandy soils.  
 Do NOT store with foodstuffs.

This product will kill or seriously injure many desirable forms of vegetation. Do not apply directly to flowers, fruits, vegetables, grapes, ornamentals, cotton or other desirable plants. Do not use when there is hazard from drifting mists. (Concentrate sprays are less likely to drift.)

Vapors from this product may injure susceptible plants in the immediate vicinity. Avoid contamination of water used for domestic purposes and irrigation purposes. Excessive amounts of this product in the soil may temporarily inhibit seed germination and plant growth.

Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not store near fertilizers, seeds, insecticides or fungicides.

Avoid contamination of feed and foodstuffs.

**CONTENTS:**

For control of broadleaf weeds in lawns, golf courses, park grounds and similar non-crop turf areas.

<b>ACTIVE INGREDIENT:</b>	
*Dimethylamine salt of 2,4-Dichlorophenoxyacetic acid .....	26.4%
<b>INERT INGREDIENTS:</b> .....	73.6%
<b>TOTAL</b> .....	<b>100.0%</b>

Equivalent to 22.1% 2,4-Dichlorophenoxyacetic acid, or 2 lbs. 2,4-D acid per gallon.

**CAUTION: KEEP OUT OF REACH OF CHILDREN**  
 See Additional Precautions on Side Panels

**CONTROLS THESE TURF WEEDS:**

- |             |                            |
|-------------|----------------------------|
| Buck Horn   | Red Sorrel                 |
| Plantains   | Wild Garlic                |
| Curled Dock | Wild Onion                 |
| Dandelion   | Many Other broadleaf weeds |

**DIRECTIONS FOR USE**

Use 2 to 3 quarts per acre (1½ to 2½ ounces in 1-2 gallons water, per 1,000 sq. ft) in sufficient water for uniform spray coverage. Repeat applications may be necessary. Use lower rate for easy-to-kill species, i.e. dandelion and plantains. Use higher rate for more effective general control of weeds.

Use preferably in the spring or fall when weeds are actively growing. The killing action is more rapid when applied on warm clear sunny days than when weather is cool, cloudy or wet.

**NOTE:** Do not use on newly seeded lawns. Do not mow turf for several days before or after treatment. In some areas Dichondra, clover, bent grasses, carpet buffalo, and St. Augustine grasses may be injured.

Sold By

**SHALCO CHEMICAL CORP.**  
 P. O. BOX 2508 - 2421 LEXINGTON AVE.  
 TOLEDO, OHIO 43606

Do not reuse empty drums. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place away from water supplies. Do not burn.

Spray equipment used in applying this product should be thoroughly cleaned before using for any other purpose. Use repeated flushing with soap and warm water or suitable chemical cleaner. It is best to use a separate sprayer for application of insecticides and fungicides.

**NOTICE**

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

DO NOT Store with Foodstuffs.

EPA Reg. No. 11511-23  
 EPA Est. 11511-OH-1

ACCEPTED WITH COMMENTS